



US007192755B1

(12) **United States Patent**
Mori et al.

(10) **Patent No.:** **US 7,192,755 B1**
(45) **Date of Patent:** **Mar. 20, 2007**

(54) **NICOTIANAMINE SYNTHASE AND GENE ENCODING THE SAME**

(75) Inventors: **Satoshi Mori**, Chiba-ken (JP); **Kyoko Higuchi**, Gunma (JP); **Kazuya Suzuki**, Tokyo (JP); **Naoko Nishizawa**, Tokyo (JP); **Hiroshi Nakanishi**, Tokyo (JP)

(73) Assignee: **Japan Science and Technology Corporation**, Saitama-Ken (JP)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **09/674,337**

(22) PCT Filed: **Apr. 30, 1999**

(86) PCT No.: **PCT/JP99/02305**

§ 371 (c)(1),
(2), (4) Date: **Jul. 26, 2001**

(87) PCT Pub. No.: **WO99/57249**

PCT Pub. Date: **Nov. 11, 1999**

(30) **Foreign Application Priority Data**

Apr. 30, 1998 (JP) 10-137685

(51) **Int. Cl.**

C12N 9/10 (2006.01)
C12N 9/00 (2006.01)
A01H 9/00 (2006.01)
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **435/193**; 435/183; 530/350;
800/295; 800/320; 800/320.2

(58) **Field of Classification Search** 435/183,
435/193; 530/350; 800/295, 320, 320.2

See application file for complete search history.

(56) **References Cited**

FOREIGN PATENT DOCUMENTS

WO 99/60107 11/1999

OTHER PUBLICATIONS

Attwood et al. Which craft is best in bioinformatics? *Comput. Chem.* 2001, vol. 25(4), pp. 329-339.*
Ponting, C.P. Issues in predicting protein function from sequence. *Brief. Bioinform.* Mar. 2001, vol. 2(1), pp. 19-29.*
Accession Q9ZQV8.*
Higuchi et al. *Plant Physiol.* Feb. 1999;119(2):471-80.*
K. Higuchi et al., *Journal of Experimental Botany*, 46(289):1061-1063 (1995).
K. Higuchi et al., *Plant Soil*, 165:173-179 (1994).
K. Higuchi et al., *Plant Soil*, 178:171-177 (1996).
S. Mori, *Soil Science and Plant Nutrition*, 43:975-980 (1997).
H. Ling et al., *Molecular and General Genetics*, 252:87-92 (1996).
K. Higuchi et al., *Journal of Plant Nutrition*, 19(8-9):1235-1239 (1996).
Mori et al., *EMBL Nucleotide Sequence*, Accession No. AB010086 (1999).

Mori et al., *EMBL Nucleotide Sequence*, Accession No. AB011265 (1999).

Mori et al., *EMBL Nucleotide Sequence*, Accession No. AB011264 (1999).

Mori et al., *EMBL Nucleotide Sequence*, Accession No. AB011266 (1999).

Mori et al., *EMBL Nucleotide Sequence*, Accession No. AB011268 (1999).

Mori et al., *EMBL Nucleotide Sequence*, Accession No. AB011269 (1999).

Mori et al., *EMBL Nucleotide Sequence*, Accession No. AB019525 (1999).

Higuchi et al., *Plant Physiology, American Society of Plant Physiologists*, 119:471-479 (1999).

K. Suzuki et al., *Database EMBL Sequence Library*, Accession No. AB021935 (1999).

Plant and Soil, vol. 165, No. 2, issued 1994, Kyoko Higuchi et al.; Purification and characterization of nicotianamine synthase from Fe-deficient barley roots, pp. 173-179.

Plant Physiology, vol. 199, No. 2, issued Feb. 1999, Kyoko Higuchi et al., Cloning of Nicotianamine Synthase Genes, Novel Genes Involved in the Biosynthesis of Phytosiderophores, pp. 417-479.

Database GenBank, Accession No. AB019525, Feb. 11, 1999, Mori S. "*Horedeum vulgare* hvnas7 mRNA for nicotianamine synthase 7, complete cds".

Database GenBank, Accession No. AB021746, Mar. 30, 1999, Mari S. "*Oryza sativa* osnas1 mRNA for nicotianamine synthase 1, complete cds".

Database GenBank, Accession No. AB021934, Feb. 11, 1999, Suzuki K. "*Arabidopsis thaliana* gene for nicotianamine synthase, complete cds."

Frances H. Arnold, "Design by Directed Evolution", *Acc. Chem. Res.*, 31:125-131 (1998).

R.W. Old et al., Fourth Edition, Principles of Gene Manipulation, an Introduction to Genetic Engineering, Blackwell Scientific Publications, Studies in Microbiology vol 2, Chapter 4, pp. 75-98 (1980).

J. Sambrook et al., Molecular Cloning, A Laboratory Manual, Second Edition, Cold Spring Harbor Laboratory Press, "Generation of Deletions and Insertions", pp. 15.3-15.113 (1989).

* cited by examiner

Primary Examiner—Tekchand Saidha

Assistant Examiner—Christian L. Fronda

(74) *Attorney, Agent, or Firm*—Peter F. Corless; Jonathan M. Spaks; Edwards Angell Palmer and Dodge, LLP

(57) **ABSTRACT**

A nicotianamine synthase is isolated and purified. Then the gene of this enzyme is cloned and the base sequence and amino acid sequence thereof are determined. This gene is employed in constructing plants, in particular, grass plants highly tolerant to iron deficiency. A nicotianamine synthase involved in the mugineic acid biosynthesis pathway; the amino acid sequence thereof; a gene encoding the same; a vector containing this gene; cells transformed by the vector; a process for producing nicotianamine by using the same; plants transformed by the gene encoding the nicotianamine synthase; and an antibody against the nicotianamine synthase.

20 Claims, 18 Drawing Sheets

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6

	GCG TTC AGA GGC TTC CAG AGT TCT TCC GGT CAC CAA GAA GCA TTT GAT CAT AAC	54
19	ATG GAT GCC CAG AAC AAG GAG GTC GCT GCT CTG ATC GAG AAG ATC GCC GGT ATC M <u>D A Q N K E V A A L I E K I A G I</u>	108
37	CAG GCC GCC ATC GCC GAG CTG CCG TCG CTG AGC CCG TCC CCC GAG GTC GAC AGG <u>Q A A I A E L P S L S P S P E V D R</u>	162
55	CTC TTC ACC GAC CTC GTC ACG GCC TGC GTC CCG CCG AGC CCC GTC GAC GTG ACG L F T D L V T A C V P P S P V D V T	216
73	AAG CTC AGC CCG GAG CAC CAG AGG ATG CCG GAG GCT CTC ATC CCG TTG TGC TCC K L S P E H Q R M <u>R E A L I R L C S</u>	270
91	GCC GCC GAG GGG AAG CTC GAG GCG CAC TAC GCC GAC CTG CTC GCC ACC TTC GAC A A E G K L E A H Y A D L L A T F D	324
109	AAC CCG CTC GAC CAC CTC GGC CTC TTC CCG TAC TAC AGC AAC TAC GTC AAC CTC N P L D H L G L F P Y Y S N Y V N L	378
127	AGC AGG CTG GAG TAC GAG CTC CTG GCG CCG CAC GTG CCG GGC ATC GCG CCG GCG S R L E Y E L L A R H V P G I A P A	432
145	CGC GTC GCC TTC GTC GGC TCC GGC CCG CTG CCG TTC AGC TCG CTC GTC CTC GCC R V A F V G S G P L P F S S L V L A	486
163	GCG CAC CAC CTG CCC GAG ACC CAG TTC GAC AAC TAC GAC CTG TGC GGC GCG GCC A H H L P E T Q F D N Y D L C G A A	540
181	AAC GAG CCG GCC AGG AAG CTG TTC GGC GCG ACG GCG GAC GGC GTC GGC GCG CGT N E R A R K L F G A T A D G V G A R	594
199	ATG TCG TTC CAC ACG GCG GAC GTC GCC GAC CTC ACC CAG GAG CTC GGC GCC TAC M <u>S F H T A D V A D L T Q E L G A Y</u>	648
217	GAC GTG GTC TTC CTC GCC GCG CTC GTC GGC ATG GCA GCC GAG GAG AAG GCC AAG <u>D V V F L A A L V G M A A E E K A K</u>	702
235	GTG ATT GCC CAC CTG GGC GCG CAC ATG GTG GAG GGG GCG TCC CTG GTC GTG CCG <u>V I A H L G A H M V E G A S L V V R</u>	756
253	AGC GCA CCG CCC CCG GGC TTT CTT TAC CCC ATT GTC GAC CCG GAG GAC ATC AGG S A R P R G F L Y P I V D <u>P E D I R</u>	810
271	CGG GGT GGG TTC GAG GTG CTG GCC GTG CAC CAC CCG GAA GGT GAG GTG ATC AAC <u>R G G F E V L A V H H P E G E V I N</u>	864
289	TCT GTC ATC GTC GCC CGT AAG GCC GTC GAA GCG CAG CTC AGT GGG CCG CAG AAC S V I V A R K A V E A Q L S G C P Q N	918
307	GGA GAC GCG CAC GCA CCG GGC GCG GTG CCG TTG GTC AGC CCG CCA TGC AAC TTC G D A H A R G A V P L V S P P C N F	972
325	TCC ACC AAG ATG GAG GCG AGC GCG CTT GAG AAG AGC GAG GAG CTG ACC GCC AAA S T K M E A S A L E K S E E L T A K	1026
	GAG CTG GCC TTT TGA TTG AAG AGT GCG CGT GGT CAT TCT GTC GCC TGC GAT CGT E L A F *	1080
	GGT AAC TTT CCT ACT CGT GTG TGT TTT GAT GTT TGT GCC TGT AAG AGT TAT GCT	1134
	TCC GGC CTT GTG CTG TTA ATT TAC ACG CGT TAC ATG TAG TAC TTG TAT TTA TAC	1188
	CTG GAA TAA CCG TAT GTA ACA TAA ATA TTA GTG GGA TTT GAA GTG TAA TGC TAA	1242
	ATA ATA AGA AAA CTT GAT GCA GAC ATT CAA AAA AAA AAA AAA AAA AAA AA	

Fig. 7

HvNAS4 MDGGSE - EVDALVQKITGLHAAIAKLPSLSPSPDVDALFTDLVTACVPPSPVDVTKLAP
HvNAS7 MDAQSK - EVDALVQKITGLHAAIAKLPSLSPSPDVDALFTDLVTACVPPSPVDVTKLAP
HvNAS6 MDAQNK - EVDALVQKITGLHAAIAKLPSLSPSPDVDALFTDLVTACVPPSPVDVTKLGS
HvNAS2 MAAQNN - QEVDALVEKITGLHAAIAKLPSLSPSPDVDALFTELVTACVPPSPVDVTKLGP
HvNAS3 MAAQNNKDVAAALVEKITGLHAAIAKLPSLSPSPDVDALFTELVTACVPPSPVDVTKLGP
HvNAS1 MDAQNK - EVAALIEKIAGIAAAIAELPSLSPSPEVDRLFTDLVTACVPPSPVDVTKLSP
HvNAS5 MEAENG - EVAALVEKITGLHAAISKLPALSPSPQVDALFTELVAACVPSSPDVTKLGP
* * * * *
HvNAS4 EAQAMREGLIRLCSEAEGKLEAHYSDMLAAFDNPLDHLGVFPYYSNYINLSKLEYELLAR
HvNAS7 EAQAMREGLIRLCSEAEGKLEAHYSDMLAAFDNPLDHLGVFPYYSNYINLSKLEYELLAR
HvNAS6 EAQEMREGLIRLCSEAEGKLEAHYSDMLAAFDNPLDHLGMFPYYSNYINLSKLEYELLAR
HvNAS2 EAQEMREGLIRLCSEAEGKLEAHYSDMLAAFDKPLDHLGMFPYYSNYINLSKLEYELLAR
HvNAS3 EAQEMREGLIRLCSEAEGKLEAHYSDMLAAFDNPLDHLGIFPYYSNYINLSKLEYELLAR
HvNAS1 EHQRREALIRLCSAAEGKLEAHYADL LATAFDNPLDHLGLFPYYSNYVNL SRLEYELLAR
HvNAS5 EAQEMRQDLIRLCSAAEGLLEAHYSDML TALDSPLDHLGRFPYFDNYVNL SKLEHDL LAG
* * * * *
HvNAS4 YVPGRHRPARVAFIGSGPLPFSSYVLAARHLPDTVFDNYDL CGAANDRATRLFRADKD - V
HvNAS7 YVPGGIAPARVAFIGSGPLPFSSYVLAARHLPDTVFDNYVPVRAANDRATRLFRADKD - V
HvNAS6 YVPGGIARPAVAFIGSGPLPFSSYVLAARHLPDAMFDNYDLCSAANDRASKLFRADKD - V
HvNAS2 YVPGGYRPARVAFIGSGPLPFSSVLAARHLPDTMFDNYDL CGAANDRASKLFRADRD - V
HvNAS3 YVRR - HRPARVAFIGSGPLPFSSVLAARHLPDTMFDNYDL CGAANDRASKLFRADTD - V
HvNAS1 HVPG - IAPARVAFIGSGPLPFSSVLAARHLPETQFDNYDL CGAANERARKLFGATADGV
HvNAS5 HVAA - - - PARVAFIGSGPLPFSSVLAARHLPDTRFDNYDRCSVANGRAMKLVGAADGEGV
* * * * *
HvNAS4 GARMSFHTADVADLTDELATYDVVFLAALVGMAAEDKAKVIAHLGAHMADGAALV - - ARH
HvNAS7 GARMSFHTADVADLTDELATYDVVFLAALVGMAAEDKGGDPHLGAHMADGAALVR - SAH
HvNAS6 GARMSFHTADVADLTRELAAYDVVFLAALVGMAAEDKAKVIPHLAGHMADGAALVV - RSA
HvNAS2 GARMSFHTADVADLAGELAKYDVVFLAALVGMAAEDKAKVIAHLGAHMADGAALVVRSAH
HvNAS3 GARMSFHTADVADLASELAKYDVVFLAALVGMAAEDKAKVIAHLGAHMADGAALVVRSAH
HvNAS1 GARMSFHTADVADLTQELGAYDVVFLAALVGMAAEEKAKVIAHLGAHMVEGASLVV - RSA
HvNAS5 RSRMAFHTAEVDTLTAELGAYDVVFLAALVGMTSKEKADAIAHLGKHMADGAVLREALH
* * * * *
HvNAS4 GARGFLYPIDPQDIGRGGFEVLAVCHPD - DDVNSVIIAQSNDVHEYGLGSGR - - GGR
HvNAS7 GARGFLYPIDPQDIGRGGFEVLAVCHPD - DDVNSVIIAQSNDVHEYGLGSGR - - GGR
HvNAS6 QARGFLYPIDPQDIGRGGFEVLAVCHPD - DDVNSVIIAHSKDVHANERPNGR - - GGQ
HvNAS2 GARGFLYPIDPQDIGRGGFEVLAVCHPD - DDVNSVIIAQSNDVHADGLGSGRGAGGQ
HvNAS3 GARGFLYPIDPQDIGRGGFEVLAVCHPD - DDVNSVIIAQSKEVHADGLGSGRAGGRQ
HvNAS1 RPRGFLYPIDPEDIIRGGFEVLAVHP - - GEVINSVIVARKAVEAQLSGPQMGD - - - - A
HvNAS5 GARAFLYPVVELDDVGRGGFQVLAVHPAGDEVFNSFTVARKVKMSA - - - - -
* * * * *
HvNAS4 YARGTVPVVSPPCRFG - EMVADVTD - - KREEFANAEEVAF
HvNAS7 YARG - TVPVVSPPCRFG - EMVADVTD - - KREEFANAEEVAF
HvNAS6 YRGA - - VPVVSPPCRFG - EMVADVTH - - KREEFANAEEVAF
HvNAS2 YARG - TVPVVSPPCRFG - EMVADVTDNHRDEFANAEEVAF
HvNAS3 YARG - TVPVVSPPCRFG - EMVADVTDNHRDEFANAEEVAF
HvNAS1 HARG - AVPLVSPPCNFSTKMEASALE - - KSEELTAKELAF
* * * * *

Fig. 8




Fig. 9




Fig. 10




Fig. 11




Fig. 12




Fig. 13




Fig. 14




Fig. 15




Fig. 16




Fig. 17





Fig. 18



1

NICOTIANAMINE SYNTHASE AND GENE ENCODING THE SAME

TECHNICAL FIELD

The present invention relates to a nicotianamine synthase involved in the mugineic acid biosynthetic pathway, the amino acid sequence thereof, a gene encoding the same, a vector, a process for producing nicotianamine by using the same, plants transformed by the gene encoding the nicotianamine synthase, and an antibody against the nicotianamine synthase.

BACKGROUND ART

Graminaceous plants that absorb by chelating the insoluble state Fe(III) in soil using mugineic acid and adopt so called the Strategy-II mechanism of Fe acquisition secrete Fe chelators (phytosiderophores) from their roots to solubilize sparingly soluble Fe in the rhizosphere (Roemheld, 1987). The amount of the secreted phytosiderophores increases under Fe-deficiency stress. The mugineic acid family is the only examples of phytosiderophores known so far (Takagi, 1976). Tolerance to Fe deficiency in graminaceous plants is thought to depend on a quantity of mugineic acid family secreted by plants (Takagi et al. 1984, Roemheld and Marschner 1986, Marschner et al. 1987, Mori et al. 1987, Kawai et al. 1988, Mori et al. 1988, Mihashi and Mori 1989, and Shingh et al. 1993).

The biosynthetic pathway of mugineic acid in plants is shown in FIG. 1. S-adenosylmethionine is synthesized from methionine by S-adenosylmethionine synthase. Subsequently, three molecules of S-adenosylmethionine are combined to form one molecule of nicotianamine by nicotianamine synthase. The generated nicotianamine is then converted to 3"-keto acid by nicotianamine aminotransferase, and 2'-deoxymugineic acid is synthesized by the subsequent action of a reductase. A further series of hydroxylation steps produces the other mugineic acid derivatives including mugineic acid from the deoxymugineic acid (Mori and Nishizawa 1987, Shojima et al. 1989, Shojima et al. 1990 and Ma and Nomoto 1993).

A compound in FIG. 1, a compound in the lower right, wherein R₁ and R₂ are hydrogen and R₃ is hydroxyl, is mugineic acid. A compound wherein R₁ is hydrogen and R₂ and R₃ are hydroxyl, is 3-hydroxymugineic acid. Also a compound wherein R₂ is hydrogen and R₁ and R₃ are hydroxyl, is 3-epihydroxymugineic acid.

Three S-adenosylmethionine synthase genes were isolated from barley roots, but these genes were not induced by Fe deficiency (Takizawa et al. 1996). A gene lds3, which is obtained from the barley by differential screening, is suspected to be a gene, which converts deoxymugineic acid to mugineic acid by hydroxylation and is strongly induced by Fe-deficiency (Nakanishi et al. 1993). Further, nicotianamine aminotransferase was purified and isolated from Fe-deficient barley roots, and two nicotianamine aminotransferase genes, Naat-A and Naat-B, were isolated (Takahashi et al. 1997). Naat-A expression was induced in Fe-deficient roots.

The synthesis of nicotianamine from S-adenosylmethionine is similar to polyamine synthesis from decarboxy-S-adenosylmethionine. In contrast to polyamine synthase, however, nicotianamine synthase catalyzes the combination of three S-adenosylmethionine molecules and the azetidinium ring formation at the same time (FIG. 1). Such the nicotianamine synthase is a novel type of enzyme. Previously, we

2

reported the partial purification of nicotianamine synthase from the roots of Fe-deficient barley and expression pattern of the activity (Higuchi et al. 1994, Higuchi et al. 1995, Kanazawa et al. 1995, Higuchi et al. 1996a and Higuchi et al. 1996b). Since nicotianamine synthase is easily decomposed during extraction and purification, it has been difficult to purify sufficient quantities for amino acid sequencing.

The present invention has an object to provide a plant, especially graminaceous plant, highly tolerant to Fe-deficiency, as a result of isolating and purifying a nicotianamine synthase, being cloned the gene of this enzyme, determining the base sequence and amino acid sequence thereof, and using said enzyme.

DISCLOSURE OF INVENTION

The present invention relates to a nicotianamine synthase shown in SEQ ID NO: 1 comprising amino acid sequence shown in SEQ ID NO: 1, or amino acid sequence having deletion in a part thereof, being substituted by the other amino acids or being added with the other amino acids.

The present invention relates to the gene encoding said amino acid sequence of nicotianamine synthase.

The present invention also relates to a vector comprising said gene, and a transformant transformed by the said vector.

The present invention relates to a process for production of nicotianamine using the said transformant.

The present invention further relates to plants, especially graminaceous plants, to which said gene is introduced, and fruits obtained by growing said plants.

The present invention relates to a process for extraction of said nicotianamine synthase in the presence of thiol protease inhibitor, preferably E-64.

Further, the present invention relates to an antibody against said nicotianamine synthase.

BRIEF DESCRIPTION OF DRAWING

FIG. 1 shows the biosynthetic pathway of mugineic acid family.

FIG. 2 shows a comparison of nicotianamine synthase purification from Fe-dependent and control barley roots.

FIG. 3 shows a preparative SDS-PAGE (sodium dodecyl sulfate-polyacrylamide gel electrophoresis, hereinafter designates as SDS-PAGE) around 30-35 kDa. The horizontal bar indicates relative enzyme activity detected from the gels.

FIG. 4 shows elution pattern of nicotianamine synthase activity from the gel-filtration column.

The large closed circles (●) indicates enzyme activity.

FIG. 5 shows a comparison with a six partial amino acid sequence determined by nicotianamine synthase originated from barley and similar sequence of graminaceous plants obtained by computer search of the database. Identical amino acid residue is shown in ":" (SEQ ID NOS 30-37, respectively in order of appearance).

FIG. 6 shows full length of HvNAS1 cDNA (SEQ ID NO: 2) and amino acid sequence (SEQ ID NO: 1) deduced therefrom. The underlined sequences indicate the identical partial amino acid sequences of fragments in the above FIG. 5. Numbers of the nucleotide sequence are indicated to the right of each row. Amino acid numbers are indicated on the left of each row.

FIG. 7 shows comparison of the deduced amino acid sequences of the above 7 cDNA obtained from barley (SEQ ID NOS 7, 13, 11, 3, 5, 1, and 9, respectively in order of appearance). Asterisks "*" indicates identical amino acid

3

residues in all sequences used to generate consensus sequences SEQ ID NOS: 23–29.

FIG. 8 shows results of thin layer chromatographic (TLC) analysis of nicotianamine synthase activity obtained from *E. coli* crude extract expressing a fused protein of maltose binding protein—HvNAS1.

FIG. 9 shows Northern—hybridization analysis of HvNAS1 as a probe.

FIG. 10 shows Southern—hybridization analysis of HvNAS1 as a probe.

FIG. 11 shows Western-blot analysis of crude enzyme used for detection of nicotianamine synthase activity.

FIG. 12 shows Western-blot analysis of total protein extracted by trichloroacetic acid/acetone.

FIG. 13 shows comparison of nicotianamine synthase purification from Fe-deficient barley and control barley after DEAE-Sepharose FF.

FIG. 14 shows comparison of nicotianamine synthase purification from Fe-deficient barley and control barley after Ether Toyopearl 650M.

FIG. 15 shows results of thin layer chromatographic (TLC) analysis of nicotianamine synthase activity obtained from *E. coli* crude extract expressing a fused protein of maltose binding protein—OsNAS1.

FIG. 16 shows Northern—hybridization analysis of OsNAS1 as a probe.

FIG. 17 shows results of thin layer chromatographic (TLC) analysis of nicotianamine synthase activity obtained from *E. coli* crude extract expressing a fused proteins of maltose binding protein—AtNAS1, AtNAS2 or AtNAS3.

FIG. 18 shows results of RT-PCR of total RNA extracted from the aboveground parts and roots of *Arabidopsis thaliana*. Right group indicates positive control.

BEST MODE FOR CARRYING OUT THE INVENTION

We have tried to isolate nicotianamine synthase (Higuchi et al. *Plant & Soil*, Vol. 165, p. 173–179, 1994), and since nicotianamine synthase was easily decomposed and was difficult to isolate and purify, we were unable to obtain sufficient amounts of protein to determine its partial amino acid sequence. Subsequently, it was found that a thiol protease inhibitor E-64 (hereinafter designates as E-64) was very effective in suppressing degradation of nicotianamine synthase (Higuchi et al. *Plant & Soil*, Vol. 178, p. 171–177, 1996 a).

In the present invention, as a result that frozen roots were crushed to a fine powder in liquid N₂ and then rapidly homogenized with buffer containing 0.1 mM thiol protease inhibitor E-64, nicotianamine synthase protein could be isolated and its gene could also be isolated.

Further, the enzyme of the present invention recovered its activity by removal of SDS after SDS-PAGE treatment, but the rate of recovery was very low (Higuchi et al. *Plant & Soil*, Vol. 165, p. 173–179, 1994). Consequently, degree of purification should be increased up before treatment of SDS-PAGE. Then the column chromatography procedures were further improved.

We have also found that the enzyme of the present invention is relatively hydrophobic and a buffer containing a mild surface active agent CHAPS increased the rate of recovery. Several ion-exchange chromatography carriers were tested, and DEAE-Sepharose FF and DEAE Sephacel were found to be the most effective. In addition to TSK gel

4

Butyl Toyopearl, another hydrophobic chromatography carrier, TSK gel Ether Toyopearl 650M, effectively removed impurities of the 30–35 kDa.

The enzyme of the present invention has been reported that it was the peptide of 30–35 kDa, the activity of which was recovered by removing SDS after SDS-PAGE treatment, and the activity was detected as a broad molecular weight range of 30–35 kDa (refer to FIG. 3). FIG. 3 shows a result of preparative SDS-PAGE in the fractions showing enzyme activity. SDS-PAGE was carried out using 11% acrylamide slab gels. A portion of the gel was stained with Coomassie brilliant blue and the rest of the gel was stained with Cu. The gel containing proteins between 30–35 kDa in size was cut into seven fragments (indicated by the short lines). The thick bars in FIG. 3 indicate relative enzymatic activities detected from each gel fragment.

In order to identify nicotianamine synthase peptide from the proteins having these molecular weights, the peptides, which were contained in the nicotianamine synthase fractions, purified from Fe-deficient and control barley roots were compared using SDS-PAGE. From each barley root 200 g, the present enzyme was purified according to the method described in example 3 hereinbelow.

The enzyme activity of the control was a quarter of the Fe-deficient roots.

The peptide composition of the active enzyme fraction from each purification step of the present enzyme was analyzed and compared by SDS-PAGE, and results are shown in FIG. 2, FIG. 13 and FIG. 14. FIG. 2, FIG. 13 and FIG. 14 show comparison with the active fraction from the purification step of Fe-deficient barley roots 200 g [in the figure, shown with (-)], and the active fraction from the purification step of the control barley roots 200 g [in the figure, shown with (+)]. SDS-PAGE was carried out using 12.5% acrylamide slab gels (Laemmli, *Nature* Vol. 227, p. 680–685, 1970). Gels were stained with Coomassie brilliant blue. FIG. 2 shows a step before DEAE-Sepharose. The upper row shows enzyme from Fe-deficient barley roots and the lower row shows enzyme from control roots. In each lane, lanes 1, crude extract, 200 μg of protein; lanes 2, after Butyl Toyopearl 650M, 100 μg of protein; lanes 3, after hydroxyapatite, 20 μg of protein; and lanes 4, after Butyl Toyopearl 650M, 15 μg of protein, are shown.

FIG. 13 shows after DEAE-Sepharose FF, each lane, 25 μg of protein. FIG. 14 shows after Ether Toyopearl 650M; in which left shows inactive fraction, and right shows active fraction, and 1/25 of each fraction is electrophoresed.

As a result, almost no difference was observed in both Fe-deficient and control roots before DEAE-Sepharose step (refer to FIG. 2). After the DEAE-Sepharose step it became clear that the 30- and 31-kDa peptides were induced by Fe-deficiency (refer to FIG. 13). After the Ether Toyopearl step, the 31 kDa peptide was eliminated from the active fraction. The 32 and 33 kDa peptides were found to be newly induced by Fe-deficiency (refer to FIG. 14). Activities were detected from the 32 and 33 kDa peptides, but no activity was detected from 30 kDa peptide (refer to FIG. 3).

Molecular weight of the enzyme of the present invention was determined by gel-filtration.

Estimated molecular weight of nicotianamine synthase by gel-filtration was reported to be 40,000–50,000 (Higuchi et al. *Plant & Soil*, Vol. 165, p. 173–179, 1994). But this did not correspond with the value estimated by SDS-PAGE.

In the present study, the buffer containing CHAPS effectively increased the resolution and molecular weight of the present enzyme was estimated to be 35,000 (refer to FIG. 4). This corresponds well to the value estimated by SDS-PAGE.

5

FIG. 4 shows elution pattern of nicotianamine synthase from the gel-filtration column. The black circles (●) indicate the enzyme activity and the solid line indicates absorption at 280 nm. The active fraction after hydroxyapatite chromatography was applied to a Sephacryl S300HR (Pharmacia) column (1.5 cm×71 cm, 125 ml), equilibrated with developing buffer (50 mM Tris, 1 mM EDTA, 0.1 M KCl, 0.05% CHAPS, 0.1 mM p-APMSF and 3 mM DTT, pH 8.0). Molecular weight markers used were thyroglobulin (Mr 670,000), γ -globulin (Mr 158,000), ovalbumin (Mr 44,000), and myoglobin (Mr 17,000). The linear flow was 10 cm/hour.

Partial amino acid sequence was determined from purified nicotianamine synthase.

The above explained 30 kDa, 32 kDa and 33 kDa peptides were purified from 1 kg of Fe-deficient barley roots by using a method in example 3 hereinbelow. These were partially degraded using a method in example 4 hereinbelow. Although 32- and 33-kDa peptides could not be completely separated from each other, these might have similar sequence or 32 kDa peptide was presumed to be the degradation product of 33 kDa peptide, and both of them were degraded in together.

The determined partial amino acid sequences indicated that these peptides were very similar in each other (FIG. 5). Further, since the molecular weights of the 33 kDa and 32 kDa (1) fragments had almost unchanged molecular weight as compared with before degradation, this sequence might be N-terminal region of the present enzyme. As a result of computer search of the database, a gene of unknown function having very similar sequence to these sequences was found to exist in *Oryza sativa* and *Alahidopsis thaliana*. Especially, EST-cDNA clones D23792 and D24790 of *Oryza sativa* were very similar with 80.0% identity in a 33-amino acid overlap in the former and 68.4% identity in a 19-amino acid overlap in the latter (FIG. 5).

FIG. 5 shows a comparison with a six partial amino acid sequence determined by nicotianamine synthase originated from barley and similar sequence of graminaceous plants obtained by computer search of the database. Identical amino acid residue is shown in “:”. The part of nucleotide sequences indicated by the arrows was applied for the sequences of primer used in PCR.

Cloning and nucleotide sequences of cDNA clones encoding nicotianamine synthase were performed and determined.

PCR amplification of total cDNA prepared from Fe-deficient barley roots using degenerate primers designed from the partial amino acid sequence obtained from the method explained hereinbefore was performed, but the objective DNA could not amplified. Then the primers having single nucleotide sequence (shown by arrows in FIG. 5) from sequences of *Oryza sativa*, D23792 and D24790, were synthesized and PCR amplification was performed. The 205 bp fragment was amplified by PCR using NF and NR primers and the 274 bp fragment was amplified by PCR using 1F and 1R primers, and these contained the objective sequences. A cDNA library prepared using poly (A)⁺ RNA from Fe-deficient barley roots was screened and 19 positive clones using the 205 bp fragment probe and 88 positive clones using the 274 fragment bp probe were obtained.

Among the thus obtained clones, the clone designated as HvNAS1, contained a translated region of 985 bp and amino acid sequence deduced therefrom was 328 amino acids residue, with deduced molecular weight of 35,144. This corresponded well with the value estimated by SDS-PAGE

6

and gel-filtration. The partial amino acid sequences of the 32 kDa and 33 kDa peptides were included totally in HvNAS1 (FIG. 6).

FIG. 6 shows full length of HvNAS1 cDNA and amino acid sequence deduced therefrom. The underlined sequences indicate the identical partial amino acid sequences of fragments in the above FIG. 5. Numbers of the nucleotide sequence are indicated to the right of each row. Amino acid numbers are indicated on the left of each row.

The predicted pI of 5.2 matched the value estimated by native isoelectric focusing electrophoresis well. The six clones having very similar sequence other than HvNAS1, i.e. HvNAS2, HvNAS3, HvNAS4, HvNAS5, HvNAS6 and HvNAS7, were also obtained (Table 1, FIG. 7).

FIG. 7 shows comparison of the deduced amino acid sequences of the above 7 cDNA obtained from barley. Asterisks “*” indicates identical amino acid residues in all sequences.

The nucleotide sequences of these clones are shown in SEQ ID NO: 2 (HvNAS1), SEQ ID NO: 4 (HvNAS2), SEQ ID NO: 6 (HvNAS3), SEQ ID NO: 8 (HvNAS4), SEQ ID NO: 10 (HvNAS5), SEQ ID NO: 12 (HvNAS6) and SEQ ID NO: 14 (HvNAS7), respectively. The amino acid sequences of these amino acid sequences are shown in SEQ ID NO: 1 (HvNAS1), SEQ ID NO: 3 (HvNAS2), SEQ ID NO: 5 (HvNAS3), SEQ ID NO: 7 (HvNAS4), SEQ ID NO: 9 (HvNAS5), SEQ ID NO: 11 (HvNAS6) and SEQ ID NO: 13 (HvNAS7), respectively.

TABLE 1

Properties of nas clones						
Clone	Number of Amino Acid Residues	Molecular Weight	pI	Identity to nas 1 (%)	Identity to nas 2 (%)	Identity to nas 4 (%)
HvNAS 1	328	35144	5.20	—	—	—
HvNAS 2	336	35839	5.07	72	—	—
HvNAS 3	336	36013	5.47	72	95	—
HvNAS 4	330	35396	4.91	73	89	—
HvNAS 5	283	30148	5.22	61	61	59
HvNAS 6	329	35350	5.07	74	89	88
HvNAS 7	330	35244	4.98	70	86	91

The partial amino acid sequences determined from the 30 kDa peptide were all included in HvNAS5. The 5'- and 3'-non-translated regions of these clones were not similar with each other.

D23792 and D24790 similar to nicotianamine synthase of *Oryza sativa* were found with about 80% identity to HvNAS1. AC003114 and AB005245 of *Arbidopsis thaliana* were found with about 45% identity to HvNAS1.

The obtained HvNAS1 protein was expressed in *E. coli*.

The PCR amplification of HvNAS1 ORF was cloned with vector pMAL-c2 to express HvNAS1 fused with C-terminal of maltose binding protein. The expression of fused protein is strongly induced by IPTG.

The crude extract was obtained from the transformed *E. coli*, and nicotianamine synthase activity was assayed in the state of the fused protein. The crude extract from the strain transformed with only the vector could not be detected the activity, whereas in case of inserted with HvNAS1 ORF, the activity was detected. Result is shown in FIG. 8.

FIG. 8 shows results of thin layer chromatographic (TLC) analysis of nicotianamine synthase obtained from *E. coli* crude extract expressing a fused protein of maltose binding protein—HvNAS1. In FIG. 8, lane 1: a standard nicotianamine synthase; lane 2: *E. coli* expressing maltose binding

protein (SAM); and lane 3: *E. coli* expressing maltose binding protein—HvNAS1 fused protein.

Northern hybridization analysis conducted by the method described in example 7 hereinbelow indicated that this gene was strongly induced in Fe-deficient roots (FIG. 9). This coincides with expression pattern of the present enzyme activity (Higuchi et al. 1994). FIG. 9 shows a result of Northern hybridization analysis using HvNAS1 as a probe. Total RNA was extracted from after one week of Fe-deficient treatment and control barley leaves and roots, and in each lane, 5 µg of RNA were electrophoresed.

Southern hybridization analysis of the barley genome DNA was performed according to the method described in example 8 hereinafter mentioned. Cutting of DNA with BamHI, EcoRI or HindIII produced plurality of fragments, however none of the clones obtained at present could be digested by BamHI and EcoRI, consequently nicotianamine synthase gene might exist with multiple copies in genomes of barley and rice (FIG. 10).

FIG. 10 shows Southern—hybridization analysis of HvNAS1 as a probe. Genomic DNAs from barley and rice were digested with BamHI (lanes B), EcoRI (lanes R) and HindIII (lanes H) and 10 µg thereof were electrophoresed in each lane.

Further, using antigen prepared by the method described in example 9 hereinbelow, Western-blot analysis was performed according to the method described in example 10. It was found that the present enzyme protein was rapidly decomposed during the operation in the crude extract prepared for detecting the present enzyme activity (FIG. 11). The staining patterns coincided with the fact that the present enzyme activity was detected on the broad range between 30–35 kDa after SDS-PAGE (refer to FIG. 3).

FIG. 11 shows Western-blot analysis of crude enzyme used for detection of activity. SDS-PAGE was performed using 12.5% acrylamide slab gel. Protein 100 µg was electrophoresed.

The crude extract obtained from denatured protein according to the method described in example 10 hereinbelow was detected as almost single band with 35–36 kDa (FIG. 12). This value coincided with the deduced value from the amino acid sequence.

FIG. 12 shows Western-blot analysis of total protein extracted by trichloroacetic acid/acetone. SDS-PAGE was performed using 12.5% acrylamide slab gel. Protein 100 µg was electrophoresed. Proteins 200 µg extracted from roots and proteins 500 µg extracted from leaves were electrophoresed.

Western-blot analysis after 2-dimension electrophoresis reveals to detect several spots. This coincided with the fact of obtaining plurality of nicotianamine synthase gene. All spots were induced in Fe-deficient roots.

As a result that cDNA library from Fe-deficient rice roots poly (A)⁺ RNA was screened using probes prepared by cutting HvNAS1 with restriction enzymes ApaI and XhoI, 20 clones were obtained. These clones were divided into 3 types of clones according to their sequences, and among them, only one type contains ORF full length, which was designated as OsNAS1. Nucleotide sequence of OsNAS1 is shown in SEQ ID NO: 16 and amino acid sequence is shown in SEQ ID NO: 15.

PCR amplification of OsNAS1 ORF was cloned with a vector pMAL-c2 to express a form fused with maltose binding protein C-terminal. The fused protein is strongly induced its expression by IPTG.

Crude extract from the transformed *E. coli* with the fused protein was obtained and nicotianamine synthase activity

was assayed in the state of the fused protein. The same activity with HvNAS1 was detected. Result is shown in FIG. 15. FIG. 15 shows results of thin layer chromatographic (TLC) analysis of nicotianamine synthase obtained from *E. coli* crude extract expressing a fused protein of maltose binding protein—OsNAS1. In FIG. 15, lane 1: a standard nicotianamine (NA); lane 2: an extract from *E. coli* expressing maltose binding protein—OsNAS1 fused protein; and lane 3: an extract from *E. coli* expressing maltose binding protein—HvNAS1 fused protein.

Northern hybridization analysis conducted by the method described in example 7 hereinbelow indicated that in contrast to barley, the expression was induced in rice by Fe-deficient treatment not only in roots but also in leaves (FIG. 16). FIG. 16 shows a result of Northern hybridization analysis using OsNAS1 ORF as a probe. Total RNA was extracted from after two weeks of Fe-deficient treatment and control rice leaves and roots, and in each lane, 5 µg of RNA were electrophoresed.

Nucleotide sequence of *Arabidopsis thaliana* similar to HvNAS1 obtained by computer search of the database was used as a primer. PCR amplification for genome DNA of *Arabidopsis thaliana* resulted to obtain three nicotianamine synthase genes. These were designated as AtNAS1, AtNAS2 and AtNAS3.

Nucleotide sequence of these genes are shown in SEQ ID NO: 18 (AtNAS1), SEQ ID NO: 20 (AtNAS2) and SEQ ID NO: 22 (AtNAS3). These amino acid sequences are shown in SEQ ID NO: 17 (AtNAS1), SEQ ID NO: 19 (AtNAS2) and SEQ ID NO: 21 (AtNAS3).

AtNAS1, AtNAS2 and AtNAS3 ORF were amplified with PCR and were cloned with a vector pMAL-c2. Each of them was tried to be expressed in the form of fusing with maltose binding protein C-terminal. The expression of the fused protein was strongly induced by IPTG.

Crude extract from the transformed *E. coli* with the fused protein was obtained and nicotianamine synthase activity was assayed in the state of the fused protein. The activity was detected. Result is shown in FIG. 17. FIG. 17 shows results of TLC analysis of nicotianamine synthase activity obtained from *E. coli* crude extract expressing a fused protein of maltose binding protein—AtNAS. In FIG. 17, lanes 1: a standard nicotianamine (NA) and S-adenosylmethionine; lanes 2: an extract from *E. coli* expressing only maltose binding protein; lanes 3: an extract from *E. coli* expressing maltose binding protein—AtNAS1 fused protein; lanes 4: an extract from expressing maltose binding protein—AtNAS2 fused protein; and lanes 5: an extract from *E. coli* expressing maltose binding protein—AtNAS3 fused protein.

RT-PCR was conducted according to the method described in example 11 hereinbelow. It was found that AtNAS1 was expressed in the roots and the aboveground parts of *Arabidopsis thaliana*, whereas AtNAS2 was expressed neither in the roots nor in the aboveground parts, and AtNAS3 was expressed only in the roots (FIG. 18). In FIG. 18, lane M shows molecular weight marker. Gene expression was conducted in the aboveground parts, roots and positive controls. In the figure, lanes C: AtNAS1 and AtNAS2 ORF full length were amplified; lanes 1: AtNAS1 specific amplification fragments; lanes 2: AtNAS2 specific amplification fragments; and lanes 3: AtNAS3 specific amplification fragments.

The amount of secreted mugineic acid is reported increased up to 20 mg mugineic acid/g roots dry weight/day (Takagi, 1993). Crude nicotianamine synthase activity detected by the present inventors was sufficient to fulfill it.

Since the present enzyme proteins exist in more than several types and 30 kDa peptide without activity exists, it can be speculated that as a result of aggregation of these peptides, the constructed structure, which is preferable for binding with 3 molecules of S-adenosylmethionine, reveals maximum activity. The molecular weight estimated by gel-filtration was 35,000 (FIG. 4).

Increase in activity by re-aggregation of subunits has not been observed at present. Since the fused protein with maltose binding protein and subunits showed its activity, we have at present an idea that the present enzyme might be a monomer. However, the possibility that large activity can be revealed by constructing multimer, can not completely be denied.

The reaction mechanism synthesizing nicotianamine from S-adenosylmethionine may be similar to methyl transfer reaction using S-adenosylmethionine as a methyl donor, and a reaction synthesizing spermidine and spermine from decarboxylated S-adenosylmethionine. The common catalytic domain of these enzymes has been discussed in relation to equivalent amino acids configuration occupying similar positions in higher-order structures (Hashimoto et al. 1998 and Schluckebier et al. 1995).

In future, catalytic domain may be elucidated as the results of comparison with nicotianamine synthase from other plant species or X-ray crystallography.

Induction of nicotianamine synthase activity by Fe-deficiency is a specific phenomenon in graminaceous plants, and is essential for mass production of mugineic acid family. *Oryza sativa* is a plant, in which secretion of mugineic acid family is the least among major graminaceous plants, consequently it is very weak for Fe-deficiency in calcareous soil.

Consequently, as a result of creating transformant *Oryza sativa* having tolerance to Fe-deficiency by introducing nicotianamine synthase gene of the present invention into the graminaceous plants, especially *Oryza sativa*, and expressing large amount at the Fe-deficiency, cultivation of rice in the calcareous soil can be possible.

Heretofore, in the graminaceous plants, nicotianamine has been thought to have only a role as a precursor for synthesis of mugineic acid family. However, since the present invention has elucidated that nicotianamine synthase gene constituted the multiple gene family, it may play other important roles in the graminaceous plants.

In plants, which lack the ability to secrete mugineic acid family, except for graminaceous plants, it has been proposed that nicotianamine plays a key role as an endogenous chelator of divalent metal cations, such as Fe²⁺, Cu²⁺, Zn²⁺ and Mn²⁺, and that it contributes to the homeostasis of those metals (Stephan et al. 1994). Consequently, it may play the same role in the graminaceous plants.

Nicotianamine synthase activity is not induced in dicots, and expression of gene of the present invention may not be induced by Fe-deficiency. We have cloned nicotianamine synthase genes of *Arabidopsis thaliana*. Composition of promoter regions in these genes can elucidate the mechanism of gene expression caused by Fe-deficiency, and the gene of the present invention may play important function not only in the graminaceous plants but also in the dicots.

SEQ ID NO: 1 shows amino acid sequence of nicotianamine synthase of the present invention.

The present invention includes nicotianamine synthase having amino acid sequence shown in SEQ ID NO: 1. However, the present invention is not limited within the above nicotianamine synthase. The nicotianamine synthase of the present invention includes, unless it loses nicotian-

amine synthase activity, the peptides, in which a part of the amino acid sequence of said peptide is deleted, preferably 50% or less, more preferably 30% or less, or more further preferably 10% or less in the total amino acids, or is substituted by other amino acids, or to which other amino acids are further added, or in which these deletion, substitution and addition may be combined.

Nucleotide sequence coding nicotianamine synthase of the present invention is shown in SEQ ID NO: 2.

The present invention also includes not only a gene coding nicotianamine synthase shown in SEQ ID NO: 2 but also genes coding nicotianamine synthase mentioned hereinabove.

The vector of the present invention introducing the above gene is not specifically limited, and various vectors can be introduced. Preferable vector is the expression vector.

Various cells can be transformed conventionally by using recombinant vector of the present invention. Mass production of nicotianamide can be performed by using the thus obtained transformant. These methods are well known in the person skilled in the art.

Examples of hosts for introducing the gene of the present invention are bacteria, yeasts and cells. Preferable host is plants, especially the graminaceous plant.

Method for introducing gene is not limited. It can be made by using vector or can be directly introduce in genome.

Antibody of the present invention against nicotianamine synthase can be prepared conventionally by using nicotianamine synthase of the present invention. Antibody can be a polyclonal antibody or, if necessary, monoclonal antibody.

Further, a selective breeding of plants, preferably graminaceous plants, can be made by using gene of the present invention. Especially, the gene of the present invention can be applied for improvement of varieties, which can grow even in Fe-deficient soil.

EXAMPLES

The following examples illustrate the present invention, but are not construed as limiting the present invention.

Example 1

Preparation of Plant Material

Seeds of barley (*Hordeum vulgare* L. cv Ehimehadakamugi No. 1) were germinated on wet filter paper and transferred into the standard hydroponic culture solution (Mori and Nishizawa, 1987) in a glass house at natural temperature under natural light. The pH of the hydroponic culture solution was adjusted at 5.5 by 0.5 N HCl everyday. When the third leaves developed, the plants were transferred to the hydroponic culture solution without containing Fe. The pH was maintained at 7.0 by 0.5 N NaOH everyday. The control plants were also cultured in the standard culture solution continuously. The culture solution was renewed once in every week. Two weeks after starting Fe-deficient treatment, when severe iron chlorosis significantly appeared on the 4th and 5th leaves, roots were harvested and frozen in liquid N₂ and stored at -80° C. until use.

Example 2

Assay of Nicotianamine Synthase Activity

Modified assay method reported previously by the present inventors (Higuchi et al. 1996a) was used. Enzyme solutions

11

were equilibrated with reaction buffer [50 mM Tris, 1 mM EDTA, 3 mM dithiothreitol (hereinafter designates as DTT), 10 μ M (p-amidinophenyl) methanesulfonyl fluoride (hereinafter designates as p-APMSF) and 10 μ M trans-epoxysuccinyl-leucylamido-(4-guanidino) butane (hereinafter designates as E-64), pH 8.7]. Buffer exchange was performed by using ultrafiltration unit, Ultrafree C3LGC NMWL10000 (Millipore Co.). S-adenosylmethionine labeled with 14 C in carboxyl group (Amersham Inc.) was added to the enzyme solution at the final concentration of 20 μ M and kept at 25° C. for 15 minutes. The reaction products were separated by thin layer chromatography on silica gel LK6 (Whatman Inc.) using developer (phenol:butanol:formic acid:water=12:3:2:3). Radioactivity of the reaction products was detected by image Analyzer BAS-2000 (Fuji Film Co.). The protein content was assayed by Bradford method using Protein Assay Kit (Bio Rad Inc.).

Example 3

Purification of Nicotianamine Synthase

The following operations were performed at 4° C. and E-64 was added to fractions containing nicotianamine synthase at the final concentration of 10 μ M.

The frozen roots were crushed into a fine powder in liquid N₂ and homogenized in a household juicer with 200 ml of extraction buffer [0.2 M Tris, 10 mM EDTA, 5% (v/v) glycerol, 10 mM DTT, 0.1 mM E-64, 0.1 mM p-APMSF and 5% (w/v) insoluble polyvinylpyrrolidone (PVP), pH 8.0] per 100 g of roots. The homogenate was centrifuged for 30 minutes at 22,500 \times g to obtain supernatant. Ammonium sulfate was added to the supernatant to yield a final concentration of 0.4 M and allowed to stand for 1 hour. Again, the mixture was centrifuged for 30 minutes at 22,500 \times g to obtain supernatant.

The supernatant was loaded onto a TSK gel Butyl Toyopearl 650M column (10 ml bed volume per 100 g of roots), equilibrated with the adsorption buffer [20 mM Tris, 1 mM EDTA, 3 mM DTT, 0.4 M (NH₄)₂SO₄ and 0.1 mM p-APMSF, pH 8.0] and eluted with elution buffer [10 mM Tris, 1 mM EDTA, 3 mM DTT, 0.1 mM p-APMSF, 5% glycerol and 0.05% 3-[(3-chloramidopropyl) dimethyl-ammonio] propanesulfonic acid (hereinafter designates as CHAPS), pH 8.0].

KCl was added to the active fraction to give a final concentration of 0.4 M, and 1 M potassium phosphate buffer (pH 8.0) was added to a final concentration of 1 mM of KCl. A hydroxyapatite 100–350 mesh (Nacalai Tesque), equilibrated with the adsorption buffer (1 mM K-P, 10 mM KCl, 3 mM DTT and 0.1 mM p-APMSF, pH 8.0), was prepared at 10 ml per protein 100 mg and the fractions containing nicotianamine synthase were loaded. Nicotianamine synthase was passed through without adsorption. The passed through fraction was loaded onto TSK gel Butyl Toyopearl 650M column (1 ml bed volume per 10 mg of protein), and nicotianamine synthase was eluted in the manner described above.

The active fraction was loaded onto a DEAE-Sepharose FF column (5 ml bed volume per 25 mg of protein, Pharmacia) equilibrated with the adsorption buffer (20 mM Tris, 1 mM EDTA, 3 mM DTT, 0.1 mM p-APMSF and 0.05% CHAPS, pH 8.0) and eluted with stepwise gradient elution of potassium chloride concentration of 0.05 M, 0.1 M, 0.15 M and 0.2 M. Nicotianamine synthase was eluted at 0.15 M of KCl concentration.

12

The active fraction was loaded onto the Ether Toyopearl 650M column (10 ml bed volume per 100 g of roots), equilibrated with adsorption buffer [20 mM Tris, 1 mM EDTA, 3 mM DTT, 1.2 M (NH₄)₂SO₄ and 0.1 mM p-APMSF, pH 8.0]. Nicotianamine synthase was not adsorbed and passed through from the column. The passed through fraction was loaded onto TSK gel Butyl Toyopearl 650M column and fractions containing nicotianamine synthase was eluted. The peptides in the active fraction containing nicotianamine synthase, which was purified by the above column chromatographic treatments, were separated by sodium dodecyl sulfate—polyacrylamide gel electrophoresis (hereinafter designates as SDS-PAGE) using 11% acrylamide slab gels. After SDS-PAGE the gel was stained with 0.3 M copper chloride (Dzandu et al. 1988), and the separated bands were cut out. The gel fragments were destained with 0.25 M EDTA/0.25 M Tris (pH 9.0) and homogenized with the extraction buffer (1% SDS, 25 mM Tris and 192 mM glycine). Each homogenate was electroeluted with SDS-free buffer (25 mM Tris and 192 mM glycine) and peptide was recovered.

Example 4

Determination of Partial Amino Acid Sequence

The isolated nicotianamine synthase was digested chemically with cyanogen bromide (Gross 1967).

After SDS-PAGE treatment, 10-fold volume of 70% (v/v) formic acid and 1% (w/v) cyanogen bromide were added to gel fragments containing nicotianamine synthase and decomposed at 4° C. for overnight. After completion of digestion, the liquid part was collected and dried in vacuo. The dried substance was dissolved in SDS-PAGE sample buffer, and allowed to stand at room temperature for overnight, then the digested product was separated by SDS-PAGE using 16.5% acrylamide gel containing Tricine (Schagger and Jagow, 1987). The peptides were transferred onto a PVDF membrane by electroblotting (Towbin et al. 1979) and stained with amido black. The stained bands were cut out and the amino acid sequence was determined from N-terminal side of each peptide by Edman degradation in gas-phase sequencer (model 492A protein sequencer, Applied Biosystems Inc.).

Example 5

Cloning of Nicotianamine Synthase Genes

PCR amplification was conducted for cDNA originated from Fe-deficient barley roots using primers, which were synthesized based on the obtained partial amino acid sequence. A pYH23 cDNA library prepared from the poly (A)⁺RNA of Fe-deficient barley roots was screened with the thus obtained DNA fragments of PCR product, which was labeled with [α -³²P]dATP using the random primer kit (Takara Shuzo Co.), as the primers. The isolated cDNA clones were sequenced by cycle sequencing kit (Shimadzu Bunko Co.) using Shimadzu DNA sequencer DSQ-2000L.

PCR amplification was conducted for genomic DNA of *Arabidopsis thaliana* using primers, which were synthesized based on nucleotide sequences of AC003114 and AB005245 of *Arabidopsis thaliana*. The thus obtained DNA fragments were sequenced by cycle sequencing kit (Shimadzu Bunko Co.) using Shimadzu DNA sequencer DSQ-1000L.

The determined nucleotide sequence is shown in SEQ ID NO: 2.

13

Example 6

Expression of NAS1 Protein in *E. coli*

A fragment, in which EcoRI site was introduced into the upstream of the first ATG of the HvNAS1 cDNA and PstI and BamHI sites were introduced into the downstream of the stop codon of the HvNAS1 cDNA, was amplified by PCR. The thus obtained amplified product was subcloned in the pBluescriptII SK- using EcoRI site and BamHI site, and the correct nucleotide sequence was confirmed. The fragment between EcoRI site and PstI site was cloned into pMAL-c2 to make expression in the form of fusing the HvNAS1 to the C-terminal of maltose binding protein.

A fragment, in which EcoRI site was introduced into the upstream of the first ATG of the OsNAS1 and HindIII site was introduced into the downstream of the stop codon of the OsNAS1, was amplified by PCR. The thus obtained amplified product was subcloned in the pBluescriptII SK- using EcoRI site and HindIII site, and the correct nucleotide sequence was confirmed. The fragment between EcoRI site and HindIII site was cloned into pMAL-c2 to make expression in the form of fusing the OsNAS1 to the C-terminal of maltose binding protein.

A fragment, in which EcoRI site was introduced into the upstream of the first ATG of the AtNAS1, AtNAS2 and AtNAS3 and XbaI site was introduced into the downstream of the stop codon of the AtNAS1, AtNAS2 and AtNAS3, was amplified by PCR. The thus obtained amplified products were subcloned in the pBluescriptII SK-, and the correct nucleotide sequences were confirmed. The fragment between EcoRI site and XbaI site was cloned into pMAL-c2 to make expression in the form of fusing the AtNAS1, AtNAS2 and AtNAS3 to the C-terminal of maltose binding proteins, respectively.

E. coli strain XL1-Blue was used as a host for expressing the said fused protein. pMAL-c2-HvNAS1 and pMAL-c2, respectively, were introduced into YL1-Blue. The thus obtained recombinant bacteria were cultured in LB medium containing ampicillin and tetracycline, each 50 µg/ml, at 37° C. until the OD 600 of the culture reached 0.5. Isopropyl β-D-thiogalactopyranoside (IPTG) was added to the final concentration of 0.3 mM, and continuously cultured at 37° C. for 3 hours, and collected bacterial cells. Cells were suspended in 10 mM Tris buffer containing 0.2 M NaCl, 1 mM EDTA, 3 mM DTT and 0.1 mM E-64, pH 7.4 and frozen with liquid nitrogen. This was melted in ice water and ultrasonication for 15 seconds was repeated for 10 times. Nicotianamine synthase activity of the thus obtained crude extract was assayed according to the method described in example 2 and the enzyme activity was confirmed.

Example 7

Northern Hybridization

Northern hybridization of barley RNA was performed using DNA fragment, which was prepared by cutting HvNAS1 cDNA with HindIII and NotI and labeled with [α -³²P]dATP, as a probe. Total RNA was extracted from barley (Naito et al. 1988). The extracted RNA was separated by 1.4% agarose gel electrophoresis, and blotted onto Hybond-N⁺ membranes (Amersham). Northern hybridization of rice RNA was performed using OsNAS1 ORF, which was labeled with [α -³²P]dATP, as a probe. Total RNA was extracted from rice. The extracted RNA was separated by 1.4% agarose gel electrophoresis, and blotted onto Hybond-

14

N⁺ membranes (Amersham). The membrane was hybridized with the probe in 0.5 M Church phosphate buffer (Church and Gilbert 1984), 1 mM EDTA, 7% (w/v) SDS with 100 µg/ml salmon sperm DNA at 65° C. for overnight. The membrane was washed with buffer containing 40 mM Church phosphate buffer and 1% (w/v) SDS at 65° C. for 10 minutes. After the washing was repeated once again, the membrane was washed with buffer containing 0.2×SSPE and 0.1% (w/v) SDS at 65° C. for 10 minutes. Radioactivity was detected using the image analyzer BAS-2000.

Results are shown in FIG. 9 and FIG. 16.

Example 8

Southern Hybridization

Genomic DNA was extracted from leaves of barley and rice. The extract was digested with BamHI, EcoRI or HindIII, separated on a 0.8% (w/v) agarose gel electrophoresis, and transferred onto Hybond-N⁺ membranes (Amersham). The hybridization was performed according to the method described in example 7 and radioactivity was detected.

Result is shown in FIG. 10.

Example 9

Preparation of Polyclonal Antibody

Two rats were immunized using the antigen containing about 100 µg of isolated nicotianamine synthase. The antigen was the same sample as that determined the partial amino acid sequence. The complete Freund's adjuvant was used at the first immunization and the incomplete Freund's adjuvant was used since the second immunization. All the constituents of the blood were corrected after the rats were immunized four times, and the obtained serum was preserved at -80° C.

Example 10

Western Blotting Analysis

Total protein was extracted using trichloroacetic acid and acetone (Damerval et al. 1986). The plants were crashed in the liquid nitrogen until powder was obtained, and mixed with acetone containing 0.1% (v/v) 2-mercaptoethanol. The protein was precipitated by allowing to stand at -20° C. for 1 hour, and the precipitate was collected by centrifugation at 16,000×g for 30 minutes. The precipitate was suspended in acetone containing 0.1% (v/v) 2-mercaptoethanol and allowed to stand at -20° C. for 1 hour, then collected the precipitate by centrifugation at 16,000×g for 30 minutes. The precipitate was dried in vacuo, and dissolved in the sample buffer [9.5 M urea, 2% (w/v) Triton X-100 and 5% (v/v) 2-ME], then centrifuged at 16,000×g for 10 minutes to obtain the supernatant. The proteins contained in the supernatant were separated by SDS-PAGE or the denaturing two-dimensional electrophoresis (O'Farrell 1975) and transferred onto PVDF membrane. Western blotting analysis was performed by applying the primary antibody containing anti-nicotianamine synthase antibody prepared in example 9 and the secondary antibody containing horse radish binding anti-mouse IgG (H+L) goat antibody (Wako Pure Chemicals Co.) on the membrane and coloring with diaminobenzidine.

15

Result is shown in FIG. 12. SDS-PAGE was performed using 12.5% acrylamide slab gel. Protein 100 µg was electrophoresed. Proteins of roots 200 µg and leaves 500 µg were electrophoresed.

Example 11

RT-PCR

Total RNA was extracted from *Arabidopsis thaliana*. RT-PCR was performed with 1 µg RNA as a template by using the EZ rTh RNA PCR kit (Parkin Elmer Inc.). Specific primers for AtNAS1, AtNAS2 and AtNAS3, respectively, were used.

Result is shown in FIG. 18.

16

INDUSTRIAL APPLICABILITY

Various cells are transformed according to the conventional method by using recombinant vectors of the present invention. Mass production of nicotianamine can be performed by using the obtained transformant. These methods can be performed according to the method known in the person skilled in the art.

Selective breeding of plants, preferably graminaceous plants can also be performed using genes of the present invention. Especially, genes of the present invention can be applied for improving varieties, which can grow on Fe-deficient soil.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 37

<210> SEQ ID NO 1

<211> LENGTH: 328

<212> TYPE: PRT

<213> ORGANISM: Hordeum vulgare

<400> SEQUENCE: 1

```

Met Asp Ala Gln Asn Lys Glu Val Ala Ala Leu Ile Glu Lys Ile Ala
 1           5           10           15

Gly Ile Gln Ala Ala Ile Ala Glu Leu Pro Ser Leu Ser Pro Ser Pro
 20           25           30

Glu Val Asp Arg Leu Phe Thr Asp Leu Val Thr Ala Cys Val Pro Pro
 35           40           45

Ser Pro Val Asp Val Thr Lys Leu Ser Pro Glu His Gln Arg Met Arg
 50           55           60

Glu Ala Leu Ile Arg Leu Cys Ser Ala Ala Glu Gly Lys Leu Glu Ala
 65           70           75           80

His Tyr Ala Asp Leu Leu Ala Thr Phe Asp Asn Pro Leu Asp His Leu
 85           90           95

Gly Leu Phe Pro Tyr Tyr Ser Asn Tyr Val Asn Leu Ser Arg Leu Glu
 100          105          110

Tyr Glu Leu Leu Ala Arg His Val Pro Gly Ile Ala Pro Ala Arg Val
 115          120          125

Ala Phe Val Gly Ser Gly Pro Leu Pro Phe Ser Ser Leu Val Leu Ala
 130          135          140

Ala His His Leu Pro Glu Thr Gln Phe Asp Asn Tyr Asp Leu Cys Gly
 145          150          155          160

Ala Ala Asn Glu Arg Ala Arg Lys Leu Phe Gly Ala Thr Ala Asp Gly
 165          170          175

Val Gly Ala Arg Met Ser Phe His Thr Ala Asp Val Ala Asp Leu Thr
 180          185          190

Gln Glu Leu Gly Ala Tyr Asp Val Val Phe Leu Ala Ala Leu Val Gly
 195          200          205

Met Ala Ala Glu Glu Lys Ala Lys Val Ile Ala His Leu Gly Ala His
 210          215          220

Met Val Glu Gly Ala Ser Leu Val Val Arg Ser Ala Arg Pro Arg Gly
 225          230          235          240

Phe Leu Tyr Pro Ile Val Asp Pro Glu Asp Ile Arg Arg Gly Gly Phe
 245          250          255

```

-continued

Glu Val Leu Ala Val His His Pro Glu Gly Glu Val Ile Asn Ser Val
 260 265 270
 Ile Val Ala Arg Lys Ala Val Glu Ala Gln Leu Ser Gly Pro Gln Asn
 275 280 285
 Gly Asp Ala His Ala Arg Gly Ala Val Pro Leu Val Ser Pro Pro Cys
 290 295 300
 Asn Phe Ser Thr Lys Met Glu Ala Ser Ala Leu Glu Lys Ser Glu Glu
 305 310 315 320
 Leu Thr Ala Lys Glu Leu Ala Phe
 325

<210> SEQ ID NO 2
 <211> LENGTH: 1295
 <212> TYPE: DNA
 <213> ORGANISM: Hordeum vulgare
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (55)..(1041)

<400> SEQUENCE: 2

gcggttcagag gcttccagag ttcttccggt caccaagaag catttgatca taac atg 57
 Met
 1
 gat gcc cag aac aag gag gtc gct gct ctg atc gag aag atc gcc ggt 105
 Asp Ala Gln Asn Lys Glu Val Ala Ala Leu Ile Glu Lys Ile Ala Gly
 5 10 15
 atc cag gcc gcc atc gcc gag ctg ccg tcg ctg agc ccg tcc ccc gag 153
 Ile Gln Ala Ala Ile Ala Glu Leu Pro Ser Leu Ser Pro Ser Pro Glu
 20 25 30
 gtc gac agg ctc ttc acc gac ctc gtc acg gcc tgc gtc ccg ccg agc 201
 Val Asp Arg Leu Phe Thr Asp Leu Val Thr Ala Cys Val Pro Pro Ser
 35 40 45
 ccc gtc gac gtg acg aag ctc agc ccg gag cac cag agg atg cgg gag 249
 Pro Val Asp Val Thr Lys Leu Ser Pro Glu His Gln Arg Met Arg Glu
 50 55 60 65
 gct ctc atc cgc ttg tgc tcc gcc gcc gag ggg aag ctc gag gcg cac 297
 Ala Leu Ile Arg Leu Cys Ser Ala Ala Glu Gly Lys Leu Glu Ala His
 70 75 80
 tac gcc gac ctg ctc gcc acc ttc gac aac ccg ctc gac cac ctc ggc 345
 Tyr Ala Asp Leu Leu Ala Thr Phe Asp Asn Pro Leu Asp His Leu Gly
 85 90 95
 ctc ttc ccg tac tac agc aac tac gtc aac ctc agc agg ctg gag tac 393
 Leu Phe Pro Tyr Tyr Ser Asn Tyr Val Asn Leu Ser Arg Leu Glu Tyr
 100 105 110
 gag ctc ctg gcg cgc cac gtg ccg ggc atc gcg ccg gcg cgc gtc gcc 441
 Glu Leu Leu Ala Arg His Val Pro Gly Ile Ala Pro Ala Arg Val Ala
 115 120 125
 ttc gtc ggc tcc gcc ccg ctg ccg ttc agc tcg ctc gtc ctc gcc gcg 489
 Phe Val Gly Ser Gly Pro Leu Pro Phe Ser Ser Leu Val Leu Ala Ala
 130 135 140 145
 cac cac ctg ccc gag acc cag ttc gac aac tac gac ctg tgc ggc gcg 537
 His His Leu Pro Glu Thr Gln Phe Asp Asn Tyr Asp Leu Cys Gly Ala
 150 155 160
 gcc aac gag cgc gcc agg aag ctg ttc gcc gcg acg gcg gac gcc gtc 585
 Ala Asn Glu Arg Ala Arg Lys Leu Phe Gly Ala Thr Ala Asp Gly Val
 165 170 175
 ggc gcg cgt atg tcg ttc cac acg gcg gac gtc gcc gac ctc acc cag 633
 Gly Ala Arg Met Ser Phe His Thr Ala Asp Val Ala Asp Leu Thr Gln
 180 185 190

-continued

gag ctc ggc gcc tac gac	gtg gtc ttc ctc gcc	gcg ctc gtc ggc atg	681
Glu Leu Gly Ala Tyr Asp	Val Val Phe Leu Ala	Ala Leu Val Gly Met	
195	200	205	
gca gcc gag gag aag gcc	aag gtg att gcc cac	ctg ggc gcg cac atg	729
Ala Ala Glu Glu Lys Ala	Lys Val Ile Ala His	Leu Gly Ala His Met	
210	215	220 225	
gtg gag ggg gcg tcc ctg	gtc gtg cgg agc gca	cgg ccc cgc ggc ttt	777
Val Glu Gly Ala Ser Leu	Val Val Arg Ser Ala	Arg Pro Arg Gly Phe	
	230	235 240	
ctt tac ccc att gtc gac	ccg gag gac atc agg	cgg ggt ggg ttc gag	825
Leu Tyr Pro Ile Val Asp	Pro Glu Asp Ile Arg	Arg Gly Gly Phe Glu	
	245	250 255	
gtg ctg gcc gtg cac cac	ccg gaa ggt gag gtg	atc aac tct gtc atc	873
Val Leu Ala Val His His	Pro Glu Gly Glu Val	Ile Asn Ser Val Ile	
	260	265 270	
gtc gcc cgt aag gcc gtc	gaa gcg cag ctc agt	ggg ccg cag aac gga	921
Val Ala Arg Lys Ala Val	Glu Ala Gln Leu Ser	Gly Pro Gln Asn Gly	
	275	280 285	
gac gcg cac gca cgg gcc	gcg gtg ccg ttg gtc	agc ccg cca tgc aac	969
Asp Ala His Ala Arg Gly	Ala Val Pro Leu Val	Ser Pro Pro Cys Asn	
290	295	300 305	
ttc tcc acc aag atg gag	gcg agc gcg ctt gag	aag agc gag gag ctg	1017
Phe Ser Thr Lys Met Glu	Ala Ser Ala Leu Glu	Lys Ser Glu Glu Leu	
	310	315 320	
acc gcc aaa gag ctg gcc	ttt tga ttgaagagtg	cgcgtggtca tctgtcgcc	1071
Thr Ala Lys Glu Leu Ala	Phe		
	325		
tgcgatcgtg gtaactttcc	tactcgtgtg tgttttgatg	tttgtgcctg taagagttat	1131
gcttcgpgcc ttgtgctggt	aatttacacg cgttacatgt	agtacttgta tttatacctg	1191
gaataacggt atgtaacata	aatattagtg ggatttgaag	tgtaatgcta aataataaga	1251
aaacttgatg cagacattca	aaaaaaaaa aaaaaaaaaa	aaaa	1295

<210> SEQ ID NO 3

<211> LENGTH: 335

<212> TYPE: PRT

<213> ORGANISM: Hordeum vulgare

<400> SEQUENCE: 3

Met Ala Ala Gln Asn Asn	Gln Glu Val Asp Ala	Leu Val Glu Lys Ile
1	5	10 15
Thr Gly Leu His Ala Ala	Ile Ala Lys Leu Pro	Ser Leu Ser Pro Ser
	20	25 30
Pro Asp Val Asp Ala Leu	Phe Thr Glu Leu Val	Thr Ala Cys Val Pro
	35	40 45
Pro Ser Pro Val Asp Val	Thr Lys Leu Gly Pro	Glu Ala Gln Glu Met
	50	55 60
Arg Glu Gly Leu Ile Arg	Leu Cys Ser Glu Ala	Glu Gly Lys Leu Glu
65	70	75 80
Ala His Tyr Ser Asp Met	Leu Ala Ala Phe Asp	Lys Pro Leu Asp His
	85	90 95
Leu Gly Met Phe Pro Tyr	Tyr Asn Asn Tyr Ile	Asn Leu Ser Lys Leu
	100	105 110
Glu Tyr Glu Leu Leu Ala	Arg Tyr Val Pro Gly	Gly Tyr Arg Pro Ala
	115	120 125
Arg Val Ala Phe Ile Gly	Ser Gly Pro Leu Pro	Phe Ser Ser Phe Val
	130	135 140

-continued

Leu Ala Ala Arg His Leu Pro Asp Thr Met Phe Asp Asn Tyr Asp Leu
 145 150 155 160

Cys Gly Ala Ala Asn Asp Arg Ala Ser Lys Leu Phe Arg Ala Asp Arg
 165 170 175

Asp Val Gly Ala Arg Met Ser Phe His Thr Ala Asp Val Ala Asp Leu
 180 185 190

Ala Gly Glu Leu Ala Lys Tyr Asp Val Val Phe Leu Ala Ala Leu Val
 195 200 205

Gly Met Ala Ala Glu Asp Lys Ala Lys Val Ile Ala His Leu Gly Ala
 210 215 220

His Met Ala Asp Gly Ala Ala Leu Val Val Arg Ser Ala His Gly Ala
 225 230 235 240

Arg Gly Phe Leu Tyr Pro Ile Val Asp Pro Gln Asp Ile Gly Arg Gly
 245 250 255

Gly Phe Glu Val Leu Ala Val Cys His Pro Asp Asp Asp Val Val Asn
 260 265 270

Ser Val Ile Ile Ala Gln Lys Ser Lys Asp Val His Ala Asp Gly Leu
 275 280 285

Gly Ser Gly Arg Gly Ala Gly Gly Gln Tyr Ala Arg Gly Thr Val Pro
 290 295 300

Val Val Ser Pro Pro Cys Arg Phe Gly Glu Met Val Ala Asp Val Thr
 305 310 315 320

Gln Asn His Lys Arg Asp Glu Phe Ala Asn Ala Glu Val Ala Phe
 325 330 335

<210> SEQ ID NO 4
 <211> LENGTH: 1342
 <212> TYPE: DNA
 <213> ORGANISM: Hordeum vulgare

<400> SEQUENCE: 4

```

ctcctgtgcc tgtcctgagg taccaagaac accagtgaaa tggctgcccc gaacaaccag    60
gaggtggatg ccctgggtga gaagatcacc gggctccatg ccgcaatcgc caagctgccg    120
tcgctcagcc catccccgga cgtcgacgcg ctcttcacgg agctggtcac ggcgtgogtt    180
ccaccgagtc cagtggacgt gaccaagctc gggccggagg cgcaggagat gcgggagggc    240
ctcatccgcc tatgtccga ggcgagggg aagctggagg cgcactactc cgacatgctc    300
gccgccttcg acaagccgct ggatcacctc ggcctgttcc cctactacia caactacatc    360
aacctcagca agctcgagta cgagctcctg gcccgctacg tgcctggcgg ctatcgcccg    420
gcgcgcgtcg cgttcacggt ctcgggcccg ctgcccgtca gctcctttgt cctggccgcg    480
cgccaactgc ccgacacat gttogacaac tatgaactgt gcggtgcggc caacgatcgc    540
gccagcaagc tcttcgcgcg ggatcgcgac gtgggtgccc gcatgtcgtt ccacacggcc    600
gacgtcgcgg acctcgcccg cgagctcgcc aagtacgacg ttgtcttctt ggccgcactc    660
gtcggcatgg ccgagcagga caagcgaag gtgatcgcgc acctcggcgc acacatggca    720
gacggggcgg ccctcgtcgt gcgcagcgca cacggagcgc gcgggttctt gtaccgatc    780
gtcgaccccc aggacatcgg ccgagcggg ttcgaggtgc tggccgtgtg ccaccccgac    840
gacgacgtgg tgaactccgt catcatcgca cagaagtcca aggaogtga tgccgatgga    900
cttgccagcg ggcgtggtgc cgggtggacag tacgcgcggg gcacggtgcc tgttgcagc    960
cccccgctga ggttcggcga gatggtggcg gacgtgacct agaaccacia gagagacgag   1020
tttgccaacg ccgaagtggc cttttgatcg ttcgctgcga ggggtgcat ccatgatcca   1080
    
```

-continued

```

tccatcacctc gttctgtgat tgcacaaagc ttgcaatcgt atgcatttca agtcacgtgt 1140
tgcttctatc caataatgta cgtgtgggtgt ttacacgcga atgtcttgta gacctttgta 1200
tgtgtacaag tgaattttaa ttcacaagta catataatgg tcaccattga aaagatgttt 1260
agtgtgtgtt ttccaatata tgtttgtgta aggttcatca tctaataaaa tatgtttgga 1320
acccaaaaaa aaaaaaaaaa aa 1342

```

```

<210> SEQ ID NO 5
<211> LENGTH: 335
<212> TYPE: PRT
<213> ORGANISM: Hordeum vulgare

```

```

<400> SEQUENCE: 5

```

```

Met Ala Ala Gln Asn Asn Asn Lys Asp Val Ala Ala Leu Val Glu Lys
 1          5          10          15
Ile Thr Gly Leu His Ala Ala Ile Ala Lys Leu Pro Ser Leu Ser Pro
 20          25          30
Ser Pro Asp Val Asp Ala Leu Phe Thr Glu Leu Val Thr Ala Cys Val
 35          40          45
Pro Pro Ser Pro Val Asp Val Thr Lys Leu Gly Pro Glu Ala Gln Glu
 50          55          60
Met Arg Glu Gly Leu Ile Arg Leu Cys Ser Glu Ala Glu Gly Lys Leu
 65          70          75          80
Glu Ala His Tyr Ser Asp Met Leu Ala Ala Phe Asp Asn Pro Leu Asp
 85          90          95
His Leu Gly Ile Phe Pro Tyr Tyr Ser Asn Tyr Ile Asn Leu Ser Lys
100          105          110
Leu Glu Tyr Glu Leu Leu Ala Arg Tyr Val Arg Arg His Arg Pro Ala
115          120          125
Arg Val Ala Phe Ile Gly Ser Gly Pro Leu Pro Phe Ser Ser Phe Val
130          135          140
Leu Ala Ala Arg His Leu Pro Asp Thr Met Phe Asp Asn Tyr Asp Leu
145          150          155          160
Cys Gly Ala Ala Asn Asp Arg Ala Ser Lys Leu Phe Arg Ala Asp Thr
165          170          175
Asp Val Gly Ala Arg Met Ser Phe His Thr Ala Asp Val Ala Asp Leu
180          185          190
Ala Ser Glu Leu Ala Lys Tyr Asp Val Val Phe Leu Ala Ala Leu Val
195          200          205
Gly Met Ala Ala Glu Asp Lys Ala Lys Val Ile Ala His Leu Gly Ala
210          215          220
His Met Ala Asp Gly Ala Ala Leu Val Val Arg Ser Ala His Gly Ala
225          230          235          240
Arg Gly Phe Leu Tyr Pro Ile Val Asp Pro Gln Asp Ile Gly Arg Gly
245          250          255
Gly Phe Glu Val Leu Ala Val Cys His Pro Asp Asp Asp Val Val Asn
260          265          270
Ser Val Ile Ala Gln Lys Ser Lys Glu Val His Ala Asp Gly Leu
275          280          285
Gly Ser Ala Arg Gly Ala Gly Arg Gln Tyr Ala Arg Gly Thr Val Pro
290          295          300
Val Val Ser Pro Pro Cys Arg Phe Gly Glu Met Val Ala Asp Val Thr
305          310          315          320

```

-continued

Gln Asn His Lys Arg Asp Glu Phe Ala Asn Ala Glu Val Ala Phe
 325 330 335

<210> SEQ ID NO 6
 <211> LENGTH: 1314
 <212> TYPE: DNA
 <213> ORGANISM: Hordeum vulgare

<400> SEQUENCE: 6

```

ctacttcact cacactagtg cccagaaaga aggctgcaat ggctgcccag aacaacaaca    60
aggatgtcgc tgccctgggtg gagaagatca ccgggctcca cgccgccatc gccaaagtgc    120
cgtcgcctcag cccatccccg gacgtcgcgc cgctcttcac cgagctggtc acggcgtgcg    180
ttcccccgag ccccgtagac gtgaccaagc tcggccccga ggcgcaggag atgcgggagg    240
gcctcatccg cctctgtccc gaggccgagg ggaagctgga ggcgcaactac tccgacatgc    300
tcggcgcctt cgacaacccg ctggatcacc tcggcatctt cccctactac agcaactaca    360
tcaacctcag caagctggag tacgagctcc tggcacgcta cgtccggcgg catcgcccgg    420
cccgcgtcgc gttcatcggc tccggccccg tcgccgttcag ctcccttctc ctggccgcgc    480
ggccacctgcc cgacaccatg tttgacaact acgacctttg cggcgcgggc aacgatcgcg    540
ccagcaagct cttccgcgcg gacacggacg tgggtgcccg catgtcgttc cacacggccg    600
acgtcgcgga cctcgcacgc gagctcgcca agtacgacgt cgtcttctct gcccgcctcg    660
tcggcatggc cgccgaggac aaggccaagg tgatcgcgca cctcggcgca cacatggcag    720
acggggcggc cctcgtcgtg cgcagcgcac acggagcgcg cgggttcctg tacccgattg    780
tcgacccccg ggacatcggc cgcggcgggg tcgaggtgct ggccgtgtgc caccocgacg    840
acgacgtggt gaactccgtc atcatcgcac agaagtcaa ggaggtgcat gccgatggac    900
ttggcagcgc gcgtggtgcc ggtcgacagt acgcgcgcgg cacggtgccg gttgtcagcc    960
ccccgtgcag gttcggtag atggtggcgg atgtgaccca gaaccacaag agagacgagt    1020
ttgccaacgc cgaagtggcc tttgatcga tcgtcgccaa gggacaataa atgaacgtgg    1080
atgtggtagg gtaatttggc tacctcgtg cttgatcgtc tgcaatatgt gcacattttc    1140
ctactaccgc tgcttatgca tttcaagcca tgtgatgttg gtatccaata aagtatgtgt    1200
agggtttaca cgaaatgtc tttacacctt gtacgtgtaa gtgttgacaa cgatgaattt    1260
cagttcacia ttaataaata gtataatga ttcaaaaaaa aaaaaaaaaa aaaa    1314

```

<210> SEQ ID NO 7
 <211> LENGTH: 329
 <212> TYPE: PRT
 <213> ORGANISM: Hordeum vulgare

<400> SEQUENCE: 7

```

Met Asp Gly Gln Ser Glu Glu Val Asp Ala Leu Val Gln Lys Ile Thr
  1          5          10          15

Gly Leu His Ala Ala Ile Ala Lys Leu Pro Ser Leu Ser Pro Ser Pro
  20          25          30

Asp Val Asp Ala Leu Phe Thr Asp Leu Val Thr Ala Cys Val Pro Pro
  35          40          45

Ser Pro Val Asp Val Thr Lys Leu Ala Pro Glu Ala Gln Ala Met Arg
  50          55          60

Glu Gly Leu Ile Arg Leu Cys Ser Glu Ala Glu Gly Lys Leu Glu Ala
  65          70          75          80

His Tyr Ser Asp Met Leu Ala Ala Phe Asp Asn Pro Leu Asp His Leu

```


-continued

				85					90					95					
Gly	Val	Phe	Pro	Tyr	Tyr	Ser	Asn	Tyr	Ile	Asn	Leu	Ser	Lys	Leu	Glu				
				100					105					110					
Tyr	Glu	Leu	Leu	Ala	Arg	Tyr	Val	Pro	Gly	Arg	His	Arg	Pro	Ala	Arg				
				115					120					125					
Val	Ala	Phe	Ile	Gly	Ser	Gly	Pro	Leu	Pro	Phe	Ser	Ser	Tyr	Val	Leu				
				130					135					140					
Ala	Ala	Arg	His	Leu	Pro	Asp	Thr	Val	Phe	Asp	Asn	Tyr	Asp	Leu	Cys				
				145					150					155					
Gly	Ala	Ala	Asn	Asp	Arg	Ala	Thr	Arg	Leu	Phe	Arg	Ala	Asp	Lys	Asp				
				165					170					175					
Val	Gly	Ala	Arg	Met	Ser	Phe	His	Thr	Ala	Asp	Val	Ala	Asp	Leu	Thr				
				180					185					190					
Asp	Glu	Leu	Ala	Thr	Tyr	Asp	Val	Val	Phe	Leu	Ala	Ala	Leu	Val	Gly				
				195					200					205					
Met	Ala	Ala	Glu	Asp	Lys	Ala	Lys	Val	Ile	Ala	His	Leu	Gly	Ala	His				
				210					215					220					
Met	Ala	Asp	Gly	Ala	Ala	Leu	Val	Ala	Arg	His	Gly	Ala	Arg	Gly	Phe				
				225					230					235					
Leu	Tyr	Pro	Ile	Val	Asp	Pro	Gln	Asp	Ile	Gly	Arg	Gly	Gly	Phe	Glu				
				245					250					255					
Val	Leu	Ala	Val	Cys	His	Pro	Asp	Asp	Asp	Val	Val	Asn	Ser	Val	Ile				
				260					265					270					
Ile	Ala	Gln	Lys	Ser	Asn	Asp	Val	His	Glu	Tyr	Gly	Leu	Gly	Ser	Gly				
				275					280					285					
Arg	Gly	Gly	Arg	Tyr	Ala	Arg	Gly	Thr	Val	Val	Pro	Val	Val	Ser	Pro				
				290					295					300					
Pro	Cys	Arg	Phe	Gly	Glu	Met	Val	Ala	Asp	Val	Thr	Gln	Lys	Arg	Glu				
				305					310					315					
Glu	Phe	Ala	Asn	Ala	Glu	Val	Ala	Phe											
				325															

<210> SEQ ID NO 8
 <211> LENGTH: 1249
 <212> TYPE: DNA
 <213> ORGANISM: Hordeum vulgare

<400> SEQUENCE: 8

```

ccactaccga ctaccgtagt accgtgctc agagctcatc actggtcagg taccaagaag      60
acataaaaat ggacggccag agcgaggagg tcgacgcctt tgtccagaag atcaccggcc      120
tccacgcccgc catcgccaag ctgccctcgc tcagcccgtc cccggacgtc gacgcgtctt      180
tcaccgacct ggtaaccgcg tgcgtgcccc cgagccccgt ggacgtgacc aagctcgccc      240
cggaggcgca ggcgatgcgg gagggcctca tccgcctctg ctccgaggcc gagggcaagc      300
tggaggcgca ctactccgac atgctcgccg ccttcgacaa cccgctcgac cacctcggcg      360
tcttccccta ctacagcaac tacatcaacc tcagcaagct tgagtacgag ctctcgcgcg      420
gtctacgtgc cggcaggcat cgcccggccc gcgtcgcctt catcggctcc ggcccgtgctc      480
cgttcagctc ctacgtcctc gccgcgcgcc acctgcccga caccgtgttc gacaactacg      540
acctgtgcgg cgcggccaac gaccgcgcga ccaggctgtt ccgcgcggac aaggacgtcg      600
gcgcccgcat gtcgttccac accgccgacg tcgcggaact caccgacgag ctcgctacgt      660
acgacgtcgt cttcctggcc gcgctcgtgg gcatggcccg cgaggacaag gccaaggtga      720
    
```

-continued

```

tcgcgcacct tggcgcgcac atggcgggacg gggcgggccct cgttgcgcgcg cacggcgcgc 780
gtgggttcct ctacccgatc gtcgatcccc aggacatcgg tcgaggcggg ttcgaggtgc 840
tcgccgtgtg tcaccccgac gacgacgtgg tgaactccgt catcatcgca caaaagagca 900
acgacgtgca cgagtatgga cttggcagcg ggcgtggtgg acggtacgcg cgaggcacgg 960
tggtgccggt ggtcagccca ccctgcaggt tcggcgagat ggtggcagac gtgaccaga 1020
agagagagga gtttgccaac gcggaagtgg ccttctgatt gctgctgaat cgcttgtgat 1080
cgtacgtggt aatitttcta ctactctcc tcctaccacc acctatcacc tatgtatgca 1140
tttcaagtcg tgtgtgttt gtatccaata atgtaagtga gatgtttaca cgcgcaaaaa 1200
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1249
    
```

```

<210> SEQ ID NO 9
<211> LENGTH: 282
<212> TYPE: PRT
<213> ORGANISM: Hordeum vulgare
    
```

<400> SEQUENCE: 9

```

Met Glu Ala Glu Asn Gly Glu Val Ala Ala Leu Val Glu Lys Ile Thr
  1             5             10
Gly Leu His Ala Ala Ile Ser Lys Leu Pro Ala Leu Ser Pro Ser Pro
             20             25             30
Gln Val Asp Ala Leu Phe Thr Glu Leu Val Ala Ala Cys Val Pro Ser
             35             40             45
Ser Pro Val Asp Val Thr Lys Leu Gly Pro Glu Ala Gln Glu Met Arg
             50             55             60
Gln Asp Leu Ile Arg Leu Cys Ser Ala Ala Glu Gly Leu Leu Glu Ala
             65             70             75             80
His Tyr Ser Asp Met Leu Thr Ala Leu Asp Ser Pro Leu Asp His Leu
             85             90             95
Gly Arg Phe Pro Tyr Phe Asp Asn Tyr Val Asn Leu Ser Lys Leu Glu
             100            105            110
His Asp Leu Leu Ala Gly His Val Ala Ala Pro Ala Arg Val Ala Phe
             115            120            125
Ile Gly Ser Gly Pro Leu Pro Phe Ser Ser Leu Phe Leu Ala Thr Tyr
             130            135            140
His Leu Pro Asp Thr Arg Phe Asp Asn Tyr Asp Arg Cys Ser Val Ala
             145            150            155            160
Asn Gly Arg Ala Met Lys Leu Val Gly Ala Ala Asp Glu Gly Val Arg
             165            170            175
Ser Arg Met Ala Phe His Thr Ala Glu Val Thr Asp Leu Thr Ala Glu
             180            185            190
Leu Gly Ala Tyr Asp Val Val Phe Leu Ala Ala Leu Val Gly Met Thr
             195            200            205
Ser Lys Glu Lys Ala Asp Ala Ile Ala His Leu Gly Lys His Met Ala
             210            215            220
Asp Gly Ala Val Leu Val Arg Glu Ala Leu His Gly Ala Arg Ala Phe
             225            230            235            240
Leu Tyr Pro Val Val Glu Leu Asp Asp Val Gly Arg Gly Gly Phe Gln
             245            250            255
Val Leu Ala Val His His Pro Ala Gly Asp Glu Val Phe Asn Ser Phe
             260            265            270
Ile Val Ala Arg Lys Val Lys Met Ser Ala
             275            280
    
```

-continued

<210> SEQ ID NO 10
 <211> LENGTH: 1044
 <212> TYPE: DNA
 <213> ORGANISM: Hordeum vulgare

<400> SEQUENCE: 10

```

gtgacatgga ggccgaaaac ggcgaggtgg ctgctctggt cgagaagatc accggtctcc    60
acgccgccat ctccaagctc ccggcactaa gcccgctctcc tcaagtcgac gcgctcttca    120
ccgagctggt tgcggcgtgc gtcccatcaa gcccggtgga cgtgaccaag ctcggcccg    180
aggcgcagga gatcggcgag gacctcatcc gtctctgctc ggccgccgag gggctgctcg    240
aggcgacta ctccgacatg ctaccgcgt tggacagccc gctcgaccac ctcgcccgct    300
tcccttactt cgacaactac gtcaacctca gcaagctcga gcacgatctt ctggcaggtc    360
acgtggcggc cccggcccgc gtggcggttca tcgggtcggg gccactgccg ttcagctcgc    420
tcttccttgc gacgtaccac ctgccggaca cccggttcga caactacgac cggtgacgcg    480
tggcgaatgg ccgggcgatg aagctggctc gcgcggcgga cgagggcgtg cgatcacgca    540
tggcggttca cacggccgaa gtcacggacc tcacggctga gctcggcgct tacgacgtgg    600
tcttcctggc cgcgctcgtg ggaatgacgt ccaaggagaa ggccgacgcc atagcgcact    660
tggggaagca catggcagat ggggcgggtc tcgtgcgcga agcgtgcac ggggcgcgag    720
cgttcctgta tcctgtcgtg gagctggacg atgtcggggc tgggtgggtc caagtgtcgg    780
ccgtgcacca ccctgcaggc gatgaggtgt tcaactcatt catagttgcc cggaaggatga    840
aatgagtgc ttaattaag aaaagggtga gcctgtctgc ttgtgcaaat ggtgtctcac    900
attgataata accagatgat accctgcaca ttgatggggg tactgcagta tgtttcaatg    960
aggctcgtgt gtatcaaata tgagtattg gcttaataat atcagcgaat atgtttcgat   1020
taaaaaaaaa aaaaaaaaaa aaaa                                           1044
  
```

<210> SEQ ID NO 11
 <211> LENGTH: 328
 <212> TYPE: PRT
 <213> ORGANISM: Hordeum vulgare

<400> SEQUENCE: 11

```

Met Asp Ala Gln Asn Lys Glu Val Asp Ala Leu Val Gln Lys Ile Thr
  1          5          10          15
Gly Leu His Ala Ala Ile Ala Lys Leu Pro Ser Leu Ser Pro Ser Pro
          20          25          30
Asp Val Asp Ala Leu Phe Thr Asp Leu Val Thr Ala Cys Val Pro Pro
          35          40          45
Ser Pro Val Asp Val Thr Lys Leu Gly Ser Glu Ala Gln Glu Met Arg
          50          55          60
Glu Gly Leu Ile Arg Leu Cys Ser Glu Ala Glu Gly Lys Leu Glu Ala
          65          70          75          80
His Tyr Ser Asp Met Leu Ala Ala Phe Asp Asn Pro Leu Asp His Leu
          85          90          95
Gly Met Phe Pro Tyr Tyr Ser Asn Tyr Ile Asn Leu Ser Lys Leu Glu
          100          105          110
Tyr Glu Leu Leu Ala Arg Tyr Val Pro Gly Gly Ile Ala Arg Pro Ala
          115          120          125
Val Ala Phe Ile Gly Ser Gly Pro Leu Pro Phe Ser Ser Tyr Val Leu
          130          135          140
  
```

-continued

Ala Ala Arg His Leu Pro Asp Ala Met Phe Asp Asn Tyr Asp Leu Cys
 145 150 155 160
 Ser Ala Ala Asn Asp Arg Ala Ser Lys Leu Phe Arg Ala Asp Lys Asp
 165 170 175
 Val Gly Ala Arg Met Ser Phe His Thr Ala Asp Val Ala Asp Leu Thr
 180 185 190
 Arg Glu Leu Ala Ala Tyr Asp Val Val Phe Leu Ala Ala Leu Val Gly
 195 200 205
 Met Ala Ala Glu Asp Lys Ala Lys Val Ile Pro His Leu Gly Ala His
 210 215 220
 Met Ala Asp Gly Ala Ala Leu Val Val Arg Ser Ala Gln Ala Arg Gly
 225 230 235 240
 Phe Leu Tyr Pro Ile Val Asp Pro Gln Asp Ile Gly Arg Gly Gly Phe
 245 250 255
 Glu Val Leu Ala Val Cys His Pro Asp Asp Asp Val Val Asn Ser Val
 260 265 270
 Ile Ile Ala His Lys Ser Lys Asp Val His Ala Asn Glu Arg Pro Asn
 275 280 285
 Gly Arg Gly Gly Gln Tyr Arg Gly Ala Val Pro Val Val Ser Pro Pro
 290 295 300
 Cys Arg Phe Gly Glu Met Val Ala Asp Val Thr His Lys Arg Glu Glu
 305 310 315 320
 Phe Thr Asn Ala Glu Val Ala Phe
 325

<210> SEQ ID NO 12
 <211> LENGTH: 1352
 <212> TYPE: DNA
 <213> ORGANISM: Hordeum vulgare

<400> SEQUENCE: 12

ctccacttcg ctctgtgccc tcaggtagcc acaacataca gtattaaaat ggatgcccag 60
 aacaaggagg ttgatgcctt ggtccagaag atcaccggcc tccacgcccgc catcgccaag 120
 ctgcccgtccc tcagcccata acccgacgtc gacgcgctct tcaccgacct ggtcaccgcg 180
 tgggtccccc cgagcccctg ggaogtgacc aagctcgggt cggagggcga ggagatgagg 240
 gagggcctca tccgcctctg ctccgaggcc gaggggaagc tggagggcga ctactccgac 300
 atgctggcgg ccttcgacaa cccgctcgac cacctcgcca tgttccccta ctacagcaac 360
 tacatcaacc tcagcaagct ggagtagcag ctctctggcg gctacgtgcc gggcgccatc 420
 gcccgcccgc ctgtcgcggt catcggtccc ggcccgtgac cgttcagctc ctacgtcctc 480
 gccgctcgcc acctgcccga cgccatgttc gacaactacg acctgtgtag cgcggccaac 540
 gaccgtgcga gcaagctggt ccgocgggac aaggacgtgg gcgcccgcac gtctttccac 600
 accgcccagc tagcggacct caccgcgag ctccgcccgt acgacgtcgt cttcctggcc 660
 gcgctcgtgg gcatggctgc cgaggacaag gcccaaggtga ttccgcacct cggcgcgcac 720
 atggcggagc gggcgcccct cgtcgtgccc agtgccgagg cacgtggggt cctctaaccg 780
 atcgtcgatc cccaggacat cggtcgaggc gggtttgagg tgctggccgt gtgtaccccc 840
 gacgatgagc tgggtaactc cgtcatcacc gcacacaagt ccaaggacgt gcatgccaat 900
 gaacgtccca acggggctgg tggacagtac cggggcgcgg taccgggtgt cagcccgcgg 960
 tgcaggttcg gtgagatggt ggcggacgtg acccacaaga gagaggagtt caccaacgcg 1020

-continued

```

gaagtggcct tctgatcggt gcgaggggaat gaaaatgaag gtggacgtgt gtggtcagca 1080
tccatacgtg gctgctgtct tcacgcttg caatcgctact actacctacc tatgcagttc 1140
aagtcacgtg ttgtcaatgt aagtgtgatg tttacactag tctatgaaag gcagggcaga 1200
cgagggtagt gtgccaagta acagtgtgtc attataggtg taagtgttga gaataagacc 1260
atTTTTgttc acaaatagta tgatgtaatc ggtgtcatat tcgtattgag tacattgttc 1320
aagttggttg ctaaaaaaaaa aaaaaaaaaa aa 1352

```

<210> SEQ ID NO 13

<211> LENGTH: 329

<212> TYPE: PRT

<213> ORGANISM: Hordeum vulgare

<400> SEQUENCE: 13

```

Met Asp Ala Gln Ser Lys Glu Val Asp Ala Leu Val Gln Lys Ile Thr
  1          5          10          15
Gly Leu His Ala Ala Ile Ala Lys Leu Pro Ser Leu Ser Pro Ser Pro
          20          25          30
Asp Val Asp Ala Leu Phe Thr Asp Leu Val Thr Ala Cys Val Pro Pro
          35          40          45
Ser Pro Val Asp Val Thr Lys Leu Ala Pro Glu Ala Gln Ala Met Arg
          50          55          60
Glu Gly Leu Ile Arg Leu Cys Ser Glu Ala Glu Gly Lys Leu Glu Ala
          65          70          75          80
His Tyr Ser Asp Met Leu Ala Ala Phe Asp Asn Pro Leu Asp His Leu
          85          90          95
Gly Val Phe Pro Tyr Tyr Ser Asn Tyr Ile Asn Leu Ser Lys Leu Glu
          100          105          110
Tyr Glu Leu Leu Ala Arg Tyr Val Pro Gly Gly Ile Ala Pro Ala Arg
          115          120          125
Val Ala Phe Ile Gly Ser Gly Pro Leu Pro Phe Ser Ser Tyr Val Leu
          130          135          140
Ala Ala Arg His Leu Pro Asp Thr Val Phe Asp Asn Tyr Val Pro Val
          145          150          155          160
Arg Ala Ala Asn Asp Arg Ala Thr Arg Leu Phe Arg Ala Asp Lys Asp
          165          170          175
Val Gly Ala Arg Met Ser Phe His Thr Ala Asp Val Ala Asp Leu Thr
          180          185          190
Asp Glu Leu Ala Thr Tyr Asp Val Val Phe Leu Ala Ala Leu Val Gly
          195          200          205
Met Ala Ala Glu Asp Lys Gly Gln Gly Asp Pro His Leu Gly Ala His
          210          215          220
Met Ala Asp Gly Ala Ala Leu Val Arg Ser Ala His Gly Ala Arg Gly
          225          230          235          240
Phe Leu Tyr Pro Ile Val Asp Pro Gln Asp Ile Gly Arg Gly Gly Phe
          245          250          255
Glu Val Leu Ala Val Cys His Pro Asp Asp Asp Val Val Asn Ser Val
          260          265          270
Ile Ile Ala Gln Lys Ser Lys Asp Met Phe Ala Asn Gly Pro Arg Asn
          275          280          285
Gly Cys Gly Gly Arg Tyr Ala Arg Gly Thr Val Pro Val Val Ser Pro
          290          295          300
Pro Cys Arg Phe Gly Glu Met Val Ala Asp Val Thr Gln Lys Arg Glu
          305          310          315          320

```

-continued

Glu Phe Ala Lys Ala Glu Val Ala Phe
325

```

<210> SEQ ID NO 14
<211> LENGTH: 1371
<212> TYPE: DNA
<213> ORGANISM: Hordeum vulgare
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (8)
<223> OTHER INFORMATION: a, c, g, t, unknown or other

<400> SEQUENCE: 14
ggagcggnac gcggtggcga ggtgggcact accgtagtac cgtgcctcag agctcatcac      60
tggtcaggta ccaagaagac ataaaaatgg acgccagag caaggaggtc gacgcccttg      120
tccagaagat caccggcctc cagcgcgcca tcgccaagct gccctcgctc agcccgtccc      180
cggacgtcga cgcgctcttc accgacctgg tcaccgctg cgtgcccccg agcccgtgg      240
acgtgaccaa gctcgccccg gaggcgcagg cgatcgggga gggcctcctc cgcctctgct      300
ccgaggccga gggcaagctg gaggcgcact actccgacat gctcgccgcc ttcgacaacc      360
cgctcgacca cctcggcgtc tccccctact acagcaacta catcaacctc agcaagctcg      420
agtacgagct cctcgcgcgc tacgtgcccg gcggcatcgc cccggcccgc gtcgcttca      480
tcggctccgg cccgctcccg ttcagctcct acgtcctcgc cgcgcgccac ctgcccgaca      540
ccgtgttcga caactacgta cctgtgcgcg cggccaacga ccgcgcgacc aggctgttcc      600
gcgcggacaa ggacgtcggc gcccgcatgt cgttccacac cgccgacgtc gcggacctca      660
ccgacgagct cgctacgtac gacgtcgtct tcttgcccgc gctcgtgggc atggccgccc      720
aggacaaggg ccaaggtgat ccgcaccttg gcgcgcacat ggcgacggg gcggccctcg      780
tccgcagcgc gcacggggcg cgtgggttcc tctaccgat cgctgatccc caagacattg      840
gtcgaggcgg gttcgaggtg ctgcgcgtgt gtcaccccga cgacgacgtg gtgaactccg      900
tcatcatcgc gcagaagtct aaggacatgt ttgccaatgg acctcgcaac ggggtgtggtg      960
gacggtacgc gcgaggcacg gtgccggtgg tcagcccgcc ctgcaggttc ggcgagatgg      1020
tggcagacgt gaccagaag agagaggagt ttgccaaggc ggaagtggcc ttctgattgc      1080
tgcgaggta ccatccgtat gccgctgcta ctttcaata tcttgcaatc gtaggtggcg      1140
atcttctac tcttgttacg acctttcaaa tcatatgttg tttgtaccca ataagtgaag      1200
tgtgttgctt acacgcgcac gtcttgta ca ctcggtctct agaaggcagg gcagatcaag      1260
agactgtgca aaggaaaaga aatgtgtgtt gttgtaggtg tatgagttgg gagtaagatg      1320
attctagttc acaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa a      1371
    
```

```

<210> SEQ ID NO 15
<211> LENGTH: 324
<212> TYPE: PRT
<213> ORGANISM: Oryza sativa
    
```

```

<400> SEQUENCE: 15
Met Glu Ala Gln Asn Gln Glu Val Ala Ala Leu Val Glu Lys Ala Gly
 1           5           10           15
Leu His Ala Ala Ser Lys Leu Pro Ser Leu Ser Pro Ser Ala Glu Val
           20           25           30
Asp Ala Leu Phe Thr Asp Leu Val Thr Ala Cys Val Pro Ala Ser Pro
 35           40           45
    
```

-continued

Val	Asp	Val	Ala	Lys	Leu	Gly	Pro	Glu	Ala	Gln	Ala	Met	Arg	Glu	Glu
	50					55					60				
Leu	Arg	Leu	Cys	Ser	Ala	Ala	Glu	Gly	His	Leu	Glu	Ala	His	Tyr	Ala
	65				70					75					80
Asp	Met	Leu	Ala	Ala	Phe	Asp	Asn	Pro	Leu	Asp	His	Leu	Ala	Arg	Phe
				85					90					95	
Pro	Tyr	Tyr	Gly	Asn	Tyr	Val	Asn	Leu	Ser	Lys	Leu	Glu	Tyr	Asp	Leu
			100					105					110		
Leu	Val	Arg	Tyr	Val	Pro	Gly	Ala	Pro	Thr	Arg	Val	Ala	Phe	Val	Gly
		115					120					125			
Ser	Gly	Pro	Leu	Pro	Phe	Ser	Ser	Leu	Val	Leu	Ala	Ala	His	His	Leu
	130					135					140				
Pro	Asp	Ala	Val	Phe	Asp	Asn	Tyr	Asp	Arg	Cys	Gly	Ala	Ala	Asn	Glu
	145				150					155					160
Arg	Ala	Arg	Arg	Leu	Phe	Arg	Gly	Ala	Asp	Glu	Gly	Leu	Gly	Ala	Arg
				165					170						175
Met	Ala	Phe	His	Thr	Ala	Asp	Val	Ala	Thr	Leu	Thr	Gly	Glu	Leu	Gly
			180					185					190		
Ala	Tyr	Asp	Val	Val	Phe	Leu	Ala	Ala	Leu	Val	Gly	Met	Ala	Ala	Glu
		195					200					205			
Glu	Lys	Ala	Gly	Val	Ala	His	Leu	Gly	Ala	His	Met	Ala	Asp	Gly	Ala
	210					215					220				
Ala	Leu	Val	Val	Arg	Thr	Ala	His	Gly	Ala	Arg	Gly	Phe	Leu	Tyr	Pro
	225				230					235					240
Val	Asp	Pro	Glu	Asp	Val	Arg	Arg	Gly	Gly	Phe	Asp	Val	Leu	Ala	Val
			245						250					255	
Cys	His	Pro	Glu	Asp	Glu	Val	Asn	Ser	Val	Val	Ala	Arg	Lys	Val	Gly
		260						265					270		
Ala	Ala	Ala	Ala	Ala	Ala	Ala	Ala	Arg	Arg	Asp	Glu	Leu	Ala	Asp	Ser
		275					280					285			
Arg	Gly	Val	Val	Leu	Pro	Val	Val	Gly	Pro	Pro	Ser	Thr	Cys	Cys	Lys
	290					295					300				
Val	Glu	Ala	Ser	Ala	Val	Glu	Lys	Ala	Glu	Glu	Phe	Ala	Ala	Asn	Lys
	305				310					315					320

Glu Leu Ser Val

<210> SEQ ID NO 16

<211> LENGTH: 1372

<212> TYPE: DNA

<213> ORGANISM: Oryza sativa

<400> SEQUENCE: 16

ctccatttgg ttgtcatttt caactataat ccaccacaac tcgtgcaaca tcagctcact	60
cgtgttccca accgogacaa agcttcacag atggaggctc agaaccaaga ggtcgctgcc	120
ctggtcgaga agatcgccgg cctccacgcc gccatctcca agctgccgctc gctgagccca	180
tccgcccagg tggacgcgct cttcacccgac ctcgtcacgg cgtgcgtccc ggcgagcccc	240
gtcgacgtgg ccaagctcgg cccggaggcg caggcgatgc gggaggagct catccgctc	300
tgctccgccc ccgagggcca cctcgaggcg cactacgccg acatgctcgc cgccttegac	360
aaccgctcg accaactcgc ccgcttcccg tactacggca actacgtcaa cctgagcaag	420
ctggagtacg acctoctcgt ccgctacgct cccggcattg cccccaccg cgtcgccttc	480
gtcgggtcgg gcccgctgcc gttcagctcc ctcgctcctg ctgcgcacca cctgcccggac	540

-continued

```

gcggtgttcc acaactacga ccggtgcggc gcggccaacg agcgggagag gaggtgttcc 600
cgcggcggcc acgagggcct cgcgcgcgcc atggcggtcc acaccgccga cgtggcgacc 660
ctgacggggg agctcgcgcc gtacgacgtc gtgttcctgg cggcgctcgt gggcatggcg 720
gccgaggaga agggccgggt gatcgcgcac ctgggcgccc acatggcgga cggcgcgggc 780
ctcgtcgtgc ggacggcgca cggggcgccc gggttcctgt acccgatcgt cgatcccag 840
gacgtcagcc gtggcggggt cgacgttctg gcggtgtgcc acccgagga cgaggtgatc 900
aactccgtca tcgtcgcccg caaggtcggg gccgcggccc ccgcccggc ggcgcgcaga 960
gacgagctcg cggactcgcg cggcgtggtt ctgccggtgg tcgggcccgc gtccacgtgc 1020
tgcaaggtgg agggcagcgc ggttgagaag gcagaagagt ttgccgcaa caaggagctg 1080
tccgtctaac agcccggacg tcgaaaggcg cactatatta tggcaataaa tcatttgatt 1140
atacttatgc tgcatttgcg aagctaaggt atactatgca agccatatgt ttgtgttcgt 1200
acgtgttgtt tgggacgtac agttgtgttg ttgtacgtcg tgaagtactg aagtgttcac 1260
agtagatcac aagttcacag caatcaatga ggaccctgta agccagtgta aacgaggaac 1320
atgccatctg tgtatgacag tgagaaatta tataagaaaa acattttgtg ac 1372

```

<210> SEQ ID NO 17

<211> LENGTH: 320

<212> TYPE: PRT

<213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 17

```

Met Ala Cys Gln Asn Asn Leu Val Val Lys Gln Ile Ile Asp Leu Tyr
  1             5             10            15
Asp Gln Ile Ser Lys Leu Lys Ser Leu Lys Pro Ser Lys Asn Val Asp
          20            25            30
Thr Leu Phe Gly Gln Leu Val Ser Thr Cys Leu Pro Thr Asp Thr Asn
          35            40            45
Ile Asp Val Thr Asn Met Cys Glu Glu Val Lys Asp Met Arg Ala Asn
          50            55            60
Leu Ile Lys Leu Cys Gly Glu Ala Glu Gly Tyr Leu Glu Gln His Phe
          65            70            75            80
Ser Thr Ile Leu Gly Ser Leu Gln Glu Asp Gln Asn Pro Leu Asp His
          85            90            95
Leu His Ile Phe Pro Tyr Tyr Ser Asn Tyr Leu Lys Leu Gly Lys Leu
          100           105           110
Glu Phe Asp Leu Leu Ser Gln His Ser Ser His Val Pro Thr Lys Ile
          115           120           125
Ala Phe Val Gly Ser Gly Pro Met Pro Leu Thr Ser Ile Val Leu Ala
          130           135           140
Lys Phe His Leu Pro Asn Thr Thr Phe His Asn Phe Asp Ile Asp Ser
          145           150           155           160
His Ala Asn Thr Leu Ala Ser Asn Leu Val Ser Arg Asp Pro Asp Leu
          165           170           175
Ser Lys Arg Met Ile Phe His Thr Thr Asp Val Leu Asn Ala Thr Glu
          180           185           190
Ala Leu Asp Gln Tyr Asp Val Val Phe Leu Ala Ala Leu Val Gly Met
          195           200           205
Asp Lys Glu Ser Lys Val Lys Ala Ile Glu His Leu Glu Lys His Met
          210           215           220
Ala Pro Gly Ala Val Leu Met Leu Arg Arg Ala His Ala Leu Arg Ala

```


-continued

225		230		235		240									
Phe	Leu	Tyr	Pro	Ile	Val	Asp	Ser	Ser	Asp	Leu	Lys	Gly	Phe	Gln	Leu
				245					250					255	
Leu	Thr	Ile	Tyr	His	Pro	Thr	Asp	Asp	Val	Val	Asn	Ser	Val	Val	Ile
			260					265					270		
Ala	Arg	Lys	Leu	Gly	Gly	Pro	Thr	Thr	Pro	Gly	Val	Asn	Gly	Thr	Arg
		275						280					285		
Gly	Cys	Met	Phe	Met	Pro	Cys	Asn	Cys	Ser	Lys	Ile	His	Ala	Ile	Met
	290					295					300				
Asn	Asn	Arg	Gly	Lys	Lys	Asn	Met	Ile	Glu	Glu	Phe	Ser	Thr	Ile	Glu
305					310					315					320

<210> SEQ ID NO 18
 <211> LENGTH: 963
 <212> TYPE: DNA
 <213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 18

```

atggcttgcc aaaacaatct cgttgtgaag caaatcatcg acttgtacga ccaaactctca    60
aagctcaaga gcttaaaacc ttccaaaaat gtcgacactt tgttcggaca actcgtgtcc    120
acgtgcttac ccacggatag aaacatcgat gtcacaaata tgtgtgaaga agtcaaagac    180
atgagagcta atctcatcaa gctttgtggt gaagccgaag gttatttggg gcaacacttc    240
tccacaattt tgggatcttt acaagaagac caaaaccac ttgaccattt acacatcttt    300
ccttactact ccaactacct caagctaggc aagctcgagt tcgatctcct gagccaacac    360
tcaagccatg tccccaccaa gattgccttc gtgggttcgg gtccgatgcc tctcacatcc    420
atcgtattgg ccaagtttca cctccccaac acgacgttcc acaactttga catcgactca    480
cacgcaaaca cactcgcttc aaacctcgtc tctcgcgacc cggacctctc aaaacgcatg    540
atcttcacca caacggagct actaaacgca accgaagccc ttgaccaata tgacgtcgtt    600
ttcttagcgg cgctttagg gatggacaaa gagtcaaagg tcaaagccat cgagcacttg    660
gagaaacaca tggctcctgg agctgttctt atgctaagga gggctcatgc tctcagagct    720
ttcttataat caatcgttga ctcgctgat ctcaaaggct ttcaactctt gacctctat    780
catccaaccg atgaogtggg taactcgggt gtgatgcac gtaagctcgg tggctccgacc    840
acgcccgggg ttaatgttac tcgtggatgc atgtttatgc cttgtaactg ctccaagatt    900
cacgcatgca tgaacaaccg tggtaagaag aatatgatcg aggagttagg taccatcgag    960
taa
    
```

<210> SEQ ID NO 19
 <211> LENGTH: 320
 <212> TYPE: PRT
 <213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 19

Met	Ala	Cys	Gln	Asn	Asn	Leu	Val	Val	Lys	Gln	Ile	Met	Asp	Leu	Tyr
1				5					10					15	
Asn	Gln	Ile	Ser	Asn	Leu	Glu	Ser	Leu	Lys	Pro	Ser	Lys	Asn	Val	Asp
			20					25					30		
Thr	Leu	Phe	Arg	Gln	Leu	Val	Ser	Thr	Cys	Leu	Pro	Thr	Asp	Thr	Asn
		35					40					45			
Ile	Asp	Val	Thr	Glu	Ile	His	Asp	Glu	Lys	Val	Lys	Asp	Met	Arg	Ser
	50					55					60				

-continued

His	Leu	Ile	Lys	Leu	Cys	Gly	Glu	Ala	Glu	Gly	Tyr	Leu	Glu	Gln	His
65					70					75					80
Phe	Ser	Ala	Ile	Leu	Gly	Ser	Phe	Glu	Asp	Asn	Pro	Leu	Asn	His	Leu
				85					90					95	
His	Ile	Phe	Pro	Tyr	Tyr	Asn	Asn	Tyr	Leu	Lys	Leu	Gly	Lys	Leu	Glu
			100					105					110		
Phe	Asp	Leu	Leu	Ser	Gln	His	Thr	Thr	His	Val	Pro	Thr	Lys	Val	Ala
		115					120					125			
Phe	Ile	Gly	Ser	Gly	Pro	Met	Pro	Leu	Thr	Ser	Ile	Val	Leu	Ala	Lys
		130				135					140				
Phe	His	Leu	Pro	Asn	Thr	Thr	Phe	His	Asn	Phe	Asp	Ile	Asp	Ser	His
145					150					155					160
Ala	Asn	Thr	Leu	Ala	Ser	Asn	Leu	Val	Ser	Arg	Asp	Ser	Asp	Leu	Ser
				165						170				175	
Lys	Arg	Met	Ile	Phe	His	Thr	Thr	Asp	Val	Leu	Asn	Ala	Lys	Glu	Gly
			180					185						190	
Leu	Asp	Gln	Tyr	Asp	Val	Val	Phe	Leu	Ala	Ala	Leu	Val	Gly	Met	Asp
		195					200					205			
Lys	Glu	Ser	Lys	Val	Lys	Ala	Ile	Glu	His	Leu	Glu	Lys	His	Met	Ala
	210					215					220				
Pro	Gly	Ala	Val	Val	Met	Leu	Arg	Ser	Ala	His	Gly	Leu	Arg	Ala	Phe
225					230					235					240
Leu	Tyr	Pro	Ile	Val	Asp	Ser	Cys	Asp	Leu	Lys	Gly	Phe	Glu	Val	Leu
				245					250					255	
Thr	Ile	Tyr	His	Pro	Ser	Asp	Asp	Val	Val	Asn	Ser	Val	Val	Ile	Ala
			260					265						270	
Arg	Lys	Leu	Gly	Gly	Ser	Asn	Gly	Ala	Arg	Gly	Ser	Gln	Ile	Gly	Arg
		275					280					285			
Cys	Val	Val	Met	Pro	Cys	Asn	Cys	Ser	Lys	Val	His	Ala	Ile	Leu	Asn
	290					295					300				
Asn	Arg	Gly	Met	Glu	Lys	Asn	Leu	Ile	Glu	Glu	Phe	Ser	Ala	Ile	Glu
305					310					315					320

<210> SEQ ID NO 20

<211> LENGTH: 963

<212> TYPE: DNA

<213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 20

atggccttgcc aaaacaatct cgttgtgaag caaatcatgg acttatacaa ccaaattcca	60
aaactogaga gcttaaaacc atccaagaat gtcgacactt tgttcagaca acttgtgtcc	120
acgtgcttac caacggacac gaacatcgat gtcacagaga tacacgatga aaaagtcaaa	180
gacatgagat ctcatctcat caagctttgt ggtgaagccg aaggttattt agagcaacac	240
ttttcagcaa tcttaggtct ttttgaagac aaccctctaa accatttaca catcttcccc	300
tattacaaca actatctcaa actaggcaaa ctggaattcg atctcttttc tcagcacaca	360
acctatgtcc cgaccaaagt cgcctttatt ggttcoggtc cgatgccact tacttccatc	420
gtcttggtcca agttocacct ccccaacaca acgttocaca acttogacat cgactcacac	480
gccaaacacac tcgcttcaaa cctogtttct cgtgattctg acctttccaa acgcatgatt	540
ttccacacaa ctgatgtatt aaacgctaag gaggggtag accaatacga tgttgttttc	600
ttggcagctc ttgttggtgat ggataaagag tcaaaggta aagctattga gcatttagag	660
aagcatatgg cccctggagc tgtggtgatg ctaagaagtg ctcatggtct tagagctttc	720

-continued

```

ttgtatccaa tcgttgactc ttgtgatctt aaagggtttg aggtggtaac catttatcat 780
ccgtctgacg acgtggttaa ttcggtggtc atcgcacgta agcttggtgg ttcaaatgga 840
gctcgaggca gccagatcgg acggtgtgtg gttatgcott gtaattgctc taaggccac 900
gcgatcttga acaatcgtgg tatggagaag aatttgatcg aggagtttag tgccatcgag 960
taa 963

```

```

<210> SEQ ID NO 21
<211> LENGTH: 320
<212> TYPE: PRT
<213> ORGANISM: Arabidopsis thaliana

```

```

<400> SEQUENCE: 21

```

```

Met Gly Cys Gln Asp Glu Gln Leu Val Gln Thr Ile Cys Asp Leu Tyr
 1          5          10          15
Glu Lys Ile Ser Lys Leu Glu Ser Leu Lys Pro Ser Glu Asp Val Asn
 20          25          30
Ile Leu Phe Lys Gln Leu Val Ser Thr Cys Ile Pro Pro Asn Pro Asn
 35          40          45
Ile Asp Val Thr Lys Met Cys Asp Arg Val Gln Glu Ile Arg Leu Asn
 50          55          60
Leu Ile Lys Ile Cys Gly Leu Ala Glu Gly His Leu Glu Asn His Phe
 65          70          75          80
Ser Ser Ile Leu Thr Ser Tyr Gln Asp Asn Pro Leu His His Leu Asn
 85          90          95
Ile Phe Pro Tyr Tyr Asn Asn Tyr Leu Lys Leu Gly Lys Leu Glu Phe
100          105          110
Asp Leu Leu Glu Gln Asn Leu Asn Gly Phe Val Pro Lys Ser Val Ala
115          120          125
Phe Ile Gly Ser Gly Pro Leu Pro Leu Thr Ser Ile Val Leu Ala Ser
130          135          140
Phe His Leu Lys Asp Thr Ile Phe His Asn Phe Asp Ile Asp Pro Ser
145          150          155          160
Ala Asn Ser Leu Ala Ser Leu Leu Val Ser Ser Asp Pro Asp Ile Ser
165          170          175
Gln Arg Met Phe Phe His Thr Val Asp Ile Met Asp Val Thr Glu Ser
180          185          190
Leu Lys Ser Phe Asp Val Val Phe Leu Ala Ala Leu Val Gly Met Asn
195          200          205
Lys Glu Glu Lys Val Lys Val Ile Glu His Leu Gln Lys His Met Ala
210          215          220
Pro Gly Ala Val Leu Met Leu Arg Ser Ala His Gly Pro Arg Ala Phe
225          230          235          240
Leu Tyr Pro Ile Val Glu Pro Cys Asp Leu Gln Gly Phe Glu Val Leu
245          250          255
Ser Ile Tyr His Pro Thr Asp Asp Val Ile Asn Ser Val Val Ile Ser
260          265          270
Lys Lys His Pro Val Val Ser Ile Gly Asn Val Gly Gly Pro Asn Ser
275          280          285
Cys Leu Leu Lys Pro Cys Asn Cys Ser Lys Thr His Ala Lys Met Asn
290          295          300
Lys Asn Met Met Ile Glu Glu Phe Gly Ala Arg Glu Glu Gln Leu Ser
305          310          315          320

```

-continued

<210> SEQ ID NO 22
 <211> LENGTH: 963
 <212> TYPE: DNA
 <213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 22

```

atgggttgcc aagacgaaca attggtgcaa acaatatgcg atctctacga aaagatctca    60
aagcttgaga gtctaaaacc atccgaagat gtcaacattc tcttcaagca gctcgtttcc    120
acatgcatac caccaaaacc taacatcgat gtcaccaaga tgtgtgacag agtccaagag    180
attcgactta atctcatcaa gatttggtgt ctagccgaag gtcacttaga aaaccatttc    240
tcttcgatct tgacctctta ccaagacaac ccacttcacg atttaaacad tttcccttat    300
tacaacaact attgaaact cggaaagctc gagttcgacc tcctcgaaca aaacctaaat    360
ggctttgtcc caaagagtgt ggctttcatt ggatctggtc ctcttcctct cacttccatc    420
gttcttgctt cattccatct caaagacaca atctttcaca actttgacat cgaccatca    480
gcgaactcac tcgcttctct tctggtttcc tctgatccag acatctctca acgcatgttc    540
ttccacaccg ttgatataat ggacgtgaca gagagcttaa agagcttga tgtcgtgttt    600
ctagctgctc ttgttggaat gaacaaagag gagaaagtta aagtgatcga gcatctgcag    660
aaacacatgg ctctctggtc tgtgctcatg cttaggagtg ctcatggtcc gagagcgttt    720
ctttatccga tcggtgagcc gtgtgatctt caggggttcg aggttttgtc tatttatcac    780
ccaacagatg atgttatcaa ctccgtggtg atctctaaaa agcatccagt tgtttcaatt    840
gggaatgttg gtggtcctaa ttcagcttg ctcaagcctt gcaactgttc caagaccac    900
gcgaaaatga acaagaacat gatgatcgag gagttcggag ctagggagga acagttgtct    960
taa
  
```

<210> SEQ ID NO 23
 <211> LENGTH: 32
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 consensus sequence
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (3)
 <223> OTHER INFORMATION: Variable amino acid residue
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (9)
 <223> OTHER INFORMATION: Variable amino acid residue
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (16)
 <223> OTHER INFORMATION: Variable amino acid residue
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (19)
 <223> OTHER INFORMATION: Variable amino acid residue
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (24)
 <223> OTHER INFORMATION: Variable amino acid residue

<400> SEQUENCE: 23

```

Leu Pro Xaa Leu Ser Pro Ser Pro Xaa Val Asp Arg Leu Phe Thr Xaa
  1             5             10             15
Leu Val Xaa Ala Cys Val Pro Xaa Ser Pro Val Asp Val Thr Lys Leu
  20             25             30
  
```

-continued

<210> SEQ ID NO 24
 <211> LENGTH: 16
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 consensus sequence
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (7)
 <223> OTHER INFORMATION: Variable amino acid residue
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (11)
 <223> OTHER INFORMATION: Variable amino acid residue
 <400> SEQUENCE: 24

Leu Ile Arg Leu Cys Ser Xaa Ala Glu Gly Xaa Leu Glu Ala His Tyr
 1 5 10 15

<210> SEQ ID NO 25
 <211> LENGTH: 10
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 consensus sequence
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (7)
 <223> OTHER INFORMATION: Variable amino acid residue
 <400> SEQUENCE: 25

Pro Leu Asp His Leu Gly Xaa Phe Pro Tyr
 1 5 10

<210> SEQ ID NO 26
 <211> LENGTH: 13
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 consensus sequence
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (4)
 <223> OTHER INFORMATION: Variable amino acid residue
 <400> SEQUENCE: 26

Val Ala Phe Xaa Gly Ser Gly Pro Leu Pro Phe Ser Ser
 1 5 10

<210> SEQ ID NO 27
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 consensus sequence
 <400> SEQUENCE: 27

Asp Val Val Phe Leu Ala Ala Leu Val Gly Met
 1 5 10

<210> SEQ ID NO 28
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

```

consensus sequence
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)
<223> OTHER INFORMATION: Variable amino acid residue

<400> SEQUENCE: 28

Arg Gly Gly Phe Xaa Val Leu Ala Val Xaa His Pro
  1             5             10

<210> SEQ ID NO 29
<211> LENGTH: 328
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      amino acid sequence of nicotianamine synthase
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(10)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (12)..(13)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (16)..(21)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (23)..(24)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (26)..(27)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (29)..(29)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (31)..(36)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (39)..(40)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (43)..(44)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (46)..(46)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (48)..(51)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (54)..(60)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (62)..(62)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (65)..(66)
<223> OTHER INFORMATION: Variable amino acid residue

```

-continued

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (69)..(70)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (72)..(73)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (76)..(77)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (80)..(80)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (82)..(85)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (87)..(89)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (91)..(91)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (94)..(94)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (97)..(98)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (102)..(103)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (106)..(107)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (109)..(110)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (113)..(114)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (117)..(128)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (131)..(131)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (138)..(139)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (141)..(142)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (145)..(146)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (149)..(152)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (154)..(154)

-continued

<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (156)..(161)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (164)..(165)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (167)..(168)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (170)..(179)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (182)..(182)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (184)..(184)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (186)..(194)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (196)..(198)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (206)..(206)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (210)..(213)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (215)..(219)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (222)..(223)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (226)..(227)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (230)..(230)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (232)..(238)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (240)..(240)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (245)..(245)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (247)..(254)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (257)..(257)
<223> OTHER INFORMATION: Variable amino acid residue
<220> FEATURE:
<221> NAME/KEY: MOD_RES

-continued

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 290 295 300

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 305 310 315 320

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 325

<210> SEQ ID NO 30
 <211> LENGTH: 19
 <212> TYPE: PRT
 <213> ORGANISM: Hordeum vulgare

<400> SEQUENCE: 30

Asp Ala Gln Asn Lys Glu Val Ala Ala Leu Ile Glu Lys Ile Ala Gly
 1 5 10 15

Ile Gln Ala

<210> SEQ ID NO 31
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: Hordeum vulgare

<400> SEQUENCE: 31

Arg Glu Ala Leu Ile Arg Leu
 1 5

<210> SEQ ID NO 32
 <211> LENGTH: 71
 <212> TYPE: PRT
 <213> ORGANISM: Oryza sativa

<400> SEQUENCE: 32

Met Glu Ala Gln Asn Gln Glu Val Ala Ala Leu Val Glu Lys Ile Ala
 1 5 10 15

Gly Leu His Ala Ala Ile Ser Lys Leu Pro Ser Leu Ser Pro Ser Ala
 20 25 30

Glu Val Asp Ala Leu Phe Thr Asp Leu Val Thr Ala Cys Val Pro Ala
 35 40 45

Ser Pro Val Asp Val Ala Lys Leu Gly Pro Glu Ala Gln Ala Met Arg
 50 55 60

Glu Glu Leu Ile Arg Leu Cys
 65 70

<210> SEQ ID NO 33
 <211> LENGTH: 111
 <212> TYPE: PRT
 <213> ORGANISM: Oryza sativa

<400> SEQUENCE: 33

Tyr Val Asn Leu Ser Lys Leu Glu Tyr Asp Leu Leu Val Arg Tyr Val
 1 5 10 15

Pro Gly Ile Ala Pro Thr Arg Val Ala Phe Val Gly Ser Gly Pro Leu
 20 25 30

Pro Phe Ser Ser Leu Val Leu Ala Ala His His Leu Pro Asp Ala Val
 35 40 45

Phe Asp Asn Tyr Asp Arg Cys Gly Ala Ala Asn Glu Arg Ala Arg Arg
 50 55 60

Leu Phe Arg Gly Ala Asp Glu Gly Leu Gly Ala Arg Met Ala Phe His
 65 70 75 80

65

The invention claimed is:

1. An isolated or purified enzyme exhibiting nicotianamine synthase activity, wherein the enzyme comprises a polypeptide that is at least 90% identical to SEQ ID NO:1.

2. The isolated or purified enzyme of claim 1, wherein the enzyme comprises the polypeptide having the amino acid sequence of SEQ ID NO: 1.

3. The enzyme according to claim 1, wherein the enzyme comprises the consensus amino acid sequence of ₁₉₉DVVFLAALVGM₂₀₉ (SEQ ID NO: 27).

4. The enzyme of claim 1, wherein the polypeptide further comprises all of the conserved amino acid residues of SEQ ID NO: 1 that is:

L(11), K(14), I(15), I(22), L(25), L(28), P(30), L(37), F(38), L(41), V(42), C(45), P(47), D(52), V(53), Q(61), M(63), R(64), L(67), I(68), C(71), A(74), E(75), L(78), E(79), H(81), L(86), D(90), P(92), L(93), H(95), L(96), F(99), P(100), Y(101), N(104), Y(105), L(108), L(111), E(112), L(115), L(116), A(129), F(130), G(132), S(133), G(134), P(135), L(136), P(137), S(140), L(143), A(144), H(147), L(148), F(153), N(155), A(162), N(163), A(166), L(169), R(180), M(181), F(183), T(185), L(195), D(199), V(200), V(201), F(202), L(203), A(204), A(205), V(207), G(208), M(209), K(214), H(220), L(221), H(224), M(225), G(228), A(229), L(231), R(239), F(241), L(242), Y(243), P(244), V(246), G(255), F(256), V(258), L(259), V(261), H(263), P(264), V(268), N(270), S(271), K(277) (SEQ ID NO:29).

5. The enzyme of claim 1, wherein the polypeptide has more than 95% identity with the amino acid sequence of SEQ ID NO: 1.

6. The enzyme of claim 1, wherein the nicotianamine synthase activity is measured in an assay in a comparison with the enzyme of SEQ ID NO:1.

7. The enzyme of claim 1, wherein the enzyme is isolated or purified from barley.

8. A mutated enzyme exhibiting nicotianamine synthase activity, wherein the enzyme:

a. is a polypeptide having more than 95% identity with the amino acid sequence of SEQ ID NO: 1, comprising at least one consensus sequence of SEQ ID NO: 1 that is:

(1)₂₅LPXLSPSPXVDRLFTXLVXACVPXPSPVD-VTKL₅₆ (SEQ ID NO: 23)
 (2)₆₇LIRLCSXAEGXLEAHY₈₂ (SEQ ID NO: 24)
 (3)₉₂PLDHLGXFPY₁₀₁ (SEQ ID NO: 25)
 (4)₁₂₈VAFXGSGPLPFSS₁₄₀ (SEQ ID NO: 26)
 (5)₁₉₉DVVFLAALVGM₂₀₉ (SEQ ID NO: 27)
 (6)₂₅₃RGGFXVLAVXHP₂₆₄ (SEQ ID NO: 28); and

b. has more than 25% of the relative nicotianamine synthase activity of the enzyme of SEQ ID NO: 1.

9. The enzyme of claim 8, wherein the polypeptide further comprises all of the conserved amino acid residues of SEQ ID NO: 1 that is:

L(11), K(14), I(15), I(22), L(25), L(28), P(30), L(37), F(38), L(41), V(42), C(45), P(47), D(52), V(53), Q(61), M(63), R(64), L(67), I(68), C(71), A(74), E(75), L(78), E(79), H(81), L(86), D(90), P(92), L(93), H(95), L(96), F(99), P(100), Y(101), N(104), Y(105), L(108), L(111), E(112), L(115), L(116), A(129), F(130), G(132), S(133), G(134), P(135), L(136), P(137), S(140), L(143), A(144), H(147), L(148), F(153), N(155), A(162), N(163), A(166), L(169), R(180), M(181), F(183), T(185), L(195), D(199), V(200), V(201), F(202), L(203), A(204), A(205), V(207), G(208), M(209), K(214), H(220), L(221), H(224), M(225), G(228), A(229), L(231), R(239), F(241), L(242),

66

Y(243), P(244), V(246), G(255), F(256), V(258), L(259), V(261), H(263), P(264), V(268), N(270), S(271), K(277) (SEQ ID NO: 29).

10. The enzyme of claim 8, wherein the nicotianamine synthase activity is measured in an assay in a comparison with the enzyme of SEQ ID NO:1.

11. The enzyme of claim 8, wherein the polypeptide has more than 97% identity with the amino acid sequence of SEQ ID NO: 1.

12. The enzyme of claim 1, wherein the polypeptide has more than 97% identity with the amino acid sequence of SEQ ID NO: 1.

13. An isolated, purified, or mutated enzyme exhibiting nicotianamine synthase activity, wherein the enzyme comprises an active fragment of the amino acid sequence of SEQ ID NO: 1, the active fragment comprising a polypeptide, wherein the polypeptide:

a. comprises at least one consensus sequence of SEQ ID NO: 1 that is:

(1)₂₅LPXLSPSPXVDRLFTXLVXACVPXPSPVD-VTKL₅₆ (SEQ ID NO: 23)
 (2)₆₇LIRLCSXAEGXLEAHY₈₂ (SEQ ID NO: 24)
 (3)₉₂PLDHLGXFPY₁₀₁ (SEQ ID NO: 25)
 (4)₁₂₈VAFXGSGPLPFSS₁₄₀ (SEQ ID NO: 26)
 (5)₁₉₉DVVFLAALVGM₂₀₉ (SEQ ID NO: 27)
 (6)₂₅₃RGGFXVLAVXHP₂₆₄ (SEQ ID NO: 28); and

b. has more than 25% of the relative nicotianamine synthase activity of the enzyme of SEQ ID NO:1.

14. An isolated or purified barley enzyme exhibiting nicotianamine synthase activity, wherein:

a. the enzyme is:

i. isolated or purified from barley; or
 ii. expressed directly or indirectly from a nucleic acid isolated or purified from barley; or
 iii. expressed directly or indirectly from a chimeric nucleic acid at least partially isolated or purified from barley;

b. the enzyme comprises a polypeptide having at least 90% identity with the amino acid sequence of SEQ ID NO: 1, comprising at least one consensus sequence of SEQ ID NO: 1 that is:

(1)₂₅LPXLSPSPXVDRLFTXLVXACVPXPSPVD-VTKL₅₆ (SEQ ID NO: 23)
 (2)₆₇LIRLCSXAEGXLEAHY₈₂ (SEQ ID NO: 24)
 (3)₉₂PLDHLGXFPY₁₀₁ (SEQ ID NO: 25)
 (4)₁₂₈VAFXGSGPLPFSS₁₄₀ (SEQ ID NO: 26)
 (5)₁₉₉DVVFLAALVGM₂₀₉ (SEQ ID NO: 27)
 (6)₂₅₃RGGFXVLAVXHP₂₆₄ (SEQ ID NO: 28); and

c. the enzyme has more than 25% of the relative nicotianamine synthase activity of the enzyme of SEQ ID NO: 1.

15. The enzyme of claim 14, wherein the polypeptide further comprises all of the conserved amino acid residues of SEQ ID NO: 1 that is:

L(11), K(14), I(15), I(22), L(25), L(28), P(30), L(37), F(38), L(41), V(42), C(45), P(47), D(52), V(53), Q(61), M(63), R(64), L(67), I(68), C(71), A(74), E(75), L(78), E(79), H(81), L(86), D(90), P(92), L(93), H(95), L(96), F(99), P(100), Y(101), N(104), Y(105), L(108), L(111), E(112), L(115), L(116), A(129), F(130), G(132), S(133), G(134), P(135), L(136), P(137), S(140), L(143), A(144), H(147), L(148), F(153), N(155), A(162), N(163), A(166), L(169), R(180), M(181), F(183), T(185), L(195), D(199), V(200), V(201), F(202), L(203), A(204), A(205), V(207), G(208), M(209), K(214), H(220), L(221), H(224), M(225), G(228), A(229), L(231), R(239), F(241), L(242),

67

Y(243), P(244), V(246), G(255), F(256), V(258),
L(259), V(261), H(263), P(264), V(268), N(270),
S(271), K(277) (SEQ ID NO: 29).

16. The enzyme of claim 14, wherein the polypeptide has
more than 90% identity with the amino acid sequence of 5
SEQ ID NO: 1.

17. The enzyme of claim 14, wherein the polypeptide has
more than 95% identity with the amino acid sequence of
SEQ ID NO: 1.

18. The enzyme of claim 14, wherein the nicotianamine 10
synthase activity is measured in an assay in a comparison
with the enzyme of SEQ ID NO:1.

19. An isolated or purified enzyme exhibiting nicotian-
amine synthase activity, wherein the enzyme consists of the
polypeptide set forth as SEQ ID NO: 1.

68

20. The isolated or purified enzyme of claim 1, compris-
ing at least one consensus sequence of SEQ ID NO: 1
selected from the group consisting of:

(1) ²⁵LPXLSPSPXVDRLFTXLVXACVPXSPVD-
VTKL₅₆ (SEQ ID NO: 23)

(2) ⁶⁷LIRLCSXAEGXLEAHY₈₂ (SEQ ID NO: 24)

(3) ⁹²PLDHLGXFY₁₀₁ (SEQ ID NO: 25)

(4) ¹²⁸VAFXGSGPLPFSS₁₄₀ (SEQ ID NO: 26)

(5) ¹⁹⁹DVVFLAALVGM₂₀₉ (SEQ ID NO: 27)

(6) ²⁵³RGGFVXLAVXHP₂₆₄ (SEQ ID NO: 28); and

b. has more than 25% of the relative nicotianamine
synthase activity of the enzyme of SEQ ID NO:1.

* * * * *