



US 20070160993A1

(19) **United States**

(12) **Patent Application Publication** (10) **Pub. No.: US 2007/0160993 A1**

**Watanabe** (43) **Pub. Date: Jul. 12, 2007**

(54) **NOVEL CENTROMERIC PROTEIN SHUGOSHIN**

**Publication Classification**

(75) Inventor: **Yoshinori Watanabe**, Tokyo (JP)

- (51) **Int. Cl.**
  - C12Q* 1/68 (2006.01)
  - C07H* 21/04 (2006.01)
  - C12P* 21/06 (2006.01)
  - C12N* 9/22 (2006.01)
  - C07K* 16/40 (2006.01)
- (52) **U.S. Cl.** ..... **435/6**; 435/69.1; 435/199; 435/320.1; 435/325; 530/388.26; 530/358; 536/23.2

Correspondence Address:  
**MORGAN & FINNEGAN, L.L.P.**  
**3 WORLD FINANCIAL CENTER**  
**NEW YORK, NY 10281-2101 (US)**

(73) Assignee: **JAPAN SCIENCE AND TECHNOLOGY AGENCY**, Saitama (JP)

(57) **ABSTRACT**

The present invention is to provide meiosis-specific novel kinetochore protein Sgo1 (shugoshin) derived from fission yeast *Schizosaccharomyces pombe*, and a homologue or paralogue thereof having a regulatory activity of chromosome segregation; and DNAs encoding them; as a factor ensuring the retention of unidirection and cohesion in sister centromere at meiosis I in cooperation with cohesin. To elucidate the proteins protecting Rec8 during anaphase, the present inventor screened in fission yeast genes for a gene that inhibits mitotic growth and prevents sister chromatid from the separation at anaphase, when co-expressed with Rec8. In this approach, meiosis-specific protein Sgo1 that protects (Shugo) centromeric Rec8 from the degradation at anaphase I was identified. Further, a budding yeast Sgo1 homologue and a fission yeast mitotic paralogue Sgo2 were identified.

(21) Appl. No.: **10/581,158**

(22) PCT Filed: **Nov. 24, 2004**

(86) PCT No.: **PCT/JP04/17428**

§ 371(c)(1),  
(2), (4) Date: **Jan. 30, 2007**

(30) **Foreign Application Priority Data**

Dec. 1, 2003 (JP) ..... 2003-401943  
Sep. 27, 2004 (JP) ..... 2004-279450

Fig.1




Fig.2




Fig.3




Fig.4




Fig.5




Fig.6




Fig.7








Fig.8




Fig.9




Fig.10




Fig.11






Fig.12




Fig.13




Fig.14




Fig.15




Fig.16




Fig.17





Fig.18



**NOVEL CENTROMERIC PROTEIN SHUGOSHIN**

## TECHNICAL FIELD

[0001] The present invention relates to a protector protein Sgo1 (shugoshin) of cohesin Rec8 derived from fission yeast *Schizosaccharomyces pombe*, its homologue and paralogue having a regulatory activity of chromosome segregation, and DNAs encoding them.

## BACKGROUND ART

[0002] In eukaryotes, sister chromatid cohesion is established during S phase of cell cycle and maintained throughout G2 until M phase. During mitosis, this cohesion is destroyed along the entire length of chromosome, allowing sister chromatid to segregate to the opposite sides of cell (equational division) and ensuring that each daughter cell receives one copy of each chromosome. In contrast, meiosis consists of two rounds of chromosome segregation following a single round of DNA replication, leading to the formation of four haploid gametes from one diploid germ cell. During meiosis I, homologous chromosomes (homologues) pair to recombine, forming chiasmata in which one sister chromatid from one homologue is covalently attached to a sister chromatid from the other homologue. Hence, in order for homologues to segregate at meiosis I, cohesion of sister chromatid is necessary to be dissociated along the chromosome arms to resolve chiasmata. However, sister chromatid cohesion is retained at centromere until meiosis II, and utilizes the residual centromeric cohesion when sister chromatid segregates, in the same manner as it does in mitosis. Thus, meiotic division requires sister chromatid cohesion to be dissociated in two steps. However, the molecular mechanism for protection of centromeric cohesion only during meiosis I and only at the centromere has remained to be elucidated (e.g., see *Annu Rev Genet* 35, 673-745(2001)).

[0003] There are important clues as to the molecular nature of sister chromatid cohesion, and the mechanism dissociating sister chromatid cohesion at the onset of anaphase (e.g., see *Annu Rev Genet* 35, 673-745(2001); *Curr Opin Cell Biol* 12, 297-301(2000); *Curr Biol* 13, R104-14(2003); *Annu Rev Cell Dev Biol* 17, 753-77(2001); *Genes Dev* 16, 399-414(2002)). In various eukaryotes, sister chromatid cohesion depends on a multisubunit cohesin complex including Scc1 (Rad21 in fission yeast *Schizosaccharomyces pombe*). Anaphase promoting complex (APC)-dependent degradation of the securin, Cut2/Pds1, allows to dissociate the Cut1/Esp1 endopeptidase (separase), which in turn cleaves Rad21/Scc1, dissociating sister chromatid cohesion. During meiosis, the cohesion subunit Rad21/Scc1 is replaced with a meiotic counterpart, Rec8 (e.g., see *Cell* 98, 91-103(1999); *Mol. Cell. Biol.* 19, 3515-3528(1999); *Nature* 400, 461-4(1999); *Genes Dev* 15, 1349-60(2001); *J Cell Biol* 160, 657-70(2003)). As Rec8 complex resides only at centromere after meiosis I and the depletion of Rec8 destroys centromeric cohesion, the presence of Rec8 at centromere has been thought to confer the persistence of cohesion throughout meiosis I (e.g., see *Nat Cell Biol* 1, E125-7(1999)). Several lines of evidence suggest that Rec8 along chromosome arms is cleaved by separase at anaphase I while centromeric Rec8 is specifically protected until metaphase II (e.g., see *Cell* 103, 387-98(2000); *Embo J* 22, 5643-53(2003)). Budding yeast SPO13 has been implicated

in the protection of centromeric Rec8 (e.g., see *Genes Dev* 16, 1659-71(2002); *Genes Dev* 16, 1672-81(2002)), but SPO13 is not centromeric and may function indirectly. *Drosophila* ME1-S332 is a protein residing at centromere, is required for the persistence of centromeric cohesion during meiosis I, and has features of a candidate protector of meiotic centromeric cohesion, although the details of such protection have so far not been elucidated (e.g., see *Annu Rev Cell Dev Biol* 17, 753-77(2001); *Cell* 83, 247-256(1995)). Despite the completion of genome sequencing projects on several organisms, no homologue of these proteins has emerged, preventing the formulation of a generalized view of the protection. Concurrently, studies in fission yeast have illuminated the importance of pericentromeric heterochromatin for recruiting centromeric Rec8 complexes and ensuring centromeric cohesion during meiosis I (e.g., see *Science* 300, 1152-5(2003)). However, pericentromeric heterochromatin cannot alone confer the specific protection of Rec8 at meiosis I toward meiosis II.

## DISCLOSURE OF THE INVENTION

[0004] Almost all the eukaryotes including human propagate offsprings by sexual reproduction evolutionarily predominant with a mixture of genome. Meiosis that reduces chromosome number in half is a core part of the sexual reproduction mechanism. In somatic mitosis, two kinetochores of sister chromatid are caught by spindle microtubule extended from the opposite poles, and sister chromatid is evenly segregated to the both poles by concurrently dissolving the cohesion of arms and centromeres (equational division). In contrast, in meiosis I kinetochores of sister chromatids are caught by spindle microtubule extended from the same pole, and segregated to the same pole while retaining the cohesion at centromere (meiotic division). Next, for the first time in meiosis II the cohesion of centromere site of sister chromatid is dissolved, and separated toward one pole or the other of the two poles respectively, which culminates in the generation of accurate four haploid gametes. Meiosis-specific meiotic division is a modality of chromosome segregation conserved in almost all the eukaryotes, from yeast to human, however regulatory mechanism at the molecular level has remained enigmatic for a long time. The present inventor has demonstrated that meiosis-specific chromosome cohesion factor, cohesin plays an essential role in this regulation by using fission yeast (*Nature* 400, 461-4(1999); *Science* 300, 1152-5(2003); *Nature* 409, 359-363(2001)). An object of the present invention is to provide meiosis-specific novel kinetochore protein Sgo1 (shugoshin) derived from fission yeast *Schizosaccharomyces pombe*, and a homologue or paralogue thereof having a regulatory activity of chromosome segregation; and DNAs encoding them; as a factor ensuring the retention of unidirection and cohesion in sister centromere at meiosis I in cooperation with cohesin.

[0005] Meiosis comprises two steps of specialized nuclear divisions for producing haploid gametes. To accomplish this, sister chromatid cohesion is necessary to be dissociated in a stepwise manner, first from chromosome arms at anaphase I and second from centromeres at anaphase II. In particular, the factors that protect centromeric cohesion during meiosis I have heretofore remained undissolved. To elucidate the proteins protecting Rec8 during anaphase, the present inventor screened in fission yeast genes for a gene that inhibits mitotic growth and prevents sister chromatid

from the separation at anaphase, when co-expressed with Rec8. In this approach, meiosis-specific protein that is a protector of Rec8 in fission yeast and protects (Shugo) centromeric Rec8 from the degradation at anaphase I was indentified, and named Sgo1 (Shugoshin, a Japanese for “guardian spirit”). It was also identified that shugoshin plays an important role in mitotic chromosome segregation, and then identified a budding yeast Sgo1 homologue and a fission yeast mitotic paralogue Sgo2. A marginal similarity between Sgo1 and *Drosophila* ME1-S332 was identified, and Sgo1 homologue in other eukaryotes was also identified. Shugoshin-like proteins in animal cells, which were predicted from the sequence, also have functional conservation with yeast shugoshin. The present invention has been thus completed based on this knowledge.

[0006] That is, the present invention relates to (1) a DNA encoding a following protein (a) or (b): (a) a protein consisting of an amino acid sequence shown in SEQ ID NO: 2, (b) a protein comprising an amino acid sequence where one or several amino acids are deleted, replaced or added in an amino acid sequence shown in SEQ ID NO: 2, and having a regulatory activity of chromosome segregation; (2) a DNA consisting of a base sequence shown in SEQ ID NO: 1 or a complementary sequence thereof; (3) a DNA containing part or whole of a base sequence shown in SEQ ID NO: 1 or a complementary sequence thereof, and encoding a protein that has a regulatory activity of chromosome segregation; (4) a DNA hybridizing with the DNA according to “2” under stringent conditions and encoding a protein that has a regulatory activity of chromosome segregation; (5) a protein consisting of an amino acid sequence shown in SEQ ID NO: 2; and (6) a protein consisting of an amino acid sequence where one or several amino acids are deleted, replaced or added in an amino acid sequence shown in SEQ ID NO: 2, and having a regulatory activity of chromosome segregation.

[0007] The present invention also relates to (7) a DNA encoding a following protein (a) or (b): (a) a protein consisting of an amino acid sequence shown in SEQ ID NO: 4, (b) a protein consisting of an amino acid sequence where one or several amino acids are deleted, replaced or added in an amino acid sequence shown in SEQ ID NO: 4, and having a regulatory activity of chromosome segregation; (8) a DNA consisting of a base sequence shown in SEQ ID NO: 3 or a complementary sequence thereof; (9) a DNA containing part or whole of a base sequence shown in SEQ ID NO: 3 or a complementary sequence thereof, and encoding a protein that has a regulatory activity of chromosome segregation; (10) a DNA hybridizing with the DNA according to “8” under stringent conditions and encoding a protein that has a regulatory activity of chromosome segregation; (11) a protein consisting of an amino acid sequence shown in SEQ ID NO: 4; and

[0008] (12) a protein consisting of an amino acid sequence where one or several amino acids are deleted, replaced or added in an amino acid sequence shown in SEQ ID NO: 4, and having a regulatory activity of chromosome segregation.

[0009] The present invention further relates to (13) a DNA encoding a following protein (a) or (b): (a) a protein consisting of an amino acid sequence shown in SEQ ID NO: 6, (b) a protein consisting of an amino acid sequence where one or several amino acids are deleted, replaced or added in an amino acid sequence shown in SEQ ID NO: 6, and having

a regulatory activity of chromosome segregation; (14) a DNA consisting of a base sequence shown in SEQ ID NO: 5 or a complementary sequence thereof; (15) a DNA containing part or whole of a base sequence shown in SEQ ID NO: 5 or a complementary sequence thereof, and encoding a protein that has a regulatory activity of chromosome segregation; (16) a DNA hybridizing with the DNA according to “14” under stringent conditions and encoding a protein that has a regulatory activity of chromosome segregation; (17) a protein consisting of an amino acid sequence shown in SEQ ID NO: 6; and (18) a protein consisting of an amino acid sequence where one or several amino acids are deleted, replaced or added in an amino acid sequence shown in SEQ ID NO: 6, and having a regulatory activity of chromosome segregation.

[0010] The present invention still further relates to (19) a DNA encoding a following protein (a) or (b) that has a regulatory activity of chromosome segregation: (a) a protein consisting of an amino acid sequence shown in SEQ ID NO: 8, 10, 12, 14, 16, 18 or 20, (b) a protein consisting of an amino acid sequence where one or several amino acids are deleted, replaced or added in an amino acid sequence shown in SEQ ID NO: 8, 10, 12, 14, 16, 18 or 20; (20) a DNA consisting of a base sequence shown in SEQ ID NO: 7, 9, 11, 13, 15, 17 or 19 or a complementary sequence thereof, and encoding a protein that has a regulatory activity of chromosome segregation; (21) a DNA containing part or whole of a base sequence shown in SEQ ID NO: 7, 9, 11, 13, 15, 17 or 19 or a complementary sequence thereof, and encoding a protein that has a regulatory activity of chromosome segregation; (22) a DNA hybridizing with the DNA according to “7”, “9”, “11”, “13”, “15”, “17” or “19” under stringent conditions and encoding a protein that has a regulatory activity of chromosome segregation; (23) a protein consisting of an amino acid sequence shown in SEQ ID NO: 8, 10, 12, 14, 16, 18 or 20, and having a regulatory activity of chromosome segregation; and (24) a protein consisting of an amino acid sequence where one or several amino acids are deleted, replaced or added in an amino acid sequence shown in SEQ ID NO: 8, 10, 12, 14, 16, 18 or 20, and having a regulatory activity of chromosome segregation.

[0011] Furthermore, the present invention relates to (25) a fusion protein in which the protein according to “5”, “6”, “11”, “12”, “23” or “24” is bound with a marker protein and/or a peptide tag; (26) an antibody specifically binding to the protein according to “5”, “6”, “11”, “12”, “23” or “24”; and (27) the antibody according to “26”, which is a monoclonal antibody.

#### BRIEF DESCRIPTION OF DRAWINGS

[0012] FIG. 1 is a set of pictures showing that sister chromatids are not segregated during mitosis by co-expression of Sgo1 and Rec8 in the present invention. a.) The cen2-GFP strains expressing the genes indicated by endogenous promoters (a constitutive chromatin promoter for rad21+ or rec8+, and a thiamine-repressible promoter Pnmt1 for Sgo1+) were streaked on a thiamine-depleted plate. b.) Samples of Padh1-rec8+Pnmt1-sgo1+ cells cultured for 15 hours at 30° C. after thiamine depletion. The non-segregation of cen2-GFP (asterisk) was identified in the septate junction cells. c.) The non-segregations of cen2-GFP were counted (n>100). d.) The Padh1-rec8+-GFP strains were



cultured with or without the use of Pnmt1-sgo1+ in the same manner as (b). Samples of cells at interphase and anaphase are shown.

[0013] FIG. 2 is a set of pictures showing that sister chromatid segregation was undergone in mitosis by expression of non-cleavable Rec8. The plasmid pREP41-rec8-RDRD (expressing non-cleavable Rec8 (Embo J 22, 5643-53(2003))) was integrated into the chromosome of cen2-GFP cell strains (+Rec8-RDRD), and the cells were streaked on plates with or without the presence of thiamine. The host strain cells (-Rec8-RDRD) were similarly cultured as a control. Note that Rec8-RDRD is expressed only on the thiamine-free plate. Samples of cells cultured in culture medium for 15 hours at 30° C. after the depletion of thiamine.

[0014] FIG. 3 is a set of pictures showing that sgo1 of the present invention is required to protect Rec8 and thereby cohesion at centromeres arises during anaphase of meiosis I. a.) As for one of the homologues marked with cen2-GFP, segregation during meiosis was observed in wild-type and sgo1Δ cells (n>170). A normal segregation pattern of cen2-GFP is illustrated (left). Samples of sgo1Δ cells are shown (right). b.) Separation of sister cen2-GFP dots after meiosis I (mes1Δ arrest) is evident in sgo1Δ cells. c.) The Rec8-GFP signal was observed in the indicated cells at late anaphase I (n>30) and at prometaphase II (n>100), and the frequency of centromeric Rec8-GFP displayed in the cells was counted. The spindles were visualized by expressing CFP-Atb2 (α-tubulin) (Curr Biol 11, 836-45(2001)). d.) Rec8-GFP levels throughout the indicated chromosome sites in the arrested cells were measured prior to meiosis I (mei4Δ arrest) by ChIP assay with the use of anti-GFP antibodies. The bottom panel shows *Schizosaccharomyces pombe* chromosome I schematically, and the primers (cnt, imr, dg, dh, lys1, mes1) were used there.

[0015] FIG. 4 is a set of pictures showing that Sgo1 of the present invention localizes at pericentromeric regions during meiosis I. a.) Synchronous meiosis of diploid pat1-114/pat1-114 cell strains (Embo J 22, 5643-53(2003)) was sampled, meiotic nuclear division was monitored by DAPI staining, and the protein level of Sgo1 was detected by Western blotting with the use of anti-Sgo1 antibodies. b.) Sgo1 (green) was counterstained with tubulin (red) and DAPI (4'6'-diamidino-2-phenylindole) (blue) at the indicated stages in meiotic cells. c.) A sgo1+-GFP cell co-expressing mis6+-CFP was examined under fluorescence microscopy. Sgo1-GFP (green) and Mis6-CFP (red) are merged. d.) Sgo1-GFP levels throughout the indicated chromosome sites in cells arrested at metaphase I were measured by ChIP assay with the use of anti-GFP antibodies. The same primers as for FIG. 2d in synchronism with additional primers at mat (heterochromatin region at the mating type locus) and TAS (telomere associated sequence) were used. e.) Sgo1-GFP (green) was detected at metaphase I in the indicated cells that express CFP Atb2 to visualize spindles (red). f.) Rec8-HA was expressed with or without Sgo1-FLAG in proliferating cells, and the extracts were immunoprecipitated with anti-FLAG antibody. g.) A model for the action of shugoshin in meiosis. Shugoshin protects centromeric Rec8 complexes from cleaving by separase at the onset of anaphase I, thereby preserves the centromeric cohesion until meiosis II. Shugoshin is degraded depending on APC during anaphase I.

[0016] FIG. 5 is a set of pictures showing the time-dependent change of the expression levels of Sgo1 and Rec8 in synchronous culture of haploid pat1-114 cell strains (wt), and of cut1-206 or Prad21-slp1 cells. The expression of slp1+ (a fission yeast CDC20 homologue required for APC activation (Mol Cell Biol 17, 742-50(1997))) was repressed during meiosis in Prad21-slp1 cells where slp1 promoter was replaced with rad21. Meiotic nuclear division was monitored by DAPI staining, and the protein levels of Sgo1, Rec8, and tubulin (control) were measured by western blotting with the use of anti-Sgo1, anti-Rec8 and anti-tubulin antibodies, respectively. Although cut1-206 cells together with normal kinetics led to Sgo1 degradation, Rec8 degradation was delayed. Prad21-slp1 cells showed delayed degradation of Sgo1 as well as Rec8. Arrowheads indicate a cleavage product of Rec8 by separase Cut1.

[0017] FIG. 6 is a set of pictures showing that ectopic expression of sgo1+ inhibits the growth of the cut1-206 mutant. Chromosomal sgo1+ promoter was replaced with Pnmt1 or Pnmt41 (a weaker version of Pnmt1), and the effect on the mitotic growth in cut1-206 temperature-sensitive cells was examined. The indicated cells were streaked on a plate without thiamine and cultured for 3 days at 28° C. The cut1-206 cells moderately expressing Sgo1 by Pnmt1, arrested mitotic growth even at the permissive temperature, whereas cut1+ cells grew normally.

[0018] FIG. 7 is a set of pictures showing that Sgo2 of the present invention plays an important role in mitotic at centromere. a.) Serial dilutions of the indicated cultures were spotted onto YEA plates containing 0, 5 or 10 μg/ml of TBZ, and cultured for 3 days at 30° C. b.) The indicated strains were streaked on YEA plates and cultured for 3 days at 30° C. c.) Sgo2-GFP (green) was detected at anaphase I in wild-types and in bub1Δ cells that express CFP-Atb2 to visualize spindles (red). DNA was stained with Hoechst (blue). Wild-type cells at anaphase are also shown. d.) The sgo2+-GFP mis6+-HA cells were fixed and stained with anti-GFP and anti-HA antibodies. e.) Sgo2-GFP levels were measured throughout the indicated chromosome sites in cells arrested at prometaphase or in asynchronous cells by ChIP assay.

[0019] FIG. 8 is a set of pictures showing the results of analysis of budding yeast shugoshin ScSgo1 of the present invention. a.) Budding yeast ScSGO1-GFP diploids in proliferation were fixed with methanol and counterstained with DAPI. b.) ScSGO1-Myc NDC10-HA cells were fixed, and stained with DAPI and antibodies against Myc and HA. c.) ScSGO1-GFP diploids causing meiosis in culture medium were fixed with methanol and counterstained with DAPI. d.) Serial dilutions of the indicated cultures were spotted onto YPD plates containing 0 or 15 μg/ml of benomyl. e.) Chromosome loss was analyzed in wild-types (wt) and Scsgo1Δ mutants by a colony sectoring assay. The loss of nonessential chromosome fragments resulted in a red sector in a white colony. As a positive control, ubr1Δ mutant was used (Nature 410, 955-9(2001)). The frequency of sectoring colonies is shown at the bottom (n>120). f.) Samples of segregation of cenV-GFP in Scsgo1Δ tetrads. The segregation patterns in tetrads were mostly classified as one of the three shown at the bottom. The each population (n=200) is also shown. g.) ScSGO1-Myc diploids were induced by synchronous meiosis and were examined the segregation of cenV-GFP marked on one of two homologues at meiosis I

and meiosis II. Although most of the cells caused reductional segregation pattern at meiosis I (96%, n=207), the incidence of non-segregation was high at meiosis II (34%, n=322). The cells marked with cenV-GFP on both homologues were induced to meiosis, and counterstained with anti-tubulin antibody and DAPI. Cells at late anaphase I were examined for cenV-GFP dots. ScSgo1-Myc cells frequently showed split cenV-GFP dots at either pair of sister chromatids (72%, n=138), while control wild-type cells did not (<2%, n=106).

[0020] FIG. 9 is a set of pictures showing sequences of the amino terminal coiled-coil regions and carboxyl terminal basic regions of shugoshin-like proteins in various organisms. The primary sequences of the amino terminal regions of Sgo1 are conserved in *Schizosaccharomyces pombe* (Sgo1 and Sgo2), budding yeast (ScSgo1) and *Neurospora crassa* (B23G1.060), while the sequences containing ME1-S332 in other species are not conserved, all presumably carry coiled-coil motif (predicted by COILS program (Science 252, 1162-4(1991))). See the arrowheads, asterisks and circles in the pictures.

[0021] FIG. 10 is a picture showing the results of examination of sgo1 mutations that were generated within conserved regions. Both h+sgo1Δ and h-sgo1Δcen2-GFP cells transformed with the indicated plasmid, were mixed on SPA plates and monitored for segregation of cen2-GFP at meiosis II. A plasmid pREP81 bearing a weak version of the thiamine-repressible nmt1 promoter was used to express sgo1. Control cells carrying plasmid pREP81-sgo1 (wt) showed nearly 80% the segregation at meiosis II, whereas cells expressing non-segregation sgo1 allele showed random segregation (50% segregation). Any of the mutations tested, except a non-conserved site mutation 297TA, did not complement sgo1Δ in this assay. The means of two independent experiments are shown (n>100).

[0022] FIG. 11 (a) is a picture showing schematic representation of the shugoshin family proteins. A predicted coiled-coil (red) and a conserved basic region (blue) exist in the N-terminal and C-terminal regions respectively. Further, FIG. 11 (b) is a picture showing the result of analysis in HeLa cell extracts by western blotting after transfection with siRNA.

[0023] FIG. 12 is a set of pictures showing the results that HeLa cells were stained (green) with antibody against hSgo1 or hSgo2 prepared from rabbit, concurrently stained with tubulin antibody and DAPI, and then respectively co-stained with spindle (red) and chromosome DNA (blue). Meanwhile, the cells were fixed with paraformaldehyde.

[0024] FIG. 13 is a set of pictures showing the results that HeLa cells at prometaphase and metaphase were stained with antibodies against hsgo1 or hSgo2 (green), and concurrently co-stained with antibodies against centromere protein Aurora B of chromosome localized within kinetochore from prophase to metaphase (b, d; red), and DAPI (blue). Both signals of hSgo1 and hSgo2 showed signals at the sites close to CENP-A dots on chromosome. From the above, it was revealed that both hsgo1 and hSgo2 are centromere proteins. Furthermore, both sites of Sgo1 and Aurora B were practically the same at prometaphase and metaphase, whereas Sgo2 was placed just outside Aurora B. From the above, it was revealed that both hsgo1 and hSgo2 are placed within kinetochore from prometaphase to metaphase.

[0025] FIG. 14 is a picture showing the results of RNAi experiments that targeted hsgo1 and hSgo2 respectively. The expressions in any proteins were significantly suppressed after 48 hours, thereby the cells arrested in mitosis (total in the figure) were accumulated. As the accumulation was dissolved by suppressing a spindle checkpoint factor BubR1 by RNAi, it was suggested that hSgo1 and hSgo2 directly or indirectly function during the process where spindle take kinetochore properly at centromeres.

[0026] FIG. 15 is a set of pictures showing the results, where RNAi experiments targeting hsgo1 was performed by using HeLa cells, and then the cells were mounted on a slide glass and stained with Giemsa. It was revealed that sister chromatid strongly adhered at centromere site in control cells; but in cells suppressed hsgo1, the adhesion at centromere site was weak, and easily detached by the experiment operation.

[0027] FIG. 16 is a set of pictures showing that Sgo1 and Bub1 are required for condensation at centromeres in mitosis. (a) By treatments with siRNA, chromosome spread was performed in mitotic HeLa cells stained with Giemsa. Representative spread is shown together with the occurrence rates. More than one hundred of the prophases and prometaphases were observed for each RNAi. An example of sister chromatid pair is magnified at the top. (b) After treatment with nocodazole for 4 hours, chromosome spread was observed in cells interfered with RNAi. Examples of the spread are shown with the frequency (n>100). (c) HeLa cells expressing Scc1-myc were fixed at 36 hours after the treatment with siRNAs. The cells were immunostained with anti-myc-antibody (green) and anti-centromere-antibody (ACA) (red). DNA was stained with DAPI (blue). (d) Rates of the cells showing Scc1-myc staining are shown. Cells expressing Scc1-myc in this cell line were less than 25%. Scale bar shows 10 μm.

[0028] FIG. 17 is a set of pictures showing the results of RNAi experiments targeting Bub1, respectively. (A, B) RNAi experiments targeting Bub1 were performed respectively, and resulted in disappearance of the localization of both proteins, hSgo1 and hSgo2 at centromere. (C, D) As the localization of both proteins, hSgo1 and hSgo2 at centromere was normal in RNAi experiments targeting a control, BubR1; the significance of the results of Bub1 was ensured. It is shown that Bub1 and BubR1 are similar but different proteins, and the localization of hSgo1 and hSgo2 at centromere depends on Bub1 (A, B), but not on BubR1 (C, D).

[0029] FIG. 18 is a set of pictures showing the results that a clone in which cDNA of mouse shugoshin homologous gene (SEQ ID NOS: 21 and 23) is fused with GFP gene was generated by using retroviral vector, and expressed in human HeLa cells. It was revealed that any of the GFP fusion proteins is co-localized with human kinetochore protein Bub1 in mitosis.

#### BEST MODE OF CARRYING OUT THE INVENTION

[0030] As for a protein of the present invention, a protein Sgo1 (shugoshin) comprising an amino acid sequence shown in SEQ ID NO: 2 and having a regulatory activity of chromosome segregation; a protein comprising the amino acid sequence shown in SEQ ID NO: 2 where one or several

amino acids are deleted, replaced or added, and having a regulatory activity of chromosome segregation; a paralogue Sgo2 of protein Sgo1 comprising an amino acid sequence shown in SEQ ID NO: 4 and having a regulatory activity of chromosome segregation; a protein comprising the amino acid sequence shown in SEQ ID NO: 4 where one or several amino acids are deleted, replaced or added, and having a regulatory activity of chromosome segregation; a *Saccharomyces cerevisiae* homologue ScSgo1 of protein Sgo1 comprising an amino acid sequence shown in SEQ ID NO: 6 and having a regulatory activity of chromosome segregation; a protein comprising the amino acid sequence shown in SEQ ID NO: 6 where one or several amino acids are deleted, replaced or added, and having a regulatory activity of chromosome segregation; a protein (NC) comprising an amino acid sequence shown in SEQ ID NO: 8 and having a *Neurospora crassa*-derived regulatory activity of chromosome segregation; a protein comprising the amino acid sequence shown in SEQ ID NO: 8 where one or several amino acids are deleted, replaced or added, and having a regulatory activity of chromosome segregation; a protein (At) comprising an amino acid sequence shown in SEQ ID NO: 10 or 12 and having a *Arabidopsis*-derived regulatory activity of chromosome segregation; a protein comprising the amino acid sequence shown in SEQ ID NO: 10 or 12 where one or several amino acids are deleted, replaced or added, and having a regulatory activity of chromosome segregation; a protein (Mm) comprising an amino acid sequence shown in SEQ ID NO: 14 or 16 and having a mouse-derived regulatory activity of chromosome segregation; a protein comprising the amino acid sequence shown in SEQ ID NO: 14 or 16 where one or several amino acids are deleted, replaced or added, and having a regulatory activity of chromosome segregation; a protein (Hs) comprising an amino acid sequence shown in SEQ ID NO: 18 or 20 and having a human-derived regulatory activity of chromosome segregation; and a protein comprising the amino acid sequence shown in SEQ ID NO: 18 or 20 where one or several amino acids are deleted, replaced or added, and having a regulatory activity of chromosome segregation; can be exemplified. Further, as for the regulatory activity of chromosome segregation described in the above, although it is not especially limited as long as the activities regulate chromosome segregation, for example, activities correctly regulating chromosome segregation of germ cells and/or of somatic cell division are preferable, and activities protecting (Shugo) the centromere of sister chromatid from the separation in meiosis 1 is more preferable. In addition, proteins of the present invention can be prepared by known methods based on DNA-sequence information and the like, and the derivations are not limited to yeast, mouse, human and the like. Furthermore, for example, Sgo1 (shugoshin) mutant that is a protein comprising an amino acid sequence shown in SEQ ID NO: 2 where one or several amino acids are deleted, replaced or added, and having a regulatory activity of chromosome segregation, can be prepared by ordinary methods such as known gene manipulation, point mutation and the like.

[0031] As for a DNA of the present invention, a DNA encoding a protein of the present invention that has a regulatory activity of chromosome segregation: a DNA derived from fission yeast *Schizosaccharomyces pombe*, comprising a base sequence shown in SEQ ID NO: 1 or 3 or a complementary sequence thereof; and a DNA containing

part or whole of these sequences, encoding a protein that has a regulatory activity of chromosome segregation: a DNA derived from *Saccharomyces cerevisiae*, comprising a base sequence shown in SEQ ID NO: 5 or a complementary sequence thereof; and a DNA containing part or whole of these sequences, encoding a protein that has a regulatory activity of chromosome segregation: a DNA derived from *Neurospora crassa*, comprising a base sequence shown in SEQ ID NO: 7 or a complementary sequence thereof, and encoding a protein that has a regulatory activity of chromosome segregation; and a DNA containing part or whole of these sequences, encoding a protein that has a regulatory activity of chromosome segregation: a DNA derived from *Arabidopsis*, comprising a base sequence shown in SEQ ID NO: 9 or 11 or a complementary sequence thereof, and encoding a protein that has a regulatory activity of chromosome segregation; and a DNA containing part or whole of these sequences, encoding a protein that has a regulatory activity of chromosome segregation: a DNA derived from mouse, comprising a base sequence shown in SEQ ID NO: 13 or 15 or a complementary sequence thereof, and encoding a protein that has a regulatory activity of chromosome segregation; and a DNA containing part or whole of these sequences, encoding a protein that has a regulatory activity of chromosome segregation: a DNA derived from human, comprising a base sequence shown in SEQ ID NO: 17 or 19 or a complementary sequence thereof, and encoding a protein that has a regulatory activity of chromosome segregation; and a DNA containing part or whole of these sequences, encoding a protein that has a regulatory activity of chromosome segregation: a DNA hybridizing with the above DNA under stringent conditions, encoding a protein that has a regulatory activity of chromosome segregation: and the like, can be exemplified.

[0032] These DNAs can be prepared by known methods based on DNA-sequence information, such as a gene or cDNA library of yeast, mouse, human and the like. Further, using a base sequence shown in SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, or others or a complementary sequence thereof, or part or whole of these sequences as a probe, DNA libraries of yeast, mouse, human and the like are hybridized under stringent conditions, and the intended DNA encoding a protein that has a regulatory activity of chromosome segregation can be obtained by isolating the DNAs that hybridized with the probes. As for a condition of hybridization to obtain the DNA; hybridization at 42° C., and washing treatment by a buffer containing 1×SSC and 0.1% SDS at 42° C.; preferably hybridization at 65° C., and washing treatment by a buffer containing 0.1×SSC and 0.1% SDS at 65° C.; can be exemplified. Moreover, as for an element affecting the stringency of hybridization, there are various elements other than the above described temperature conditions, those skilled in the art can actualize the stringency equivalent to that of hybridization as exemplified in the above with an appropriate combination of various elements.

[0033] As for a fusion protein of the present invention, any protein can be used as long as the protein of the present invention is bound to a marker protein and/or a peptide tag, as for a marker protein, it is not especially limited but a conventionally known marker protein, for example, alkaline phosphatase, Fc region of antibody, HRP, GFP and the like can be exemplified. Further, as for a peptide tag of the present invention, conventionally known peptide tags such

as Myc, His, FLAG and GST tags can be specifically exemplified. The fusion protein can be produced by ordinary methods; and is useful for purification of protein Sgo1 and the like by using the affinity of Ni-NTA and His tag, and for a reagent for study in the art.

**[0034]** As for an antibody specifically binding to a protein of the present invention, immunospecific antibodies such as monoclonal antibody, polyclonal antibody, chimeric antibody, single-stranded antibody, humanized antibody and the like, can be specifically exemplified. These antibodies can be produced by ordinary methods with the use of proteins such as the above-mentioned Sgo1 or part thereof as an antigen, and among them a monoclonal antibody is preferable in terms of specificity. Antibodies such as a monoclonal antibody are useful for elucidating the localization of Sgo1 and others in vivo.

**[0035]** The above-mentioned antibodies of the present invention can be generated with the use of common protocol by administering proteins of the present invention or fragments containing epitope thereof, or cells expressing the protein on their membrane surfaces, to animals (preferably non-human). For example, for preparation of a monoclonal antibody any method such as hybridoma (Nature 256, 495-497, 1975), trioma, human B cell hybridoma (Immunology Today 4, 72, 1983) and EBV-hybridoma (MONOCLONAL ANTIBODIES AND CANCER THERAPY, pp. 77-96, Alan R. Liss, Inc., 1985), by which antibodies are generated from cultures of continuous cell lines, can be used.

**[0036]** To generate a single-stranded antibody against a protein of the present invention, a method for preparation of single-stranded antibody (U.S. Pat. No. 4,946,778) can be applied. Further, to express a humanized antibody, transgenic mouse or other mammals can be used, clones that express a protein of the present invention with the use of the above-mentioned antibody can be isolated/identified, and its polypeptide can be purified by affinity chromatography. Antibodies against peptide containing proteins of the present invention or antigen epitopes thereof can be possibly used for diagnosis and treatment of cancer, or of chromosome segregation diseases such as infertility or Down's syndrome using a regulatory factor of chromosome segregation as an index.

**[0037]** Functional analysis of a protein of the present invention can be performed by using fusion proteins fused with, for example; fluorescent substances such as FITC (fluorescein isocyanate) or tetramethyl rhodamine isocyanate; radioisotopes such as <sup>125</sup>I, <sup>32</sup>P, <sup>14</sup>C, <sup>35</sup>S or <sup>3</sup>H; labelings with enzymes such as alkaline phosphatase, peroxidase,  $\beta$ -galactosidase or phycoerythrin; fluorescence emission proteins such as green fluorescent protein (GFP); or the like, to antibodies such as the above-mentioned monoclonal antibodies. As an immunological assay method with the use of antibody of the present invention, methods such as RIA, ELISA, Fluorescent antibody method, Plaque forming cell assay, Spotting method, Hemagglutination testing, Ouchterlony method can be exemplified.

**[0038]** The present invention will be explained in detail in the following by referring to the examples, but the technical scope of the present invention will not be limited to these.

## EXAMPLE 1

### [Method]

#### (Screening of Rec8 Protector)

**[0039]** The present inventor examined a gene that is toxic only when co-expressed with Rec8 in vegetative cells. The Rec8 encoding sequence that was fused with GFP was cloned into pREP82 (ura4+ marker) under the thiamine-repressible nmt1+ promoter, to construct pREP82-rec8+-GFP. A *Schizosaccharomyces pombe* cDNA library constructed by mRNA that was prepared from meiotic cells, and a pREP3 vector (nmt1+ promoter, LEU2+ marker) (Y. Akiyoshi and Y. W., unpublished) were used. The leu1 ura4-D18 cells carrying pREP82-rec8+-GFP were transformed with the cDNA library, spread on agar plates containing thiamine (promoter-off) and incubated for 3 days at 30° C. The colonies were then replicated on two thiamine-free agar plates: one that contains uracil and 5'-fluoroorthoic acid (5'-FOA) where only cells lacked the plasmid pREP82-rec8+-CFP can grow (thereby expresses a library clone alone), and the other that does not contain 5'-FOA (allows co-expression of rec8+-GFP and a library clone). The present inventor added Phloxine B, a drug that stains dead cells red, onto the both agar plates, thereby illuminated sick colonies. After incubation for two days, the colonies exhibiting sickness only on the co-expression agar plate were picked up, and the library-derived plasmids were recovered and analyzed.

#### (*Schizosaccharomyces pombe* Strains)

**[0040]** Deletion and tagging of GFP or FLAG to endogenous sgo1+ and sgo2+ were performed by a PCR-based gene targeting method (Yeast 14, 943-951(1998)). By inserting GFP into the C-terminus of the PCR-amplified sgo1+-FLAG, sgo1+-FLAG-GFP was generated and integrated into the endogenous sgo1 locus. Further, an endogenous promoter of the sgo1+ was replaced with a nmt promoter to generate Pnmt-sgo1+ or Pnmt-sgo1+-FLAG-GFP by the PCR-based gene targeting method. The proteins tagged to Sgo1-GFP or Sgo1-FLAG was deleted depending on the purpose. A mei4 $\Delta$  mutation was used to arrest meiotic cells prior to meiosis I (close to late prophase in meiosis I), and a mes1 $\Delta$  mutation was used to arrest after meiosis I, as described previously (Nature 400, 461-4(1999)).

#### (Observation of Chromosomes Marked with GFP)

**[0041]** To observe the segregation patterns of homologues at meiosis I, h90 cells retaining cen2-GFP (Embo J 22, 2284-96(2003)) were spotted on meiosis-inducing medium, SPA. To examine the segregation patterns of sister chromatids, opposite mating type cells, one marked with cen2-GFP and the other not marked, were mixed and spotted on SPA. After incubation for one day, the zygotes were monitored for GFP. Images were obtained under a microscope (Axioplan2, Zeiss) equipped with a cooled CCD camera (Quantix, Photometrics) and by using Metamorph software (Universal Imaging Corporation). Seven Z-sections for GFP signals were converted to single two-dimensional images by taking the maximum signal at each pixel position in the images.

#### (Chromatin Immunoprecipitation; ChIP)

**[0042]** Diploid sgo1+-FLAG-GFP was used for ChIP with Sgo1. To achieve a highly synchronous culture, the endogenous slp1+ promoter was replaced with the rad21+ pro-

moter that is not active during meiosis, and the cells were arrested at metaphase I. The cells were incubated in nitrogen-depleted medium for 17 hours at 30° C., and 60% the cells or less were arrested at metaphase I. For ChIP with Sgo2, *nda3-KM311 sgo2+*-GFP cells were proliferated at 30° C., and then shifted to 18° C. After incubation for 8 hours, most of the cells were arrested at metaphase. The cells were fixed with 3% para-formaldehyde for 30 minutes at 18° C., and extracts were prepared. The DNA was broken to an average size of 400 bp, and the extracts were immunoprecipitated with rabbit anti-GFP antibodies (Clontech). DNAs prepared from the whole cell crude extracts, or immunoprecipitated chromatin fractions were analyzed by quantitative PCR, with a LightCycler or a Lightcycler-DNA Master SYBR Green 1 kit (Roche Molecular Biochemicals). Antibody-minus samples were used as controls in each experiment to explain the nonspecific binding in the ChIP fractions.

#### (Preparation of Anti-Sgo1 Antibodies)

**[0043]** Sgo1+ ORF was PCR-amplified from an *Schizosaccharomyces pombe* cDNA library, and inserted into plasmids pGEX4T-2 (Pharmacia Biotech) and pET-19b (Novagen) respectively to prepare recombinant proteins GST-Sgo1 and His-Sgo1. GST-Sgo1 was used to immunize rabbit, and the raised antibodies were purified by His-Sgo1 as described previously (Embo J 22, 5643-53(2003)). Furthermore, for the purpose of analyzing proteins (SEQ ID NOs: 18 and 20; hSgo1 and hSgo2 respectively) encoding human shugoshin homologous gene (SEQ ID NOs: 17 and 19), part of hSgo1 and hSgo2 was expressed in *E. coli*, and antibodies against hSgo1 and hSgo2 were produced by injecting the protein into rabbit.

#### (Immunostaining)

**[0044]** To stain endogenous Sgo1, wild-type diploid cells cultured for 5 hours in MM-N were fixed with 3% formaldehyde for 40 min at 30° C., and stained by the method described previously (Embo J 22, 5643-53(2003)). To stain Sgo2-GFP and Mis6-HA, logarithmically growing cells were used. Sgo1 was detected by using rabbit anti-Sgo1 antibody at 1:50 and Alexa488-conjugated anti-rabbit antibody (Molecular Probes) at 1:100. Tubulin was detected by using mouse anti-tubulin antibody TAT-1 (provided by Keith Gull) at 1:200 and Cy3-tagged anti-mouse antibody (Chemicon) at 1:2000. Cells were counterstained with DAPI to visualize DNA. The Sgo2-GFP was detected by using mouse anti-GFP antibody (Roche) at 1:50 and BODIPY FL-conjugated anti-mouse antibody (Molecular Probes) at 1:100. The Mis6-HA was detected by using rabbit anti-HA antibody Y-11 (Santa Cruz) at 1:50 and Alexa488-conjugated anti-rabbit antibody at 1:100. Cells were counterstained with DAPI to visualize DNA. Further, immunostaining was performed by using rabbit anti-hSgo1 antibody and rabbit anti-hSgo2 antibody in the same manner as the above.

#### (Coimmunoprecipitation)

**[0045]** Padh-rec8+3HA Pnmt41-sgo1+-FLAG-GFP strain cells and control Padh-rec8+3HA strain cells were cultured without thiamine for 15 hours at 30° C., collected, and the extracts were prepared. To liberate chromatin-bound proteins, the extracts were treated with DNase I. After clarifying the extracts by centrifugation, the Sgo1-FLAG-GFP protein was immunoprecipitated with anti-FLAG anti-

body M2 (Sigma). The Rec8-3HA and Sgo1-FLAG-GFP were detected by anti-HA antibody Y-11 and anti-FLAG antibody M2, respectively.

#### (Analysis of Budding Yeast)

**[0046]** All sample strains except those for chromosome loss assay are derivative of SK1 (Cell 98, 91-103(1999)). The chromosome loss assay was performed as described previously (Nature 410, 955-9(2001)). The ScSGO1 gene was deleted or epitope-tagged by using PCR generated cassettes (Yeast 14, 953-961(1998)). Accurate gene targeting was checked by PCR. URA3-GFP dots marking chromosome V (cenV-GFP) were described previously (Cell 98, 91-103(1999)). Sporulation was induced by culturing diploid cells at 30° C. as described previously (Dev Cell 4, 535-48(2003)). In situ immunofluorescence was performed as described previously (Dev Cell 4, 535-48(2003)).

#### (Cell Culture)

**[0047]** HeLa cells were cultured in DMEM supplemented with 10% fetal bovine serum and 0.03% L-Glutamine. The HeLa cell strain expressing Scc1-myc was cultured with 200 µg/ml of G418 (Invitrogen) and 100 µg/ml of Hygromycin B (Wako). Expression of Scc1-myc was induced by incubation with 2 µg/ml of Doxycyclin (Sigma) for 48 hours.

#### (Preparation of Anti-Human Sgo Antibody)

**[0048]** As the information for N-terminal amino acid sequence of human Sgo1 was not obtained from the databases, the present inventor cloned a cDNA fragment that was amplified from a cDNA library (BD Biosciences) with the use of primers recognizing the cloning site of λTriplEx: CTCGGGAAGCGCGCCATTGTG (SEQ ID NO: 38) and the DNA sequence corresponding to the numbers 237-242 in amino acid sequence of Q9BVA8: CCTGGCTGAAT-CAGCTTTGGTG (SEQ ID NO: 39). The Sequencing revealed that the Sgo1 mRNA encodes a protein having 527 amino acids. To obtain polyclonal antibodies against Sgo1, a cDNA fragment encoding the numbers 109-491 in amino acid sequence of Sgo1 was amplified and inserted into the reading frames of plasmids pGEX4T-2 (Amersham) and pET19b (Novagen) to produce GST-Sgo1 and His-Sgo1 respectively, and followed by immunization of a rabbit (QIAGEN) (performed according to the manufacturer's instructions). His-Sgo1 was affinity-purified on CNBr-activated sepharose (Amersham). Antibodies against Sgo2 were raised with GST-Sgo2 (amino acid numbers 331-631) and purified with His-Sgo2 in the same manner as the above.

#### (Immunofluorescence Microscopy and Chromosome Spreading)

**[0049]** Immunofluorescent staining was performed as described in the above, by using anti-human Sgo1 (1:1000), anti-human Sgo2 antiserum (1:10000), anti-Bub1 (1:1000, MBL), anti-BubR1 (1:1000, MBL), anti-CENP-A (1:1000, MBL), anti-Aurora B AIM-1 (1:1000, BD Biosciences) and anti-tubulin DM1A (1:1000, Sigma). Immunostaining of Scc1-myc was performed as described in the above, by using anti-myc CM-100 (1:1000, Gramsch Laboratories) and ACA (1:1000, provided from Dr. Yoshinari Takasaki). As a secondary antibody, Alexa Fluor 488 goat anti-rabbit antibody (1:1000, Molecular Probes), Cy3 conjugated anti-mouse antibody (1:1000, CHEMICON), and Cy3 conjugated donkey anti-human antibody (1:1000, Jackson ImmunoRe-

search Laboratories, Inc) were used. 3 µg/ml of Hoechst 33342 or 0.5 µg/ml of DAPI were used for counter staining. Images were taken by using SlideBook or MetaMorph software.

(Chromosome Spreading)

[0050] HeLa cells during mitosis were collected by mitotic shake-off and incubated with 330 nM of nocodazole for 0 up to 4 hours. Chromosome spreading was performed as described in the above.

(Immunoblotting)

[0051] HeLa cells were boiled with the sample buffer and resolved by SDS-polyacrylamide gel electrophoresis. Proteins were transferred to Immobilon-P membrane (Millipore), followed by blocking with 5% Skim milk (Nacalai) in TBST (150 mM of NaCl, 20 mM of Tris-HCl pH7.4, 0.05% Tween-20). Antibody incubations were performed in 0.1% skim milk TBST supplemented with anti-Sgo1 antibody (1:1000), anti-Sgo2 antibody (1:1000), anti-Bub1 antibody (1:500) and anti-tubulin antibody (1:3000). Blots were produced by ECL (Amersham).

(RNAi)

[0052] As a siRNA target sequence, hSgo1:

```

hSgo1:                               (SEQ ID NO:40)
AAGUCUACUGAUAAUGUCUUATT
and
hSgo2:                               (SEQ ID NO:41)
AAGCACUACCACUUUGAAUAATT,
and
human Sgo1:                          (SEQ ID NO:42)
GUGAGCCUCUGUGAAUCAATT
and
human Sgo2:                          (SEQ ID NO:43)
GCUCUCAUGAACAAUACUTT
    
```

GCUCUCAUGAACAAUACUTT (SEQ ID NO: 43) were respectively selected on hSgo1RNA or hSgo2RNA. Furthermore, as a siRNA target sequence, GAGUGAUCACGAUUUCUAATT (SEQ ID NO: 44) was selected on other siRNA target sequence, Bub1RNA; siRNA target sequence, AACGGCAUUUGAAUAUGAAA (SEQ ID NO: 45, see JCS, 117, 1577-1589(2004)) was selected at 2 sites on a spindle checkpoint factor BubR1 RNA. These sequences were synthesized as double strand, and introduced into cells by using oligofectamine (Invitrogen). Furthermore similarly, when producing HIV vector, HeLa cells were transfected with HIV plasmid vector, pMD.G (VSV-G env expressing plasmid), pMDLg/p.RRE (the third generation packaging plasmid) and pRSV Rev (Rev expressing plasmid) by calcium phosphate method, collected the culture supernatant 48 hours after the transfection, and condensed to use as a virus vector.

EXAMPLE 2

[Results]

(Identification of Shugoshin Sgo1 in Fission Yeast)

[0053] The replacement of the mitotic cohesin, Rad21/Scc1, with the meiotic version, Rec8, is a prerequisite for protecting centromeric sister chromatid cohesion through anaphase of meiosis I (Cell 103, 1155-68(2000), Mol Cell Biol 23, 3965-73(2003)). However, when Rec8 was expressed ectopically during mitosis, Rec8 was localized largely at centromeres but disappeared at anaphase, with sister chromatids segregating to opposite sides (FIGS. 1c and d). Moreover, the ectopic expression of non-cleavable Rec8 during mitosis (note that Rec8 is cleaved by separase Cut1 during meiosis (Embo J 22, 5643-53(2003))) resulted in an inability to separate sister chromatids (see FIG. 2). Thus, in contrast to the situation during meiosis I, centromeric Rec8 is cleaved by separase during mitosis, and results in separation of sister chromatids. The present inventor thus postulated a meiosis I specific centromeric protector of Rec8 from these observations. To identify this factor, the present inventor searched for a gene that generates toxicity during mitotic growth only when co-expressed with Rec8. This screening identified a novel gene, sgo1+ (ORF: SPBP35G2.03C). The hindrance of growth by Sgo1 was significantly dependent on Rec8, as Sgo1 had little effect on growth when co-expressed with Rad21 (FIG. 1a). Co-expression of rec8+ and sgo1+ resulted in high frequency of the blocked nuclear division, as centromere-associated green fluorescent protein markers (cen2-GFP) segregated to the same side of a septated cell highly frequently (see Figs. b and c). To test the possibility that Sgo1 protects Rec8 from degradation at anaphase, the localization of Rec8 was examined in associated with Sgo1 expression, Rec8 tagged with GFP at its carboxyl terminus was expressed under the control of a constitutive adh1 promoter and induced Sgo1 by using a thiamine-repressible nmt1 promoter. Consequently it was found that the Rec8-GFP signal persisted through anaphase only when Sgo1 was co-expressed (FIG. 1d). As Sgo1 is expressed exclusively in meiosis (DNA micro array data (Nat Genet 32, 143-7(2002)), see below), it was found from the above-mentioned results, that Sgo1 is a protector of Rec8 during meiosis.

(Sgo1 Protects Centromeric Cohesion at Meiosis I)

[0054] To examine whether Sgo1 is actually required for the protection of Rec8 during meiosis, the entire ORF encoding sgo1+ was deleted, and the phenotype was examined. Sgo1Δ cells are viable and showed normal vegetative growth, consistent with the concept that sgo1+ is a meiosis-specific gene. To examine the meiotic chromosome segregation of sgo1Δ cells, centromere-linked sequences were marked with GFP (cen2-GFP) on only one of the two homologues in a zygote, and the segregation of the GFP dots were monitored during meiosis I. It was revealed that meiosis I emerged normally in sgo1Δ cells, as sister chromatid pairs generally moved together to the same side of each zygote. Therefore, monopolar attachment was intact (FIG. 3a). Moreover, by marking cen2-GFP on both chromosomes, it was determined that accurate segregation was undergone with homologues at meiosis I (data not shown). However, sister chromatid pairs failed to segregate properly at meiosis II, non-segregation was caused in 50% of the cells or less (FIG. 3a). This value is consistent with random chromosome segregation at meiosis II.

[0055] To examine centromeric cohesion, cen2-GFP marked on both homologues was monitored in zygotes arrested prior to meiosis II via a *mes1Δ* mutation. Supporting the above results, *sgo1Δ* cells frequently showed precocious division of centromeres as split cen2-GFP signals prevailed in the dyad nuclei (FIG. 3b). Finally, it was examined whether protection of Rec8 at centromeres is dependent on Sgo1 by monitoring Rec8-GFP at late anaphase I and prometaphase II. While it is significant that Rec8 signals were centromeric in wild-type cells, the Rec8 signals had largely disappeared from the centromeres at these stages in *sgo1Δ* cells (FIG. 3c). Although all phenotypes of *sgo1Δ* cells are reminiscent of heterochromatin-deficient *Schizosaccharomyces pombe*, in which Rec8 localization to the pericentromeric regions is decreased and centromeric cohesion is lost during meiosis I, leading to random division at meiosis II (Science 300, 1152-5(2003)). Chromatin binding by Rec8 was examined in cells arrested prior to meiosis I by using a chromatin immunoprecipitation (ChIP) assay. In marked contrast to heterochromatin-deficient cells, Rec8 localization was intact in *sgo1Δ* cells at the pericentromeric regions as well as all other regions tested. These results suggest that the loss of centromeric Rec8 after meiosis I is caused not by an initial defect in Rec8 localization to centromeres but rather by a defect in the preservation of centromeric Rec8 during meiosis I. The above results indicated that the Cut1 separase becomes active at the onset of anaphase I and cleaves most chromosomal Rec8, leaving only centromeric Rec8 intact (Embo J 22, 5643-53(2003)). These results indicated that Sgo1 plays an essential role in protecting centromeric cohesion throughout meiosis I by protecting cohesin Rec8 from separase cleavage.

#### (Sgo1 Localizes at Centromeres During Meiosis I)

[0056] To detect the Sgo1 protein, Sgo1-specific antibodies were produced, and the results of Western blotting indicated that Sgo1 is expressed only around at meiosis I (FIG. 4a). The results of immunofluorescence microscopy on cells at various stages of meiosis revealed that Sgo1 appears at late prophase of meiosis I and is fully localized as several punctuate dots by the point of metaphase I (FIG. 4b). These dots were co-localized with the Mis6 kinetochore protein (Cell 90, 131-143(1997)), and indicated that Sgo1 is a centromere-associating protein (FIG. 4c). At the onset of anaphase I, Sgo1 signals decrease dramatically. It was found that Sgo1 remains undegraded at centromeres in APC-depleted cells arrested at metaphase I but undergoes normal degradation in separase-defective cells (FIG. 5), and indicated that Sgo1 degradation at anaphase I is regulated more directly by the APC rather than through separase. Although residual Sgo1 signals were detectable at the centromeres in early anaphase I, they disappeared completely by the end of anaphase I (FIG. 4b). This indicates that a substantial amount of Sgo1 is required at the onset of anaphase I when separase is fully activated. However, it is considered that the amounts of Sgo1 required are smaller and smaller as anaphase I progressed. This idea is tenable when the separase, activity is quickly down-regulated or when the access to chromosomes is prevented during anaphase I. Sgo1 never reappears during meiosis II (FIG. 4b), and which is consistent with the idea that Sgo1 is required for the protection of Rec8 only during meiosis I.

[0057] The present inventor has already reported that Rec8 localization at pericentromeric regions is especially important for the persistence of centromeric cohesion throughout meiosis I (Science 300, 1152-5(2003)). If Sgo1 is a centromeric protector of Rec8, then it might be expected to localize there as well. To test this possibility, Rec8 localization was delineated more precisely by using the ChIP assay. Sgo1 actually associated with pericentromeric heterochromatin regions rather than with central core regions along the centromere sequences (FIG. 4d). As the results of immunoprecipitation experiments indicated that Sgo1 interacts with Rec8 complexes in vivo (FIG. 4f), the protection was carried out through close interaction. Concurrently, these results indicate that Sgo1 resides at pericentromeric regions and acts to protect centromeric Rec8 from the cleavage of separase at anaphase I (FIG. 4d). It was found that the localization of Rec8 does not depend on Sgo1, and vice versa (FIG. 3d, figure not shown). Actually, the Rec8 and the Sgo1 are in fact independently generated at pericentromeric regions, as for the localization, the Rec8 and the Sgo1 depend on heterochromatin and Bub1 kinase respectively (FIG. 4e). In contrast, Rec8 and Sgo1 are localized at centromeres in *swi6Δ* (heterochromatin deficient) and *bub1Δ* cells respectively (FIG. 4e). Thus by localizing independently, it can be ensured that Rec8 is protected only at centromeres not along the chromosomal arm regions.

[0058] Further, it is indicated that shugoshin shields Rec8 physically from the action of separase and counteracts the effects. On this point, even when the strong expression of Sgo1 dose not express Rec8, the mitotic growth was moderately disturbed (figure not shown); and even when the temperature is tolerated for *cut1* allele, it was found that *cut1* mutant was killed by moderate expression of Sgo1 (FIG. 6).

#### (Sgo2 is a Mitotic Sgo1 Parologue in Fission Yeast)

[0059] By a conventional BLAST search of genome databases, the present inventor identified Sgo1-like proteins from *Saccharomyces cerevisiae* and *Neurospora crassa*, and indicated that Sgo1 is a conserved protein (see below). In the same search, a *Schizosaccharomyces pombe* Sgo1 parologue which the present inventor named Sgo2, was also identified (ORF: SPAC15A10.15). The *sgo2*+gene was disrupted, and it was identified that *sgo2Δ* cells are viable but show sensitivity to the spindle destabilizing drug thiabendazole (TBZ) (FIG. 7a). As *sgo1Δ* cells never show such a defect, this phenotype is remarkable. To investigate its cellular distribution, the endogenous *sgo2*+gene was tagged with GFP. In proliferating cells, Sgo2-GFP was observed as two or three dots in the nucleus (FIG. 7d). However, Sgo2-GFP co-localized with the centromere protein Mis6 at metaphase and disappeared during anaphase (FIGS. 7c and d). The results of ChIP assays showed that Sgo2 chromatin association is detectable only on synchronous populations of mitotic cells, and that chromatin association is localized to the pericentromeric regions (FIG. 7e). By enhancing this localization, *sgo2* deletion confers a dramatic defect to chromosome segregation when the heterochromatin-deficient *swi6Δ* mutation was bound thereto, however which by itself impairs centromeric function slightly (Science 269, 1429-31(1995)) (FIG. 7b). These results indicate that Sgo2 cooperates with centromeric heterochromatin factors to ensure chromosome segregation at mitosis. Moreover, it was found that *sgo2Δ* cells have a modest increase (up to 15%) in non-segregation of homologues at meiosis I, and indi-

cated that Sgo2 is also important for promoting proper meiosis I. However, the role of Sgo2 does not overlap with that of Sgo1, as *sgo1Δ* neither causes an apparent defect at meiosis I (FIG. 3a) nor enhances a defect of *sgo2* in meiosis. (Shugoshin Localization Controlled by Bub1)

[0060] As centromeric Rec8 cannot be detected after meiosis I in fission yeast *bub1* mutants, a conserved centromere-associated kinase Bub1 is considered to function in protecting Rec8 during meiosis, (Nat Cell Biol 3, 522-6(2001)) (FIG. 3c). Although *bub1* mutation has pleiotropic effects in meiotic chromosome segregation, it is considered that Sgo1 function can be targeted by Bub1 activity. To elucidate this problem, Sgo1-GFP signals were examined in *bub1Δ* cells undergoing meiosis. Obviously, *Bub1Δ* cells were almost completely devoid of accurate centromeric Sgo1-GFP signals, instead showed a diffuse fluorescence in the nucleus (FIG. 4e). Similar results were obtained by using the *bub1-K762R* point mutation that abolishes the kinase activity (Embo J 22, 1075-87(2003)). Although substantial levels of Sgo1 protein were detected in meiotic *bub1Δ* cells by Western blot analysis (figure not shown), Bub1 dose not influence protein stability of Sgo1. Thus, the kinase activity of Bub1 is required for incorporating Sgo1 to centromeres, and the observed defects in centromeric protection in *bub1Δ* cells can be explained by impaired localization of Sgo1.

[0061] In parallel experiments, it was identified that mitotic Sgo2 localization at centromeres was similarly disturbed in *bub1* mutants (FIG. 7c). It has been indicated that loss of Bub1 function causes centromeric function to be weakened (J Cell Biol 143, 1775-87(1998)). In this regard, the *bub1-K762R* mutation shows co-lethality with *swi6Δ*, a mutation that also slightly impairs centromeric function via its role in pericentromeric heterochromatin formation. It was found that *sgo2Δ* similarly shows co-lethality with *swi6Δ* (FIG. 7b), and exhibits severe miss-segregation of chromosomes at mitosis (figure not shown). As the *sgo2Δ bub1Δ* double mutant showed no cumulative defects at all in growth or TBZ sensitivity (FIG. 7a), Sgo2 and Bub1 tandem function was confirmed to ensure chromosome segregation in mitosis by these genetic analyses. Taken all together, the above results revealed that the incorporation of Sgo1 and Sgo2 to centromeres is a crucial function of Bub1 kinase in meiosis and mitosis, respectively.

(Characteristics of a Budding Yeast Sgo1 Homologue)

[0062] The present inventor identified a single Sgo1 homologue, ScSgo1 in budding yeast (ORF: YOR073W), which has so far not been analyzed. The cellular localization of ScSgo1 was examined by tagging endogenous ScSGO1 with GFP. ScSgo1-GFP was detected mainly as a single dot in proliferating cells, but only in a limited subset of the population (FIG. 8a). Scsgo1-GFP was not detected during the G1/S period (i.e. in cells with no bud or a small bud) but appeared as a dot in G2/M (cells with a large bud and a single nucleus) and disappeared at anaphase (cells with a large bud and a stretched nucleus) (FIG. 8a). The dot is co-localized with Ndc10 kinetochore protein (FIG. 8b). During meiosis, ScSgo1-GFP was detected at the kinetochore only at metaphase I, but never during anaphase I or meiosis II (FIG. 8c). Thus, the pattern of ScSgo1 localization closely resembles that of SpSgo2 in mitosis and SpSgo1 in meiosis.

[0063] The ScSGO1 gene was disrupted to examine the function of ScSgo1. Although the Scsgo1Δ cells were viable,

they grew slowly and showed sensitivity to the spindle destabilizing drug benomyl (FIG. 8d), and indicated that centromeric function might be impaired. And then the chromosome loss rates in Scsgo1Δ cells were compared with those in wild-type cells by a colony sectoring assay. Whereas 40% of the Scsgo1Δ colonies contained red sectors (which indicate chromosome loss), less than 2% wild-type colonies contained such sectors (FIG. 8e). It was concluded that ScSgo1 plays a crucial role at centromeres to ensure mitotic chromosome segregation. At the onset of meiosis, Scsgo1Δ cells showed significant defects that many cells are arrested with a single nucleus in the meiotic condition. However, among the leaked tetranucleate products of meiosis, the distribution pattern of cenV-GFP was consistent with proper segregation at meiosis I with the exception of random segregation at meiosis II (FIG. 8f). It was also found that tagging chromosomal ScSGO1 with 13Myc at its carboxyl terminus, which by itself causes no detectable defects in mitotic growth or meiosis I, resulted in impaired segregation at meiosis II (34% non-segregation indicates 68% random segregation)(FIG. 8g). Moreover, the ScSGO1-Myc cells showed frequent separation of sister centromeres at late meiotic anaphase I (FIG. 8h), indicated that centromeric cohesion was not properly protected. Concurrently, these results support the idea that ScSgo1 plays a crucial role in protecting centromeric cohesion throughout meiosis I, and meiosis II was ensured thereby as is the case with fission yeast Sgo1.

(Conservation of Shugoshin Among Eukaryotes)

[0064] BLAST searches identified only three Sgo1-like proteins, which were all in fungi: *Schizosaccharomyces pombe* Sgo2, *Saccharomyces cerevisiae* ScSgo1, and *Neurospora crassa* B23G1.060. As the two conserved regions were found in these proteins, the related proteins are searched under conditions of two block sequences by the BLOCK MAKER and MAST programs (Nucleic Acids Res 26,309-12(1998), Bioinformatics 14, 48-54(1998)). This approach extracted several candidate proteins from various eukaryotes including fly, worm, plant, mouse and human (see SEQ ID Nos: 21-37; *drosophila* Dm, Ce, *Arabidopsis* At, mouse Mm and human Hs, respectively, in FIG. 9). Especially, this list includes *Drosophila* ME1-S332, which is previously characterized as a protein essential for preserving centromeric cohesion in meiosis (Cell 83, 247-256(1995)), although the similarity score is marginal (E-value=10). All other proteins in the list show a short stretch of similarity in the carboxyl terminal basic regions, while the primary sequences in the first block are not conserved except that they all contain a putative coiled-coil. The space and sequences between these two blocks diverge among the proteins. As these blocks were previously identified to be important for ME1-S332 function (Genes Dev. 12, 3843-3856(1998)), the importance of the conserved regions in Sgo1 was investigated. Several amino acids were changed individually to alanines in these similarity blocks and the function of the mutant proteins in vivo was examined (FIG. 10). It was found that three conserved amino acids known to be important for ME1-S332 function were also required for Sgo1 function (13N, 34V and 368S in ME1-S332; 29N, 50I and 294S in Sgo1) (marked as arrowheads in FIG. 9). Further, other conserved amino acids in the second block (293P, 296R, 298K, 299L and 300R in Sgo1) were also all required for Sgo1 function (asterisks in FIG. 9), and non-conserved residue 297T could be changed to ala-



nine without impairing function (circle in FIG. 9). These results indicated that the marginal structural similarity observed among *Schizosaccharomyces pombe* Sgo1 and other proteins in various eukaryotes is important. Plants and mammals carry two shugoshin-like proteins, suggesting the possibility that the function of shugoshin diverges to complete mitosis and meiosis as in fission yeast.

(Proteins Encoding Human Shugoshin Homologous Gene are Specifically Localized at Centromeres in Mitosis)

[0065] The present inventor previously identified two putative human Sgo proteins, Sgo1 and Sgo2 in the database, although their overall sequence homology to known Sgo proteins in any species other than human is marginal (FIG. 11a). To examine whether these proteins identified in the database are actually human Sgo homologs, the present inventor examined the localization of the proteins. For this end, the present inventor cultured rabbit polyclonal antibodies against recombinant proteins that were produced in bacteria. The obtained Sgo1 antibodies detected an up to 70 kD band (predicted molecular weight is 60 kD) in the HeLa cell extracts, and the signal was significantly reduced when cells were treated with siRNA that targets Sgo1 mRNA (FIG. 11b). Similarly, Sgo2 antibodies detected an up to 120 kD band (predicted molecular weight is 145 kD), the signal was reduced in extracts obtained from cells treated with Sgo2 siRNA (FIG. 11b). These data indicate that both Sgo1 and Sgo2 are expressed at least in proliferating HeLa cells. Next, for the purpose of analyzing proteins (SEQ ID NOs: 18 and 20, respectively hSgo1 and hSgo2) encoding human shugoshin homologous gene (SEQ ID NOs: 17 and 19) that was presumed to be human Sgo homologues, part of hSgo1 and hSgo2 was expressed in *E. coli*, and antibodies against hSgo1 and hSgo2 were produced by injecting the protein into rabbit, HeLa cells were stained with the antibodies and concurrently with tubulin antibodies and DAPI, and co-stained with spindle and chromosome DNA respectively, and the expression of hSgo1 and hSgo2 proteins that were both endogenous in proliferating cells was examined. The results are shown in FIG. 12. As shown in FIG. 12, both signals of hsgo1 and hSgo2 were also observed as dots on chromosomes from prometaphase to metaphase. As a result of the immunostaining, it was identified that both proteins, hsgo1 and hSgo2 are specifically localized at centromeres at mitotic phase. Further, HeLa cells at prometaphase and metaphase were stained with antibodies against hsgo1 or hSgo2; concurrently co-stained with antibodies against centromere protein CENP-A, and DAPI; and examined the expression of hsgo1 and hSgo2 proteins. The results are shown in FIG. 13. As shown in FIG. 13, both signals of hSgo1 and hSgo2 were observed at sites close to CENP-A dots on chromosomes. As a result of the above, it was revealed that both hsgo1 and hSgo2 are centromere proteins. Further, to examine this possibility, Aurora B, which is a passenger protein of chromosome known to be localized within kinetochore from prophase to metaphase, was stained. The sites of Sgo1 and Aurora B were practically the same at prometaphase and metaphase, whereas Sgo2 was placed just outside Aurora B (see FIG. 13). As a result of the above, it was revealed that both hsgo1 and hSgo2 are placed within kinetochores from prometaphase to metaphase. Representative views of sister kinetochore are magnified on the right. Scale bar is 10  $\mu$ m.

(Proteins Encoding Human Shugoshin Homologous Gene are Specifically Localized at Centromeres in Mitosis and Play an Important Role to Progress Chromosome Segregation)

[0066] RNAi experiments targeting hsgo1 and hSgo2 were performed respectively. The results are shown in FIG. 14. As a result, the expressions in any proteins were significantly suppressed 48 hours later, the cells arrested in mitosis (total, in figure) were accumulated as indicated in FIG. 14. As described above, it was strongly suggested that any protein localized at centromeres in mitosis plays an important role for progressing chromosome segregation. As the accumulation was dissolved by suppressing a spindle checkpoint factor BubR1 by RNAi, it was suggested that hsgo1 and hSgo2 are directly or indirectly function during the process where spindle properly takes the kinetochore at centromeres as described below.

[0067] Further, the cells for which RNAi experiments targeting hsgo1 was performed by using HeLa cells were mounted on a slide glass and stained with Giemsa. The results are shown in FIG. 15. It was revealed that sister chromatid at prophase strongly adhered at centromere site in control cells where RNAi was not performed; while in cells suppressing hsgo1 expression, where RNAi was performed, the adhesion was weak at centromere site, and easily detached. Consequently, it was demonstrated that hsgo1 has an important role to maintain the strong cohesion at centromere site in mitosis in proliferating cells. Mitotic cells where Sgo1 protein knockdown was performed by RNAi experiments were collected, and the chromosomes were spread to observe chromosome structure directly. In control cells, sister chromatids were resolved along the arm regions but showed the primary constriction at centromeres (FIG. 16a i). Amazingly, in Sgo1-depleted cells, sister chromatids were often separated along the whole chromosome length (FIG. 16a iii). In samples where sister chromatids stayed densely close, although sister chromatids did not indicate the primary constriction (FIG. 16a iv), this suggests that centromeric cohesion was lost selectively. Nocodazole treatment activates the spindle checkpoint; thereby the cell cycle is arrested at prometaphase. Such prolonged arrest in M phase causes the complete separation of the connectivity from the chromosomal arm regions. For this reason, sister chromatids are only connected at centromeres, and form 'Xshaped' chromosome (FIG. 16b, control). As expected, nocodazole-treatment caused the complete separation of sister chromatids along the chromosome length in Sgo1 RNAi cells (up to 97%) (FIGS. 16c and d). Consequently, it was demonstrated that hSgo1 plays an important role to maintain the strong cohesion at chromosomal centromere site in mitosis in proliferating cells.

[0068] RNAi experiments targeting Bub1 were performed respectively. The results are shown in FIG. 17. Consequently, the localization of either protein of the hsgo1 and hSgo2 to centromere was disappeared. This result means that the conclusion, "localization of shugoshin to centromere depends on Bub1 kinase", which was found in yeast by the present inventor, is also conserved in higher organisms.

[0069] Next, clone where cDNA of mouse shugoshin homologous genes (SEQ ID NOs: 21 and 23) was fused with GFP gene was produced by using retroviral vector and expressed in human HeLa cells. The results are shown in

FIG. 18. Consequently, it was revealed that any of the GFP fusion proteins are also co-localized with human kinetochore protein Bub1 in mitosis.

[0070] The analysis of the above hsgo1 and hSgo2 and the analysis results obtained with the use of mouse shugoshin homologous genes were strongly suggested that shugoshin-like protein in animal cells, which were predicted from the sequence, also have functional conservation with yeast shugoshin.

## INDUSTRIAL APPLICABILITY

[0071] Shugoshin of the present invention that is a regulatory factor of chromosome segregation widely conserved in eukaryotic cells, can be advantageously used for studies on the induction mechanism of cancer in somatic division, the chromosome segregation diseases such as infertility or Down's syndrome in meiotic division, and the like besides on the elucidation of mechanism in chromosome segregation.

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 45

<210> SEQ ID NO 1  
<211> LENGTH: 960  
<212> TYPE: DNA  
<213> ORGANISM: yeast

<400> SEQUENCE: 1

```

atgaactttc aatttataaa ttcaaatata aacaatgaag ataaattgcc gatggagtgc 60
ttgaaaaaga aattttataa acaaaatcgt gaaattataa aaataaatac tcagctttct 120
ataaaaatta gagaatctga aaacgaaatt caagatttga tacaagaaaa tttcactttg 180
aaaagttatt tggtaaact tgaagctcga tttcgcaatc aatctcaaac tgaggacttg 240
ttaaaaaact tctttcctga gatacaaac attcacaaaa agatttcaca agtgcaaagt 300
ttactgaaga ttatagagaa aaagtgttca tcagatttcc tcgaagcgaa tgtaaaaagt 360
caatttacia cctgtgaaaa taaagattcg aaagaagatt atcagatttt gcataataaa 420
cgcttgagat atgtatcatt taatgatgaa cttaaaagtc tcgaaacagg gcaaccattg 480
tattgttttc aagatttcca aaaaaaagtc catggtcctc cggtctctac tgaaaaacct 540
ggaaaatgta tattaaaaga taaaaccaat gccacgtaa acaaaatacc acaagatgag 600
gtgaattact cattgccgca aaaaaatc accatctttt caaaggaatt aaaagaaaac 660
gaatttgaat ccatcaacga gggcgaaact gaagaagaaa aggctaaaac atcaaatggt 720
tgtgtttgta ttccctgtaa aagtgtgaa cagataactg accttaaagg acaagcaacc 780
ggagacagct ccccatgtga ttttgaagaa tctcaaccaa ggattaatgg acgtgaaaaa 840
ctaagacgat cagtcaaagt gataaactat gcaataccca gtttgcgaac taaactacga 900
cgagactttg acttaccatc tgatagaaaa cgcaaacgac atcccagagg caaagcataa 960

```

<210> SEQ ID NO 2  
<211> LENGTH: 319  
<212> TYPE: PRT  
<213> ORGANISM: yeast

<400> SEQUENCE: 2

```

Met Asn Phe Gln Phe Ile Asn Ser Asn Ile Asn Asn Glu Asp Lys Leu
1           5           10           15
Pro Met Glu Ser Leu Lys Lys Lys Phe Leu Lys Gln Asn Arg Glu Ile
                20                25                30
Ile Lys Ile Asn Thr Gln Leu Ser Ile Lys Ile Arg Glu Ser Glu Asn
35                40                45
Glu Ile Gln Asp Leu Ile Gln Glu Asn Phe Thr Leu Lys Ser Tyr Leu
50                55                60

```

-continued

---

Val Lys Leu Glu Ala Arg Phe Arg Asn Gln Ser Gln Thr Glu Asp Leu  
 65 70 75 80  
 Leu Lys Asn Phe Phe Pro Glu Ile Gln Thr Ile His Lys Lys Ile Ser  
 85 90 95  
 Gln Val Gln Ser Leu Leu Lys Ile Ile Glu Lys Lys Cys Ser Ser Asp  
 100 105 110  
 Phe Leu Glu Ala Asn Val Lys Ser Gln Phe Thr Thr Cys Glu Asn Lys  
 115 120 125  
 Asp Ser Lys Glu Asp Tyr Gln Ile Leu His Asn Lys Arg Leu Glu Tyr  
 130 135 140  
 Val Ser Phe Asn Asp Glu Leu Lys Ser Leu Glu Thr Gly Gln Pro Leu  
 145 150 155 160  
 Tyr Cys Phe Gln Asp Phe Gln Lys Lys Val His Gly Pro Pro Ala Leu  
 165 170 175  
 Ser Glu Lys Pro Gly Lys Cys Ile Leu Lys Asp Lys Thr Asn Ala His  
 180 185 190  
 Val Asn Lys Ile Pro Gln Asp Glu Val Asn Tyr Ser Leu Pro Gln Lys  
 195 200 205  
 Asn Ile Thr Ile Phe Ser Lys Glu Leu Lys Glu Asn Glu Phe Glu Ser  
 210 215 220  
 Ile Asn Glu Gly Glu Thr Glu Glu Glu Lys Ala Lys Thr Ser Asn Val  
 225 230 235 240  
 Cys Val Cys Ile Pro Cys Lys Ser Ala Glu Gln Ile Thr Asp Leu Lys  
 245 250 255  
 Gly Gln Ala Thr Gly Asp Ser Ser Pro Cys Asp Phe Glu Glu Ser Gln  
 260 265 270  
 Pro Arg Ile Asn Gly Arg Glu Lys Leu Arg Arg Ser Val Lys Val Ile  
 275 280 285  
 Asn Tyr Ala Ile Pro Ser Leu Arg Thr Lys Leu Arg Arg Asp Phe Asp  
 290 295 300  
 Leu Pro Ser Asp Arg Lys Arg Lys Arg His Pro Arg Gly Lys Ala  
 305 310 315

<210> SEQ ID NO 3  
 <211> LENGTH: 1944  
 <212> TYPE: DNA  
 <213> ORGANISM: yeast

<400> SEQUENCE: 3

```

atgtcgaaaag catctctttc cccgaacgta gaagacttga aaaaaagca aattcgacag    60
tataagaaaa ttatacgaat aagcaaggca caatcaatta gaattaaaga attgcagtta    120
gaaaatgaac ggttgctttc ggaaaatc gatttgagga ctacagcgat aaacttgaa    180
gagcaactcg aaacogtgc aacgaaaaac gaagaaaaca aaacaaagt agctgcatta    240
cttaatcgat ttcataaga aacagataat tttttatcaa aattaagtct ttgtcagcaa    300
gaaatacaag acacctcaa accagtggag gctaacttag cttacgatgt cgatacggat    360
tctgaagacc ttgaogagga atcogtctgt aaagataccg aagaaataat tgagcaagct    420
cagcatgatg tttccttacg aaatthaagt ggaatagagg atgaaaatat aattgatgac    480
ggagaaaactg ctataaatga acaaaaaaaaa agagaagcta atgttttttc cgacacgcaa    540
tcagcacctc agctaaaatc cggcaaaagc ctcccagctg attttgaaaa tccttaaat    600
  
```

-continued

---

```

ctatccaatt cgaaacctgt aaataataat aatgaagata gagttgaagc ggttacttct 660
gaaaataaat ctatcgattc tgctcctcag gaaaaaaatc atgaatacga aatcgttagt 720
ccaaaatcat tatccaacaa aattaataat caagcagctg cacaaagaag aaccgaagaa 780
gataatgcaa atggagttgc tcaagaagaa aatgagggtt cacaagaagc tcattttcat 840
agcagaatac aatctgatac agtaatacaa agtacacca ctaaaccgaa atgggacgtt 900
gacattcaaa ataaacaaat taatctggct tctgcagcta ccaatgttac cggttatgta 960
tcggagaccg atagtcgccc caatcgcgca aactccttgg attctgctgt ccttcttctg 1020
caatcttcaa ataaaagtaa ccgaaatggg catcatatct cagatcctaa tttaaatagc 1080
tccatatcgt tgaagtttgc gcctgaagat actgcgcata attcattaac ttcacaagag 1140
aatgttgggc ctcaggttac gacgacttct ctgtcaaata tgactgttgc tgaatctcct 1200
cgtacagaca ctccaagga aataaacggg ttagtagact cttctgtcac taatgggaac 1260
gaaaatctt ctgtagaat aatgaatgac tctaacaaaa ttggactgaa tcctaaatct 1320
tttaccgacg aagagcggga aattttaaca ctttttcgaa atcctccat gagactgtca 1380
agtgaacctc catcttcaaa tggattttca atagcccatc ccaataattc tccgttacgt 1440
ccgcatcgcg tacaaggaat attgaatgct gaagatcgac cttacgaaat tgagccgtca 1500
cgtagctcct ttgctaccaa cgatcgggc tcctataata atttggaaact tctgtcatct 1560
gtaacgaatt tgaatcccc taatgagaac gatcgtgtga cgaaaactca gtcgcaaga 1620
gaaacaaaag tgaaaaggcg aagaaaagct cggattcaag aaacttctga agaaagtaca 1680
gtagtcaatg agccaaatga aaaacctgat ggaaggagcc gaagggaacg gaaaaaggtt 1740
aattacgctt tgcttgatt aaggacgaaa ttaagacgga atttcgattt accttcagat 1800
catgtaaaag ctaaaaaaac gagacgtgct cctaagaact ctgagaatga ttcagctacc 1860
aaaacagaaa ccgaaacat tacttctgaa gcaccacta cttcagaagt aaccctgaa 1920
aactccgaaa cccttaattt gtaa 1944

```

```

<210> SEQ ID NO 4
<211> LENGTH: 647
<212> TYPE: PRT
<213> ORGANISM: yeast

```

```

<400> SEQUENCE: 4

```

```

Met Ser Lys Ala Ser Leu Ser Pro Asn Val Glu Asp Leu Lys Lys Lys
1           5           10          15
Gln Ile Arg Gln Tyr Lys Glu Ile Ile Arg Ile Ser Lys Ala Gln Ser
20          25          30
Ile Arg Ile Lys Glu Leu Gln Leu Glu Asn Glu Arg Leu Leu Ser Glu
35          40          45
Asn Ile Asp Leu Arg Thr Thr Ala Ile Asn Leu Glu Glu Gln Leu Glu
50          55          60
Thr Val Gln Asn Glu Asn Glu Glu Asn Lys Thr Lys Leu Ala Ala Leu
65          70          75          80
Leu Asn Arg Phe His Glu Glu Thr Asp Asn Phe Leu Ser Lys Leu Ser
85          90          95
Leu Cys Gln Gln Glu Ile Gln Asp Thr Phe Lys Pro Val Glu Ala Asn
100         105         110

```

-continued

---

Leu Ala Tyr Asp Val Asp Thr Asp Ser Glu Asp Leu Asp Glu Glu Ser  
 115 120 125  
 Val Val Lys Asp Thr Glu Glu Ile Ile Glu Gln Ala Gln His Asp Val  
 130 135 140  
 Ser Leu Arg Asn Leu Ser Gly Ile Glu Asp Glu Asn Ile Ile Asp Asp  
 145 150 155 160  
 Gly Glu Thr Ala Ile Asn Glu Gln Lys Lys Arg Glu Ala Asn Val Phe  
 165 170 175  
 Ser Asp Thr Gln Ser Ala Pro Gln Leu Lys Ser Gly Lys Ala Leu Pro  
 180 185 190  
 Ala Asp Phe Glu Asn Pro Tyr Asn Leu Ser Asn Ser Lys Pro Val Asn  
 195 200 205  
 Asn Asn Asn Glu Asp Arg Val Glu Ala Val Thr Ser Glu Asn Lys Ser  
 210 215 220  
 Ile Asp Ser Ala Pro Gln Glu Lys Asn His Glu Tyr Glu Ile Val Ser  
 225 230 235 240  
 Pro Lys Ser Leu Ser Asn Lys Ile Asn Asn Gln Ala Ala Ala Gln Arg  
 245 250 255  
 Arg Thr Glu Glu Asp Asn Ala Asn Gly Val Ala Gln Glu Glu Asn Glu  
 260 265 270  
 Gly Ser Gln Glu Ala His Phe His Ser Arg Ile Gln Ser Asp Thr Val  
 275 280 285  
 Ile Gln Ser Thr Pro Thr Lys Arg Lys Trp Asp Val Asp Ile Gln Asn  
 290 295 300  
 Lys Gln Ile Asn Leu Ala Ser Ala Ala Thr Asn Val Thr Gly Tyr Val  
 305 310 315 320  
 Ser Glu Thr Asp Ser Arg Pro Asn Arg Ala Asn Ser Leu Asp Ser Ala  
 325 330 335  
 Val Leu Leu Val Gln Ser Ser Asn Lys Ser Asn Arg Asn Gly His His  
 340 345 350  
 Ile Ser Asp Pro Asn Leu Asn Ser Ser Ile Ser Leu Lys Phe Ala Pro  
 355 360 365  
 Glu Asp Thr Ala His Asn Ser Leu Thr Ser Gln Glu Asn Val Gly Pro  
 370 375 380  
 Gln Val Thr Thr Thr Ser Leu Ser Asn Met Thr Val Ala Glu Ser Pro  
 385 390 395 400  
 Arg Thr Asp Thr Pro Arg Glu Ile Asn Gly Leu Val Asp Ser Ser Val  
 405 410 415  
 Thr Asn Gly Asn Glu Lys Phe Ser Val Glu Ile Met Asn Asp Ser Asn  
 420 425 430  
 Lys Ile Gly Leu Asn Pro Lys Ser Phe Thr Asp Glu Glu Arg Glu Ile  
 435 440 445  
 Leu Thr Leu Phe Arg Asn Pro Pro Met Arg Leu Ser Ser Glu Pro Pro  
 450 455 460  
 Ser Ser Asn Gly Phe Ser Ile Ala His Pro Asn Asn Ser Pro Leu Arg  
 465 470 475 480  
 Pro Pro Ser Leu Gln Gly Ile Leu Asn Ala Glu Asp Arg Pro Tyr Glu  
 485 490 495  
 Ile Glu Pro Ser Arg Ser Ser Phe Ala Thr Asn Asp Thr Gly Ser Tyr  
 500 505 510  
 Asn Asn Leu Glu Leu Leu Ser Ser Val Thr Asn Leu Lys Ser Pro Asn

-continued

---

515	520	525													
Glu Asn Asp Arg Val Thr	Lys Thr Gln Ser Arg	Arg Glu Thr Lys Val													
530	535	540													
Lys Arg Arg Arg Lys Ala Arg Ile Gln Glu Thr Ser Glu Glu Ser Thr															
545	550	555	560												
Val Val Asn Glu Pro Asn Glu Lys Pro Asp Gly Arg Ser Arg Arg Glu															
	565	570	575												
Arg Lys Lys Val Asn Tyr Ala Leu Pro Gly Leu Arg Thr Lys Leu Arg															
	580	585	590												
Arg Asn Phe Asp Leu Pro Ser Asp His Val Lys Ala Lys Lys Thr Arg															
	595	600	605												
Arg Ala Pro Lys Asn Ser Glu Asn Asp Ser Ala Thr Lys Thr Glu Thr															
	610	615	620												
Ala Asn Ile Thr Ser Glu Ala Pro Thr Thr Ser Glu Val Thr Leu Glu															
625	630	635	640												
Asn Ser Glu Thr Leu Asn Leu															
	645														

<210> SEQ ID NO 5  
 <211> LENGTH: 1773  
 <212> TYPE: DNA  
 <213> ORGANISM: yeast

<400> SEQUENCE: 5

```

atgccgaaga gaaaaattgc tcctaacaag gaaagcagca ggcgtacggt ctcccacgat    60
gatttaaccc cacaataca agaatttcaa aacctaatgg atctogaatc gcaaaaagtg    120
gaaaacatca gacagtcgta ttcgaggcaa aactccctgc tggccaagga taactccata    180
ttaaaaatta aagttaatag cttggaaaaa aaaataagcc agctggtaca agaaaacgtg    240
actctacgat ctaaaacctc tataagcgaa gctatctaca gggaaacggt aagtaatcaa    300
ctacaagtca ttgaaaacgg tattattcaa agatttgacg aaatTTTTta tatgtttgag    360
aacgtacgta aaaacgaaaa tttgccagc tcgagcttaa gaacaatggt gaagagaacg    420
agttccaggt caagatcatg ctcatgttca tcaccocat actcaaaaag ttacactagg    480
ttatcaaadc acgagaataa cctgtcgcac gaatcaagtt ttaacaagga cgatggtcca    540
gatcttgagc ctaaggctaa aaaaaggaag agttctaggc ggcaatctat gtttgtatcc    600
acgagtttag aacctgaaga cgaaacgggt gaaaaogaac ccatgatgga aaattcctct    660
gtagaggtag cggcagaatc acacgagtct gcgcaagtgg aggaacaat agatgcctta    720
aacctgaag aggaaaatag cgattctgtc agtaatttta ccaattcaat tatagaatac    780
tccataccag aggagaatcc gacagaaccc gagcattcat cttctaaact agaaatattc    840
aatgacagta caaatatgct aagtacagtg cctgcaaatc ctttgccggt gcctttacca    900
ggccatccg caactttacc tactaccact agcgatgctt caacggtota tccttcatca    960
agttcttcta ctaatttca tccaaagacc aaaattaagc attccatgaa gccgcctagg    1020
atagaactga agaaaaagg tattgacgaa gtcattgccc taagtaacat gagcagcaac    1080
agcgaatat catttacgag aactagaaga actcgtggtg aagctgtaga ttacactttg    1140
ccttctttaa gagccaaat gaggaggcct tcagaaaaac ttgtggatgc tactactgtg    1200
attgatatac atgatctaca ggtttccaag agaaatcggg aaacttcaca taaaaggaaa    1260
    
```

-continued

```

agtttatccc aagattcaat acccgacgaa cgcgaattga gagaagtcgt cgtctcaaag 1320
gattatggaa ctccaaaagg gaaaaaacg gaagatgaaa tacacgagga taccgctcat 1380
ctaatgacca cttccaacaa caacagcaac aacaaaaacg aaaaaaaact aactagcaac 1440
aatagcccta aaaaatcgtc gcctttactt gacattacaa ataaatcgga gaataagaaa 1500
aagtcaacaa gaactaaaaa attgttcaaa aatgcaattg tcaataatth atctgatgaa 1560
aattctacta cgcgacctc caagtcgtca aaggaacca gtaataataa caacaattac 1620
aacaatttcg acaataacaa ttcaaacatt aataatgta atataaatc tgttagcttt 1680
agactaaatg aagatgattt agcagtattt gatttatttg gaaatgtaa ggcagtgaaa 1740
catcaaccaa aaacatatcg caccaaaaaa tga 1773

```

```

<210> SEQ ID NO 6
<211> LENGTH: 590
<212> TYPE: PRT
<213> ORGANISM: yeast

```

```

<400> SEQUENCE: 6

```

```

Met Pro Lys Arg Lys Ile Ala Pro Asn Lys Glu Ser Ser Arg Arg Thr
1           5           10           15
Val Ser His Asp Asp Leu Thr Pro Gln Ile Gln Glu Phe Gln Asn Leu
20           25           30
Met Asp Leu Glu Ser Gln Lys Val Glu Asn Ile Arg Gln Ser Tyr Ser
35           40           45
Arg Gln Asn Ser Leu Leu Ala Lys Asp Asn Ser Ile Leu Lys Ile Lys
50           55           60
Val Asn Ser Leu Glu Lys Lys Ile Ser Gln Leu Val Gln Glu Asn Val
65           70           75           80
Thr Leu Arg Ser Lys Thr Ser Ile Ser Glu Ala Ile Tyr Arg Glu Arg
85           90           95
Leu Ser Asn Gln Leu Gln Val Ile Glu Asn Gly Ile Ile Gln Arg Phe
100          105          110
Asp Glu Ile Phe Tyr Met Phe Glu Asn Val Arg Lys Asn Glu Asn Leu
115          120          125
Pro Ser Ser Ser Leu Arg Thr Met Leu Lys Arg Thr Ser Ser Arg Ser
130          135          140
Arg Ser Cys Ser Leu Ser Ser Pro Thr Tyr Ser Lys Ser Tyr Thr Arg
145          150          155          160
Leu Ser Asn His Glu Asn Asn Leu Ser His Glu Ser Ser Phe Asn Lys
165          170          175
Asp Asp Gly Pro Asp Leu Glu Pro Lys Ala Lys Lys Arg Lys Ser Ser
180          185          190
Arg Arg Gln Ser Met Phe Val Ser Thr Ser Leu Glu Pro Glu Asp Glu
195          200          205
Thr Gly Glu Asn Glu Pro Met Met Glu Asn Ser Ser Val Glu Val Pro
210          215          220
Ala Glu Ser His Glu Ser Ala Gln Val Glu Glu Thr Ile Asp Ala Leu
225          230          235          240
Asn Pro Glu Glu Glu Asn Ser Asp Ser Val Ser Asn Phe Thr Asn Ser
245          250          255
Ile Ile Glu Tyr Ser Ile Pro Glu Glu Asn Pro Thr Glu Pro Glu His
260          265          270

```

-continued

Ser Ser Ser Lys Leu Glu Ile Phe Asn Asp Ser Thr Asn Met Leu Ser  
 275 280 285  
 Thr Val Pro Ser Asn Pro Leu Pro Leu Pro Leu Pro Gly Pro Ser Ala  
 290 295 300  
 Thr Leu Pro Thr Thr Thr Ser Asp Ala Ser Thr Val Tyr Pro Ser Ser  
 305 310 315 320  
 Ser Ser Ser Thr Asn Ser His Pro Lys Thr Lys Ile Lys His Ser Met  
 325 330 335  
 Lys Pro Pro Arg Ile Glu Leu Lys Lys Lys Val Ile Asp Glu Val Met  
 340 345 350  
 Pro Val Ser Asn Met Ser Ser Asn Ser Glu Ile Ser Phe Thr Arg Thr  
 355 360 365  
 Arg Arg Thr Arg Gly Lys Ala Val Asp Tyr Thr Leu Pro Ser Leu Arg  
 370 375 380  
 Ala Lys Met Arg Arg Pro Ser Glu Lys Leu Val Asp Ala Thr Thr Val  
 385 390 395 400  
 Ile Asp Ile His Asp Leu Gln Val Ser Lys Arg Asn Arg Glu Thr Ser  
 405 410 415  
 His Lys Arg Lys Ser Leu Ser Gln Asp Ser Ile Pro Asp Glu Pro Gln  
 420 425 430  
 Leu Arg Glu Val Val Val Ser Lys Asp Tyr Gly Thr Pro Lys Gly Lys  
 435 440 445  
 Lys Thr Glu Asp Glu Ile His Glu Asp Thr Ala His Leu Met Thr Thr  
 450 455 460  
 Ser Asn Asn Asn Ser Asn Asn Lys Asn Glu Lys Lys Leu Thr Ser Asn  
 465 470 475 480  
 Asn Ser Pro Lys Lys Ser Ser Pro Leu Leu Asp Ile Thr Asn Lys Ser  
 485 490 495  
 Glu Asn Lys Lys Lys Ser Thr Arg Thr Lys Lys Leu Phe Lys Asn Ala  
 500 505 510  
 Ile Val Asn Asn Leu Ser Asp Glu Asn Ser Thr Thr Arg Pro Ser Lys  
 515 520 525  
 Ser Ser Lys Gly Thr Ser Asn Asn Asn Asn Tyr Asn Asn Phe Asp  
 530 535 540  
 Asn Asn Asn Ser Asn Ile Asn Asn Val Asn Asn Lys Ser Val Ser Phe  
 545 550 555 560  
 Arg Leu Asn Glu Asp Asp Leu Ala Val Phe Asp Leu Phe Gly Asn Gly  
 565 570 575  
 Lys Ala Val Lys His Gln Pro Lys Thr Tyr Arg Thr Lys Lys  
 580 585 590

&lt;210&gt; SEQ ID NO 7

&lt;211&gt; LENGTH: 2325

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Neurospora crassa

&lt;400&gt; SEQUENCE: 7

```

atggcccgcc tcaagaaca agccatgtcg tctgtcgcgt tgtcaacaga caatctogag    60
ctcctgcgta ggaagttcct cagacaaaac agagatattg ctcgagtcaa ttccacacag    120
tcaactccgta tccgtggggtt ggagaatgaa tgcgctcgtt tgctgtcggg aaacctcgaa    180
ctccgtgtgc aggtottgcg cctcgaaaag gagctccaag acaacgctgc gcgaagggtg    240

```



-continued

---

```

gccgatcatg cgctcgaggt caaggccaag atggagacgc agttggcggg actcagttcg 300
ctgctggcaa gcttagggga gccgccctcg aagcggcgcc tttcagaaga gaggcgatac 360
gcgcagcctc gaccgagcgt tcaccggagc cctcccttac gaagagcagc ccaggaggcc 420
gaccaggaac tactggctga gcaggaagga aggctaccgc cgatatacga gaacaagacg 480
tatgctcgag ccacaatgaa cagtgaagaa atcctggcgc tgtgcatgca ggcagacgat 540
tcgaatgact cgccagatat cggaccgccg ccagtatcta ggttgtcga ggatgatatg 600
gtcatacctt gttaccatc gccaaacaag aacgccgagg ctgaagaaac ggaaactacc 660
gagcaagtgg aagagagccc tagggctctt caagtaccgc cgtcattatc gccgcctaaa 720
ctggactacg acaggagacc aaacatgatc ctattcagcc caccocaaaga atcgagagtg 780
gcagaacctt ccaaatggtt cagtccccct ccgatggaac caccgaaaca gtccacatcg 840
gctgtaccga gtgagacaat acgagcaggc ctcaagcgaa agttgaacgg cgacaaccaa 900
aacgaacca acaaggcaac caagcttcaa caaggaaagg agaatggcaa tgagactggg 960
atcaagaaa gactctctgc ccgcgacccg cacaagagga aaagcatcaa agagaccgca 1020
acgaaaccga gagccccgct gtcagcaaag agcacgaacg agcacattgt ctctccgaag 1080
aagccggcga agccccacca agtggccgac gattttaagc cgggtgaagg gcacaaggcg 1140
tcaaagggta aagagaaagt cgacctgccc gctccggaca agaagtcagc agtagaagaa 1200
acgcaaggaa attctacgtc ggcattcacg aaagtcgaga tcctcccgcc ggctctggaa 1260
cctactcctg aagttgcaga gattcctgaa accgatattc tgatcacacc tggaacacca 1320
gagcgcgct ctgaaagcac tgtgtgacc cagcataccc cgccgccagc ccacatttca 1380
tccaatggag agagctcgcg gcctagcagc cgtgctagag cggctatcag ctatacagag 1440
cccaatctgc gcgacaagat gcgacgaccg accaaagagc tccttgatgc cgtttctggg 1500
gagggcaagt tcctacacag gccgacatcg caacagcaac agcagcaacg caaggcgac 1560
gagtcagcac cgagctcagt tagcaaggtc aaggctcagc catcgccggc ggtggatata 1620
agtagtctga ccagcagtcg gctgttgaa aaagagaagg agaaggaacc acagccggat 1680
gaaggaatat tatctccaaa cggcatcctc ccaagctcag tagacctggg aaggagaaga 1740
cgccctcat ccttctctac tgcagcccct gcaatgacaa tccttcggt ccaagaacaa 1800
tcaactctaa acctccagc cgcgacgag accgatgaaa acgcccggg cgaggctcag 1860
attcagaagg agctgagtaa tagtattaca acacggccca ggggtggaaa ggggaggcaa 1920
tcaatgagcc gttcogtacc cacgatocca acagaaaatt acgagcacga ggacgcacaa 1980
ctctcgacga actcagcctc ggtggatctt tacgactttg ctagtgtgc gtctccggat 2040
agcgcagcac cccagctaga agcagctacc ggcgatgttc ctgtaataa gaaggcacc 2100
aaagttcaa gaagagcgtc ctacagctgt tcgaccgaga caacagcaac agcatccgca 2160
aagccaagat cttccgaaa aagggcttcg atgctggtgc cgaagaaaag cttgtgggct 2220
gaagagttag cgcaggagga agaggatgag gaagatgctg gcaatgacag tggcgggtcc 2280
ttgtccaagg ggaggcctc gaggaggaga agcatgatgc tttga 2325

```

<210> SEQ ID NO 8

<211> LENGTH: 774

<212> TYPE: PRT

<213> ORGANISM: Neurospora crassa

-continued

&lt;400&gt; SEQUENCE: 8

Met Ala Arg Leu Asn Glu Gln Ala Met Ser Ser Val Ala Leu Ser Thr  
1 5 10 15  
Asp Asn Leu Glu Leu Leu Arg Arg Lys Phe Leu Arg Gln Asn Arg Asp  
20 25 30  
Ile Ala Arg Val Asn Ser Thr Gln Ser Leu Arg Ile Arg Gly Leu Glu  
35 40 45  
Asn Glu Cys Ala Arg Leu Leu Ser Glu Asn Leu Glu Leu Arg Gly Gln  
50 55 60  
Val Leu Arg Leu Glu Lys Glu Leu Gln Asp Asn Ala Ala Arg Arg Val  
65 70 75 80  
Ala Asp His Ala Leu Glu Val Lys Ala Lys Met Glu Thr Gln Leu Ala  
85 90 95  
Glu Leu Ser Ser Leu Leu Ala Ser Leu Gly Glu Pro Pro Ser Lys Arg  
100 105 110  
Arg Leu Ser Glu Glu Arg Arg Tyr Ala Gln Pro Arg Pro Ser Val His  
115 120 125  
Arg Ser Pro Pro Leu Arg Arg Ala Arg Gln Glu Ala Asp Gln Glu Leu  
130 135 140  
Leu Ala Glu Gln Glu Gly Arg Leu Pro Pro Ile Tyr Glu Asn Lys Thr  
145 150 155 160  
Tyr Ala Arg Ala Thr Met Asn Ser Glu Glu Ile Leu Ala Leu Cys Met  
165 170 175  
Gln Ala Asp Asp Ser Asn Asp Ser Pro Asp Ile Gly Pro Pro Pro Val  
180 185 190  
Ser Arg Phe Val Glu Asp Asp Met Val Ile Pro Cys Ser Pro Ser Pro  
195 200 205  
Asn Lys Asn Ala Glu Ala Glu Glu Thr Glu Thr Thr Glu Gln Val Glu  
210 215 220  
Glu Ser Pro Arg Ala Leu Gln Val Pro Pro Ser Leu Ser Pro Pro Lys  
225 230 235 240  
Leu Asp Tyr Asp Arg Arg Pro Asn Met Ile Leu Phe Ser Pro Pro Lys  
245 250 255  
Glu Ser Arg Val Ala Glu Pro Ser Lys Met Phe Ser Pro Pro Pro Met  
260 265 270  
Glu Pro Pro Lys Gln Ser Thr Ser Ala Val Pro Ser Glu Thr Ile Arg  
275 280 285  
Ala Gly Leu Lys Arg Lys Leu Asn Gly Asp Asn Gln Asn Glu Pro Asn  
290 295 300  
Lys Ala Thr Lys Leu Gln Gln Gly Lys Glu Asn Gly Asn Glu Thr Gly  
305 310 315 320  
Ile Lys Lys Gly Leu Ser Ala Arg Asp Pro His Lys Arg Lys Ser Ile  
325 330 335  
Lys Glu Thr Ala Thr Lys Pro Arg Ala Pro Leu Ser Ala Lys Ser Thr  
340 345 350  
Asn Glu His Ile Val Ser Pro Lys Lys Pro Ala Lys Pro His Gln Val  
355 360 365  
Ala Asp Asp Phe Lys Pro Val Lys Val His Lys Ala Ser Lys Gly Lys  
370 375 380  
Glu Lys Val Asp Leu Pro Ala Pro Asp Lys Lys Ser Ala Val Glu Glu



-continued

---

```

<211> LENGTH: 1671
<212> TYPE: DNA
<213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 9
atggttcgag cgacggttct gaatgctcgg gatcacgccca gtgaagggtg gcgtactaac    60
aaagctaaag gagagaaaat ggttctggaa cctccgatga acagtgcaca aagacgaaag    120
ttgggggata ttactaattt gcagaatcag aagaatctaa tgaatcaggg agcgaagcat    180
cagcaacaag ctatattaat ctcttctaaa gaaaacgctg aaaatcttca aaaggcactg    240
agaaattctt ctgaaaacac aaagctgatg aaagtcgtca tggagagaga tggaatcaaa    300
agtgatctga agaaacttag gattgaattt cagaaggttc aagaacagaa tttgctactt    360
gcccaggcta acactcgtat cttggcctg aaggtacttc agcacgaact tggttgcaag    420
aatgggtag  tcatggccag gaaaatgctg cttaaggctc aagcaaatgc ttgtggtggg    480
gcttgcaaaa cctttcagcc aaatgatgca gatcatgagc atgcttccgg gagctccaac    540
gctaactcat tgcaagaaa tgagaaagcc aacagtaaaa ggagagtctt tggaaaggaag    600
aatcccgcca attccgaggt attagatata attggcagat cgggagagac atgtcagatg    660
gaagacaaca ttgacaacaa gaagttggtc tctgatagtg acaatgatgc tgaaaacctat    720
ataaatgaca atgtccaaag caaaagatat tgtgcaggaa gacagagtag cagttctaaag    780
actcgagaag ccagccaaac agaaaccttg caaaagggtg ttgacgcca aaaaattaag    840
gggatgcaa  ggTTTTTTTt gacaaagcat tctgactggt taaaatctca agaacctgag    900
ccatctgaaa gcctatacga gtcaagggtc cttttgagaa ggcgttctgc ccggttaaaa    960
tctcaagaac ctgagccatc tgaagcttc catgactcaa tagagacaac caagaggagg    1020
aggtcggcaa taaggtctgc tatgtttaat atccaagagc tgggcgttat taaaacttg    1080
aacggtttac ctgatgatca agagattgct gcaaaggcca gatgctctgc acgtgaacag    1140
tctaccgggt ctaaaccgca agcagtagaa ccacatgaca caaagagat aatcgggaaa    1200
agcaggatat ctttgagaag acagtctgag aggtttaatt tccaagagct gggcgtgact    1260
gaaaacttga atggtccaca tgatgatcaa acgattgctg caaatgccag atgctgtgca    1320
agtgaacagt ctatcgggtc taaaccgaa gcagtagaac cacatgacat tgaagagaga    1380
atcgggaaaa tcagagtctc ttcaagaaga caatctgcaa acattgaaac tccgagagcc    1440
atcaaagaac ctgcaaatcc gcctttgcat gatgacaatg ttgaggagtc tagtcagata    1500
tcatgttcag tttcaatgga gcttaaaaga gaatcaaaga agaaccaac aggcgacgaa    1560
tcagaggaaa tgagaaaaac aactgttga agaccttcaa ggcaagctgc tgaaaaaatc    1620
aaatcgtaca agaaccttc acttaaggag aagatgagag ggggcttctg a                1671

```

```

<210> SEQ ID NO 10
<211> LENGTH: 556
<212> TYPE: PRT
<213> ORGANISM: Arabidopsis thaliana

```

```

<400> SEQUENCE: 10

Met Val Arg Ala Thr Val Leu Asn Val Gly Asp His Ala Ser Glu Gly
1          5          10          15
Val Arg Thr Asn Lys Ala Lys Gly Glu Lys Met Val Leu Glu Pro Pro
20          25          30

```

-continued

---

Met	Asn	Ser	Ala	Gln	Arg	Arg	Lys	Leu	Gly	Asp	Ile	Thr	Asn	Leu	Gln
	35						40					45			
Asn	Gln	Lys	Asn	Leu	Met	Asn	Gln	Gly	Ala	Lys	His	Gln	Gln	Gln	Ala
	50					55					60				
Ile	Leu	Ile	Ser	Ser	Lys	Glu	Asn	Ala	Glu	Asn	Leu	Gln	Lys	Ala	Leu
65					70					75					80
Arg	Asn	Ser	Ser	Glu	Asn	Thr	Lys	Leu	Met	Lys	Val	Val	Met	Glu	Arg
				85					90					95	
Asp	Gly	Ile	Lys	Ser	Asp	Leu	Lys	Lys	Leu	Arg	Ile	Glu	Phe	Gln	Lys
			100					105					110		
Val	Gln	Glu	Gln	Asn	Leu	Leu	Leu	Ala	Gln	Ala	Asn	Thr	Arg	Ile	Leu
		115					120					125			
Ala	Leu	Lys	Val	Leu	Gln	His	Glu	Leu	Gly	Cys	Lys	Asn	Gly	Leu	Val
	130					135					140				
Met	Ala	Arg	Lys	Met	Leu	Leu	Lys	Ala	Gln	Ala	Asn	Ala	Cys	Gly	Gly
145					150					155					160
Ala	Cys	Lys	Thr	Phe	Gln	Pro	Asn	Asp	Ala	Asp	His	Glu	His	Ala	Ser
				165					170					175	
Gly	Ser	Ser	Asn	Ala	Asn	Ser	Leu	Gln	Arg	Asn	Glu	Lys	Ala	Asn	Ser
			180						185					190	
Lys	Arg	Arg	Val	Ser	Gly	Arg	Lys	Asn	Pro	Ala	Asn	Ser	Glu	Val	Leu
		195					200					205			
Asp	Ile	Ile	Gly	Arg	Ser	Gly	Glu	Thr	Cys	Gln	Met	Glu	Asp	Asn	Ile
	210					215					220				
Asp	Asn	Lys	Lys	Leu	Val	Ser	Asp	Ser	Asp	Asn	Asp	Ala	Glu	Asn	His
225					230					235					240
Ile	Asn	Asp	Asn	Val	Gln	Ser	Lys	Arg	Tyr	Cys	Ala	Gly	Arg	Gln	Ser
				245					250					255	
Ser	Ser	Ser	Lys	Thr	Arg	Glu	Ala	Ser	Gln	Thr	Glu	Thr	Leu	Gln	Lys
			260					265					270		
Val	Val	Asp	Ala	Lys	Glu	Ile	Lys	Gly	Asp	Ala	Arg	Phe	Ser	Leu	Thr
		275					280					285			
Lys	His	Ser	Asp	Trp	Leu	Lys	Ser	Gln	Glu	Pro	Glu	Pro	Ser	Glu	Ser
	290					295					300				
Leu	Tyr	Glu	Ser	Arg	Phe	Pro	Leu	Arg	Arg	Arg	Ser	Ala	Arg	Leu	Lys
305					310					315					320
Ser	Gln	Glu	Pro	Glu	Pro	Ser	Glu	Ser	Phe	His	Asp	Ser	Ile	Glu	Thr
			325						330					335	
Thr	Lys	Arg	Arg	Arg	Ser	Ala	Ile	Arg	Ser	Ala	Met	Phe	Asn	Ile	Gln
			340					345					350		
Glu	Leu	Gly	Val	Ile	Gln	Asn	Leu	Asn	Gly	Leu	Pro	Asp	Asp	Gln	Glu
		355					360					365			
Ile	Ala	Ala	Lys	Ala	Arg	Cys	Ser	Ala	Arg	Glu	Gln	Ser	Thr	Gly	Ser
	370					375					380				
Lys	Pro	Glu	Ala	Val	Glu	Pro	His	Asp	Thr	Lys	Glu	Ile	Ile	Gly	Lys
385					390					395					400
Ser	Arg	Ile	Ser	Leu	Arg	Arg	Gln	Ser	Ala	Arg	Phe	Asn	Phe	Gln	Glu
			405						410					415	
Leu	Gly	Val	Thr	Glu	Asn	Leu	Asn	Gly	Pro	His	Asp	Asp	Gln	Thr	Ile
		420						425					430		
Ala	Ala	Asn	Ala	Arg	Cys	Cys	Ala	Ser	Glu	Gln	Ser	Ile	Gly	Ser	Lys



-continued

---

```

<210> SEQ ID NO 12
<211> LENGTH: 446
<212> TYPE: PRT
<213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 12

Met Asp Lys Glu Glu Thr Gln Gln Lys Glu Asn Met Leu Phe Ser Ser
 1          5          10          15

Gln Glu Tyr Ala Ala Lys Leu Gln Lys Ala Phe Pro Leu His Phe Asn
 20          25          30

Leu Glu Asn Met Thr Leu Met Lys Ala Leu Ala His Arg Asn Lys Leu
 35          40          45

Val Glu Leu Ser Gly Ile Glu Ile Gln Lys Leu Arg Ile Asn Leu Arg
 50          55          60

Ser Val Gln Glu Lys Asn Leu Gln Leu Ala Gln Ala Asn Ser Gln Met
 65          70          75          80

Leu Ala Leu Lys Asp Leu Gln His Glu Leu Gly Cys Lys Asn Ala Leu
 85          90          95

Leu Lys Val Lys Lys His Leu Glu Glu Gln Val Leu Pro Arg Thr His
 100         105         110

His Glu Ser Lys Asp Lys Val Ser Ala Ser Ala Ser Asp Gly Asp Cys
 115         120         125

Lys Ser Phe Gln Val His Asp Ile Lys His Lys Asp Thr Lys Arg Lys
 130         135         140

Arg Thr Thr Arg Ile Lys Ser Ser Val Ser Ala Asp Val Lys Pro Ile
 145         150         155         160

Pro Val Asn Asp Ser Asn Ser Lys Ala Asn Arg Lys Arg Arg Val Ser
 165         170         175

Gly Val Ile Asp Thr Thr Gly Ile Pro Glu Glu Ile Cys Gln Thr Glu
 180         185         190

Asp Asp Ile Asp Lys Gly Val Val Ser Arg Gly Val Asn Gln Asp Ile
 195         200         205

Asp Asn Val Val Asn Lys Lys Phe Val Pro Asp Ala Ala Asn Pro Val
 210         215         220

Lys Glu Ser Val His Arg Lys Arg Gln Cys Thr Arg Arg Gln Ser Thr
 225         230         235         240

Arg Phe Asp Val Gln Glu Thr Lys Gln Thr Glu Lys Leu Leu Glu Met
 245         250         255

Asp Gly Ala Lys Glu Ser Lys Glu Thr Ala Ser Phe Ser Leu Arg Arg
 260         265         270

Arg Ser Ala Arg Leu Arg His Glu Glu Ala Glu Pro Cys Lys Ser Leu
 275         280         285

His Glu Gly Asp Glu Val Arg Glu Thr Ile Lys Arg Arg Arg Val Ser
 290         295         300

Leu Arg Leu Ser Ala Arg Phe Asp Ile Gln Glu Pro His Val Thr Glu
 305         310         315         320

Thr Ser Asn Ala Asp Asp Ala Arg Ser Ile Val Ile Glu Glu Ser Ala
 325         330         335

Gly Ser Arg Ser Glu Ser Val Glu Pro Ser Glu Ser Arg His Glu Thr
 340         345         350

Lys Glu Ile Thr Arg Lys Arg Ser Phe Ser Thr Arg Arg Gln Ser Thr

```

-continued

355				360				365							
Lys	Gly	Lys	Ser	Gln	Thr	Asp	Glu	Ala	Ile	Lys	Glu	Ile	Ala	Thr	Asp
	370					375					380				
Pro	Ser	Leu	Val	Asn	Thr	Ile	Val	Gln	Glu	Cys	Asp	Gln	Glu	Thr	Glu
385					390					395					400
Ser	Lys	Asp	Lys	Pro	Lys	Ala	Asp	Glu	Asn	Glu	Gly	Met	Thr	Arg	Arg
				405					410					415	
Ser	Ser	Val	Gly	Arg	Pro	Ser	Arg	His	Ala	Ala	Glu	Lys	Val	Gln	Ser
			420					425					430		
Tyr	Arg	Glu	Val	Ser	Leu	Arg	Val	Lys	Met	Arg	Arg	Lys	Cys		
		435					440					445			

<210> SEQ ID NO 13  
 <211> LENGTH: 1554  
 <212> TYPE: DNA  
 <213> ORGANISM: mouse

<400> SEQUENCE: 13

```

atggctaagg aaaggtgtca gaaaagggtcc tttcaagata cccttgaaga cattaagaat    60
cgaatgaaag aaaaaaggaa taaaaatttg gcggggattg ggaaacgcaa gtcctttatt    120
gttgaccggg gccaaagtacc cactaacact gctacactac tgagatatta ccaagataac    180
aacaggttgt tagtcttggc tttggaaaat gagaaatcca aagtgagaga agcacaggat    240
gtcatcctgc aactgagaaa agaatgctac taccttactt gtcagctgta tgcattgaaa    300
gagaagctaa cttcccgaca aagtgaagaa actactcaga actggaaagg acgtccctca    360
gacgtggtct ccagcattga caatacgacc agggacttgt cagggaaagtc cttacagcaa    420
attgtctgtt aagaaactga ttgtccttac caaaccacag aaccaagtcc tgctgttact    480
ccagagacac aggggtgcga ttttgattca ggtaaagttg agtctactga tgaagtotta    540
cccagaacta tatctatccg tcgccattta aggaaagatt ttagtaatat aagccactcc    600
acgacttttg aggattgtaa agccagtcca agagtggcac agtctctgga agttaaagga    660
agtagatgta gagaagtaac cgtaaccctg cacagacttg aaaatgtttg tctgtggaac    720
aaagaccaaa ttagcttatg ttctagactg attaaccagc caaagattac tgaaacagaa    780
gtcattttat catctaaacc tgaacaataa gaaagcaagc ataaacgtgc acgaaaaaga    840
agagcagagc aaagaagaac caagcagaga tgcaaatcaa aatcctcatt gaggagtaag    900
gggaacaaaa acaagataa gcaggggtta ccccctacta cactggatgg aggtattggt    960
tcctgtgatg cttacgattt taatctaaaa gggacgtgcc accccacccc tttccgacaa    1020
aaaatgaaca atggctgcaa caaagaaacg gatagcagca actcagaagt gagtgacctc    1080
gaatgcagta cctctgagga tgagtctgat gacctctacc tgcctccctc caagcgcttg    1140
cgagactaca gagagtcaga gagagcagtt accaggcctc ggtctaaaag aggacttcag    1200
taccagatg ggaaagagag gaaggagtg ctgccatcta cagctcctac tggtatocca    1260
cctgagactc aagagtacc tcgttgtagc ctaaaggatg tcaccaatat cctgcagtg    1320
cctagagtga agatcaggaa gccttctctg cctccaaagc ggcgtgaaga cagcccagca    1380
gtggctctga ctaaacgcag gtgtagcacc atcaaaagct ataaagagcc aacactcgct    1440
tcaaagctaa gaagagggga ccctttcacc gacttgtgtt tcttgaattc tcctattttc    1500
aagcagaaaa ggggtatgag atgtcctaaa agaagaacca agcaaacaca gtaa    1554

```



-continued

```

<210> SEQ ID NO 14
<211> LENGTH: 517
<212> TYPE: PRT
<213> ORGANISM: mosue

<400> SEQUENCE: 14

Met Ala Lys Glu Arg Cys Gln Lys Arg Ser Phe Gln Asp Thr Leu Glu
 1           5           10           15

Asp Ile Lys Asn Arg Met Lys Glu Lys Arg Asn Lys Asn Leu Ala Gly
 20           25           30

Ile Gly Lys Arg Lys Ser Phe Ile Val Ala Pro Gly Gln Val Pro Thr
 35           40           45

Asn Thr Ala Thr Leu Leu Arg Tyr Tyr Gln Asp Asn Asn Arg Leu Leu
 50           55           60

Val Leu Ala Leu Glu Asn Glu Lys Ser Lys Val Arg Glu Ala Gln Asp
 65           70           75           80

Val Ile Leu Gln Leu Arg Lys Glu Cys Tyr Tyr Leu Thr Cys Gln Leu
 85           90           95

Tyr Ala Leu Lys Glu Lys Leu Thr Ser Arg Gln Ser Glu Glu Thr Thr
 100          105          110

Gln Asn Trp Lys Gly Arg Pro Ser Asp Val Val Ser Ser Ile Asp Asn
 115          120          125

Thr Thr Arg Asp Leu Ser Gly Lys Ser Leu Gln Gln Ile Ala Val Glu
 130          135          140

Glu Thr Asp Cys Pro Tyr Gln Thr Thr Glu Pro Ser Pro Ala Val Thr
 145          150          155          160

Pro Glu Thr Gln Gly Cys Asp Phe Asp Ser Gly Lys Val Glu Ser Thr
 165          170          175

Asp Glu Val Leu Pro Arg Thr Ile Ser Ile Arg Arg His Leu Arg Lys
 180          185          190

Asp Phe Ser Asn Ile Ser His Ser Thr Thr Leu Glu Asp Cys Lys Ala
 195          200          205

Ser Pro Arg Val Ala Gln Ser Leu Glu Val Lys Gly Ser Arg Cys Arg
 210          215          220

Glu Val Thr Val Thr Leu His Arg Leu Glu Asn Val Cys Leu Trp Asn
 225          230          235          240

Lys Asp Gln Ile Ser Leu Cys Ser Arg Leu Ile Asn Pro Ala Lys Ile
 245          250          255

Thr Glu Thr Glu Val Ile Leu Ser Ser Lys Pro Glu Gln Ile Glu Ser
 260          265          270

Lys His Lys Arg Ala Arg Lys Arg Arg Ala Glu Gln Arg Arg Thr Lys
 275          280          285

Gln Arg Cys Lys Ser Lys Ser Ser Leu Arg Ser Lys Gly Asn Lys Asn
 290          295          300

Lys Asp Lys Gln Gly Leu Pro Pro Thr Thr Leu Asp Gly Gly Ile Gly
 305          310          315          320

Ser Cys Asp Ala Tyr Asp Phe Asn Leu Lys Gly Thr Val His Pro Thr
 325          330          335

Pro Phe Arg Gln Lys Met Asn Asn Gly Cys Asn Lys Glu Thr Asp Ser
 340          345          350

Ser Asn Ser Glu Val Ser Asp Leu Glu Cys Ser Thr Ser Glu Asp Glu

```

-continued

355	360	365	
Ser Asp Asp Leu Tyr Leu	Pro Pro Ser Lys Arg	Leu Arg Asp Tyr Arg	
370	375	380	
Glu Ser Glu Arg Ala Val	Thr Arg Pro Arg Ser	Lys Arg Gly Leu Gln	
385	390	395	400
Tyr Pro Asp Gly Lys Glu	Arg Lys Glu Val Leu	Pro Ser Thr Ala Pro	
	405	410	415
Thr Gly Ile Pro Pro Glu	Thr Gln Glu Ser Pro	Arg Cys Ser Leu Lys	
	420	425	430
Asp Val Thr Asn Ile Leu	Gln Cys Pro Arg Val	Lys Ile Arg Lys Pro	
	435	440	445
Ser Leu Pro Pro Lys Arg	Arg Glu Asp Ser Pro	Ala Val Ala Leu Thr	
	450	455	460
Lys Arg Arg Cys Ser Thr	Ile Lys Ser Tyr Lys	Glu Pro Thr Leu Ala	
465	470	475	480
Ser Lys Leu Arg Arg Gly	Asp Pro Phe Thr Asp	Leu Cys Phe Leu Asn	
	485	490	495
Ser Pro Ile Phe Lys Gln	Lys Arg Gly Met Arg	Cys Pro Lys Arg Arg	
	500	505	510
Thr Lys Gln Thr Gln			
	515		

<210> SEQ ID NO 15  
 <211> LENGTH: 3495  
 <212> TYPE: DNA  
 <213> ORGANISM: mouse

<400> SEQUENCE: 15

```

atggagtacc cagggataaa agttgacact gttacctctg gaattcagag acgagtgaag    60
ggcagaattg caaagacaaa tttgaatggt tctcttgctt caaagatcaa agcaaaaata    120
ttaacaattt cttctatctt caagatctct ctaaagcaca acaacagagc attagcgcgg    180
gcccttagta aagagaaaga gaattctcga agaattacta ccgaaaagat gcaattacag    240
aaagaagtag agaaactgaa ttttgagaat acctttcttc gcttaaagtt aaataccttg    300
aataagaagc ttgtagaaat agaatcgcat gtgagcaatg atttgttaac tgcaattgaa    360
ataagcagtc tttctgagtt ccaccaaggt tcttttctcc tgtcagctac caagaaacaa    420
aggaacagta agcagtgcaa gcctgcgcat cttccatagc caagagttct gttaacttca    480
gaaaatgatg atgatgatgg tgctgatgat aatggcaga caaagtgtaa caacagaact    540
atatcaaaga cctcacctga tagtacctct tcagtatcaa gacaaccttc atccttcat    600
cagtgcaatt tgaaagcatt ccctcctaaa gaagataatc agaagacatg tgggtcaggt    660
catttagaac atacttcaag tgttgatata cttcctaagc agagccactc agatcaaagt    720
cctaagagtt ctctgagtga gatgaaaact gctccatctc ccagcctcag aagggaaaaa    780
ttatcacatg gtaatgtgac tatgaggaag aagtgtgtgt cttcaactcc agacattctg    840
tatgtgacag atttagatca ccaaccaact tcaagtccag gatcaaattg gaataatgag    900
atacatggtc atactaatga aaccagcaat aacacgcaa gaaatgccga gtgttttctt    960
gacttacctt ctgagtcttc cagtgagcct gacgcaaagc gcatggagct agtcagaag    1020
aacaccgata gctttcactt ccagaaaact gtatatgatg ccgctgatat ggagttaact    1080
    
```

-continued

---

gctactgaca taggcaagat thtagcagtt tcaaaaagca agaaaaatca aaataagaaa	1140
aaggcagact gtagaaagga gactttcaga aaagtgaaag gtgcaagctc tgataaaaag	1200
agagaaagct caaagagaga atgtaaagat ggttcagaag taggtgctga ggaagaggct	1260
gatgcagcca gagcagaaag aggcgctggt gtcctggatg gcagagggga ttcagaagag	1320
ccaaactgca tttccagtac tgagcagcca tctcaggtaa acacgcaaaa gaaaagaacc	1380
ctccagaaca gctcagatca ggagaacatt caaaatacga agaggaggca aacatatacg	1440
acagatgagc aagaggaaac aaacccttcc tccagacatt cagtcaaatt tcttcaagat	1500
ggtaaatttg atctgtgtca gaaaacccta catcataatt taagtaagcc ttctcgacag	1560
acatttgtga ttcgtaagtc agaaaaagat aacttatttc caaatcaaga agataaagac	1620
accatttctg aaaacctaga agttacaaat gaatttcata tagatgatct ttccatcgaa	1680
gctaataaaa atgtatgtga ccatgagact cagacaatgt tggacttgaa aaagtctgtc	1740
agtgtcaac aaaatcaaac aaaaataaat aagactaagc agaaaaataa tcgaaggaca	1800
aaaataatth ctgtcatgag ccaagtatat gaggacaatg ataaagatat tcacgtccta	1860
gaaaagaca actttccctt tcatacccaa gcaataaag aaaccaccag tggaaaccta	1920
gaaagttcaa aagaatttga atcacctctt cttttcaca gagacaacgg aagcttacgt	1980
gactgtaaga cccagaatgt tctggatctg cacaagcaaa ttcctgatct ataccctgat	2040
cggaatgagt cccagattag caaaatccct aggcaaaaag taaatcgcaa gacagaagta	2100
atcttctggag tgaatgttt tagtaatgac caagggttcc attgctcaga aaaggataag	2160
tctttgttac taaaaagga taaagacttc ccaggaactt taaaagactt aagtgagttt	2220
gatacgcctg ctttttghta caaagatagt gcaaaagcgt gtgattataa gtctgaaatg	2280
ctcttggggt tgaaaaaca tgacccta atgcaacctg cttgtcaaga tgattcaaaa	2340
gcaggtaaga aacttagaca aaaggtaaat cgaaaaacag aaataatthc taaaatcacc	2400
caaatacatg aaaatgatag aggaagtaca catgactcat taaataagaa gctctgtcag	2460
aaggttataa tatcaaaaat catttctcaa atgaaccaa tatatgagac tattaatgaa	2520
gatggaatg gctttaaag ctctatcaaa gattgogaag atattaaaag ttgtgacttt	2580
gggaaatca acagtaataa aaaggaaaat tatgatccaa ttcaagatcc ttgcacactg	2640
gttaaaaaaa caaagagaaa gggatcatgt aaagcagga gcagtttggc aggagctaag	2700
aacaggtgtg gtttgcagtt aacagactct tcccaggtag agtctgtccc cttagactct	2760
ggcttaagac accatccaaa cgaagcagat tctggctcctg gagagcagac taacctgcca	2820
aagatgcaga aacaaagcgc tgggaggtca ctgggagatg ctttctctgt gagtctggga	2880
aaagaaggaa gccgccagc caaagcagtt agtaaatga caccocaaatc aaagaagaga	2940
aagctccctc tcggtgttcc tcctgaaacc cacgggacgg tggagataac acccaacact	3000
gacctcgcta aggtgttga ctcccacag actgagaagg agaactatth ggagaaggag	3060
aaaattgcca agaggaagcc agatthttgt acaaaggtgt tgaaacctth atctgagaca	3120
tgttcatcta acataaagaa ttcttctctg gacagtatgt gtaagagttc gctacctttg	3180
agtatthctt ctgaaaaaac cctgatgctg gaagaaagtt cttccctgga gagtacatgc	3240
atctttcaag taggtgatgc cgctcatgag aagataacga caggcacagc taatccccac	3300
cacaggacac agaagtcgac accgggtagc agaactgccc tggctttggt ggataaccagt	3360

-continued

---

```
tctgtttcag ataccaaccc tgetaacccc gagaatgagt cagaagggca gtcttcacac 3420
ccaatgagaa ggaaaagaca gtgcgtccct ctcaacctga cagagccaag ccttagaagc 3480
aagatgagga gataa 3495
```

```
<210> SEQ ID NO 16
<211> LENGTH: 1164
<212> TYPE: PRT
<213> ORGANISM: mouse
```

```
<400> SEQUENCE: 16
```

```
Met Glu Tyr Pro Gly Ile Lys Val Asp Thr Val Thr Ser Gly Ile Gln
1 5 10 15
Arg Arg Val Lys Gly Arg Ile Ala Lys Thr Asn Leu Asn Val Ser Leu
20 25 30
Ala Ser Lys Ile Lys Ala Lys Ile Leu Asn Asn Ser Ser Ile Phe Lys
35 40 45
Ile Ser Leu Lys His Asn Asn Arg Ala Leu Ala Arg Ala Leu Ser Lys
50 55 60
Glu Lys Glu Asn Ser Arg Arg Ile Thr Thr Glu Lys Met Gln Leu Gln
65 70 75 80
Lys Glu Val Glu Lys Leu Asn Phe Glu Asn Thr Phe Leu Arg Leu Lys
85 90 95
Leu Asn Thr Leu Asn Lys Lys Leu Val Glu Ile Glu Ser His Val Ser
100 105 110
Asn Asp Leu Leu Thr Ala Ile Glu Ile Ser Ser Leu Ser Glu Phe His
115 120 125
Gln Gly Ser Phe Leu Leu Ser Ala Thr Lys Lys Gln Arg Asn Ser Lys
130 135 140
Gln Cys Lys Pro Ala His Leu Pro Tyr Ala Arg Val Leu Leu Thr Ser
145 150 155 160
Glu Asn Asp Asp Asp Asp Gly Ala Asp Asp Lys Trp Gln Thr Lys Cys
165 170 175
Asn Asn Arg Thr Ile Ser Lys Thr Ser Pro Asp Ser Thr Ser Ser Val
180 185 190
Ser Arg Gln Pro Ser Ser Leu His Gln Cys Asn Leu Lys Ala Phe Pro
195 200 205
Pro Lys Glu Asp Asn Gln Lys Thr Cys Gly Ser Gly His Leu Glu His
210 215 220
Thr Ser Ser Val Asp Ile Leu Pro Asn Glu Ser His Ser Asp Gln Ser
225 230 235 240
Pro Lys Ser Ser Leu Ser Glu Met Lys Thr Ala Pro Ser Pro Ser Leu
245 250 255
Arg Arg Glu Lys Leu Ser His Gly Asn Val Thr Met Arg Lys Lys Cys
260 265 270
Val Ser Ser Thr Pro Asp Ile Leu Tyr Val Thr Asp Leu Asp His Gln
275 280 285
Pro Thr Ser Ser Pro Gly Ser Asn Trp Asn Asn Glu Ile His Gly His
290 295 300
Thr Asn Glu Thr Ser Asn Asn Thr Gln Arg Asn Ala Glu Cys Phe Leu
305 310 315 320
Asp Leu Pro Ser Glu Ser Ser Ser Glu Pro Asp Ala Lys Arg Met Glu
325 330 335
```



-continued

---

Leu Ser Glu Phe Asp Thr Pro Ala Phe Cys Asn Lys Asp Ser Ala Lys  
                   740                                  745                                  750

Ser Cys Asp Tyr Lys Ser Glu Met Leu Leu Gly Leu Lys Lys His Asp  
                   755                                  760                                  765

Pro Asn Met Gln Pro Ala Cys Gln Asp Asp Ser Lys Ala Gly Lys Lys  
                   770                                  775                                  780

Leu Arg Gln Lys Val Asn Arg Lys Thr Glu Ile Ile Ser Lys Ile Thr  
 785                                  790                                  795                                  800

Gln Ile His Glu Asn Asp Arg Gly Ser Thr His Asp Ser Leu Asn Lys  
                                   805                                  810                                  815

Lys Leu Cys Gln Lys Val Asn Ile Ser Lys Ile Ile Ser Gln Met Asn  
                                   820                                  825                                  830

Gln Ile Tyr Glu Thr Ile Asn Glu Asp Gly Asn Gly Phe Lys Ser Ser  
                   835                                  840                                  845

Ile Lys Asp Cys Glu Asp Ile Lys Ser Cys Asp Phe Gly Glu Ile Asn  
                   850                                  855                                  860

Ser Asn Lys Lys Glu Asn Tyr Asp Pro Ile Gln Asp Pro Cys Thr Leu  
 865                                  870                                  875                                  880

Val Lys Lys Thr Lys Arg Lys Gly Ser Cys Lys Ala Gly Ser Ser Leu  
                                   885                                  890                                  895

Ala Gly Ala Lys Asn Arg Cys Gly Leu Gln Leu Thr Asp Ser Ser Gln  
                                   900                                  905                                  910

Val Gln Ser Val Pro Leu Asp Ser Gly Leu Arg His His Pro Asn Glu  
                   915                                  920                                  925

Ala Asp Ser Gly Pro Gly Glu Gln Thr Asn Leu Pro Lys Met Gln Lys  
                   930                                  935                                  940

Gln Ser Ala Gly Arg Ser Leu Gly Asp Ala Phe Ser Val Ser Leu Gly  
 945                                  950                                  955                                  960

Lys Glu Gly Ser Arg Pro Ala Lys Ala Val Ser Lys Met Thr Pro Lys  
                                   965                                  970                                  975

Ser Lys Lys Arg Lys Leu Pro Leu Gly Cys Ser Pro Glu Thr His Gly  
                                   980                                  985                                  990

Thr Val Glu Ile Thr Pro Asn Thr Asp Leu Ala Lys Ala Val Asp Ser  
                   995                                  1000                                  1005

Gln Gln Thr Glu Lys Glu Asn Tyr Leu Glu Lys Glu Lys Ile Ala  
                   1010                                  1015                                  1020

Lys Arg Lys Pro Asp Phe Cys Thr Lys Val Leu Lys Pro Leu Ser  
                   1025                                  1030                                  1035

Glu Thr Cys Ser Ser Asn Ile Lys Asn Ser Ser Leu Asp Ser Met  
                   1040                                  1045                                  1050

Cys Lys Ser Ser Leu Pro Leu Ser Ile Ser Ser Arg Lys Thr Leu  
                   1055                                  1060                                  1065

Met Leu Glu Glu Ser Ser Ser Leu Glu Ser Thr Cys Ile Phe Gln  
                   1070                                  1075                                  1080

Val Gly Asp Ala Ala His Glu Lys Ile Thr Thr Gly Thr Arg Asn  
                   1085                                  1090                                  1095

Pro His His Arg Thr Gln Lys Ser Thr Pro Gly Ser Arg Thr Ser  
                   1100                                  1105                                  1110

Leu Val Leu Val Asp Thr Ser Ser Val Ser Asp Thr Asn Pro Ala  
                   1115                                  1120                                  1125

Asn Pro Glu Asn Glu Ser Glu Gly Gln Ser Ser His Pro Met Arg

-continued

1130	1135	1140	
Arg Lys Arg Gln Cys Val Pro Leu Asn Leu Thr Glu Pro Ser Leu			
1145	1150	1155	
Arg Ser Lys Met Arg Arg			
1160			

<210> SEQ ID NO 17  
 <211> LENGTH: 1525  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 17

```

tcgcccacgc gtccgaagga ataaaaactt ggcagagatt ggcaaacgca ggtcttttat    60
agctgcacca tgccaaataa tcaccaaacac ttctacactg ctgaaaaatt accaagacaa    120
caacaaaatg ttagttag ctttgaaaa tgaaaaatcc aaagtgaag aagccaaga    180
tatcatccta cagctgagaa aagaatgta ctatctcaca tgtcagctat atgcattgaa    240
aggaaaactt acatcacac aaacagtaga acctgctcag aaccaggaaa tatgttctc    300
tggaatgac ccaatagtg atgacagctc cagaaatta tttgtgaag atttaccgca    360
aattcctctt gaagaaactg aacttccagg acaaggagaa tcatttcaa tagaagatca    420
gatacctact attcctcaag acacactggg agttgatttt gattcaggtg aagctaagtc    480
tactgataat gtcttaccta gaactgtatc tgttcgtagc agtttaaga aacattgtaa    540
cagtatatgt cagtttgata gcttgatga ttttgaaacc agtcatttgg cagggaagtc    600
ttttgaattc gaaagagttg gattttttaga cccactagta aacatgcaca tacctgaaaa    660
tgtacaacac aatgcttgtc aatggagcaa ggaccaagtt aacttatcac caaagctgat    720
tcagccagga acgtttacta aaacaaaaga agacatttta gaatctaaat ctgaacaaac    780
taaaagtaag caaagagata cacaagaaag aaaaagagaa gagaaaagaa aagctaacag    840
gagaaaatca aaacgtatgt caaaatataa agagaataaa agcgaaaata aaaaaactgt    900
tccccaaaa aaaatgcaca aatctgtcag ttccaatgat gcttacaatt ttaatttga    960
agagggtgtt catcttactc ctttccgaca aaaagtgagc aatgactcta atagagaaga    1020
aaacaacgag tctgaagtga gcctctgtga atcaagtggc tcaggagatg attccgatga    1080
cctctatttg cccacttgca agtacattca gaatcccacg agcaattcag atagaccagt    1140
caccaggcct ctagtataaa gagcaactgaa atacacagat gaaaaagaga cggaggggtc    1200
taagccaaca aaaactccta ccactacacc acctgaaact cagcagtcac ctcatcttag    1260
cctgaaggat atcaacaatg tctccttgta tctgtttgtg aaaatcagaa gactttctct    1320
ttctccaaaa aagaataaag caagcccagc agtggtctctg cctaaacgta ggtgcacagc    1380
cagcgtgaac tataaggagc ccaccctcgc ttcgaaactg agaagagggg acccttttac    1440
agatttgtgt tttttgaatt ctctatctt caagcagaaa aaggatttga gacgttctaa    1500
aaaaagtatg aaacaaatac aatga                                     1525
  
```

<210> SEQ ID NO 18  
 <211> LENGTH: 511  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 18

-continued

---

Gly Arg Val Gly Arg Pro Arg Val Arg Arg Asn Lys Asn Leu Ala Glu  
 1 5 10 15  
 Ile Gly Lys Arg Arg Ser Phe Ile Ala Ala Pro Cys Gln Ile Ile Thr  
 20 25 30  
 Asn Thr Ser Thr Leu Leu Lys Asn Tyr Gln Asp Asn Asn Lys Met Leu  
 35 40 45  
 Val Leu Ala Leu Glu Asn Glu Lys Ser Lys Val Lys Glu Ala Gln Asp  
 50 55 60  
 Ile Ile Leu Gln Leu Arg Lys Glu Cys Tyr Tyr Leu Thr Cys Gln Leu  
 65 70 75 80  
 Tyr Ala Leu Lys Gly Lys Leu Thr Ser Gln Gln Thr Val Glu Pro Ala  
 85 90 95  
 Gln Asn Gln Glu Ile Cys Ser Ser Gly Met Asp Pro Asn Ser Asp Asp  
 100 105 110  
 Ser Ser Arg Asn Leu Phe Val Lys Asp Leu Pro Gln Ile Pro Leu Glu  
 115 120 125  
 Glu Thr Glu Leu Pro Gly Gln Gly Glu Ser Phe Gln Ile Glu Asp Gln  
 130 135 140  
 Ile Pro Thr Ile Pro Gln Asp Thr Leu Gly Val Asp Phe Asp Ser Gly  
 145 150 155 160  
 Glu Ala Lys Ser Thr Asp Asn Val Leu Pro Arg Thr Val Ser Val Arg  
 165 170 175  
 Ser Ser Leu Lys Lys His Cys Asn Ser Ile Cys Gln Phe Asp Ser Leu  
 180 185 190  
 Asp Asp Phe Glu Thr Ser His Leu Ala Gly Lys Ser Phe Glu Phe Glu  
 195 200 205  
 Arg Val Gly Phe Leu Asp Pro Leu Val Asn Met His Ile Pro Glu Asn  
 210 215 220  
 Val Gln His Asn Ala Cys Gln Trp Ser Lys Asp Gln Val Asn Leu Ser  
 225 230 235 240  
 Pro Lys Leu Ile Gln Pro Gly Thr Phe Thr Lys Thr Lys Glu Asp Ile  
 245 250 255  
 Leu Glu Ser Lys Ser Glu Gln Thr Lys Ser Lys Gln Arg Asp Thr Gln  
 260 265 270  
 Glu Arg Lys Arg Glu Glu Lys Arg Lys Ala Asn Arg Arg Lys Ser Lys  
 275 280 285  
 Arg Met Ser Lys Tyr Lys Glu Asn Lys Ser Glu Asn Lys Lys Thr Val  
 290 295 300  
 Pro Gln Lys Lys Met His Lys Ser Val Ser Ser Asn Asp Ala Tyr Asn  
 305 310 315 320  
 Phe Asn Leu Glu Glu Gly Val His Leu Thr Pro Phe Arg Gln Lys Val  
 325 330 335  
 Ser Asn Asp Ser Asn Arg Glu Glu Asn Asn Glu Ser Glu Val Ser Leu  
 340 345 350  
 Cys Glu Ser Ser Gly Ser Gly Asp Asp Ser Asp Asp Leu Tyr Leu Pro  
 355 360 365  
 Thr Cys Lys Tyr Ile Gln Asn Pro Thr Ser Asn Ser Asp Arg Pro Val  
 370 375 380  
 Thr Arg Pro Leu Ala Lys Arg Ala Leu Lys Tyr Thr Asp Glu Lys Glu  
 385 390 395 400  
 Thr Glu Gly Ser Lys Pro Thr Lys Thr Pro Thr Thr Pro Pro Glu



-continued

	405		410		415										
Thr	Gln	Gln	Ser	Pro	His	Leu	Ser	Leu	Lys	Asp	Ile	Thr	Asn	Val	Ser
	420							425					430		
Leu	Tyr	Pro	Val	Val	Lys	Ile	Arg	Arg	Leu	Ser	Leu	Ser	Pro	Lys	Lys
	435						440					445			
Asn	Lys	Ala	Ser	Pro	Ala	Val	Ala	Leu	Pro	Lys	Arg	Arg	Cys	Thr	Ala
	450					455					460				
Ser	Val	Asn	Tyr	Lys	Glu	Pro	Thr	Leu	Ala	Ser	Lys	Leu	Arg	Arg	Gly
465					470					475					480
Asp	Pro	Phe	Thr	Asp	Leu	Cys	Phe	Leu	Asn	Ser	Pro	Ile	Phe	Lys	Gln
			485						490					495	
Lys	Lys	Asp	Leu	Arg	Arg	Ser	Lys	Lys	Ser	Met	Lys	Gln	Ile	Gln	
		500						505					510		

<210> SEQ ID NO 19  
 <211> LENGTH: 3798  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 19

```

atggagtgcc cagtgatgga aactggctca ctttttacct caggaattaa gagacatttg      60
aaagacaaaa gaatttcaaa gactactaag ttgaatgttt ctcttgcttc aaaaataaaa      120
acaaaaatac taaataattc ttctatcttc aaaatatctt taaagcacia caacagggca      180
ttagctcagg ctcttagtag agaaaaagag aattctcgaa gaattacaac tgaaaagatg      240
ctattgcaaa aagaagtaga gaaactgaat ttgagaaca catttcttcg cctaaagcta      300
aataacttga ataagaagct tatagacata gaagctctca tgaacaataa cttgataact      360
gcaactgaaa tgagcagtct tcttgagttc catcagagtt cctttctact gtcagctagc      420
aagaagaaac gagttagtaa acagtgcaag ttgatgcgtc ttccatttgc aagggttcca      480
ttaacttcaa atgatgatga agatgaagat aaagagaaaa tgcagtgatga caacaatatt      540
aatcaaaga cattacctga tattccctct tcaggatcaa caacacaacc tttatcaact      600
caggataatt cggaagtgtt atttcttaaa gaaaataatc aaaatgtata tggtttagat      660
gattcagaac atatttcttc tatagttgat gtacctcca gagaaagcca ttcccactca      720
gaccaaagt ctaagacttc tctaattgag gagatgagaa acgcccagtc tattggccgc      780
agatgggaga aaccatctcc tagtaatgtg actgaaagga agaagcgttg gtcactttgg      840
gaatcaaata atctttctgc agacactccc tgtgcaacag ttttagataa acaacacatt      900
tcaagtccag aattaaattg caataatgag ataaatggtc atactaatga acaaaatact      960
gaaatgcaaa gaaataaaca ggatcttcct ggcttatctt ctgagtctgc cagagaacct      1020
aatgcagagt gcatgaatca aattgaggat aatgatgact ttcaattgca gaaaactgtg      1080
tatgatgctg acatggattt aactgctagt gaagtcagca aaattgtcac agtctcaaca      1140
ggcattaaaa agaaaagtaa taaaaaaca aatgaacatg gaatgaaac tttcagaaaa      1200
gtgaaagatt ccagctctga aaaaaagaga gaaagatcaa agagacagtt taaaaatagt      1260
tcagatgtcg atattgggga aaagattgaa aacaggacag aaagatctga tgtcctggat      1320
ggcaaaaggg gtgcagaaga tcccggtttt attttcaata atgaacagct ggctcagatg      1380
aatgaacagc tggctcaggt gaatgaacta aagaaaatga cccttcaaac tggctttgaa      1440
    
```

-continued

---

caaggtgaca gagaaatgt actgtgtaat aaaaaggaga aaagaataac aaatgagcaa	1500
gaggaaacat actctttatc ccaaagtcca ggtaaatttc accaggagag taaatttgat	1560
aagggtcaga attccctaac ttgtaataaa agtaaagcct cttagacagac atttgtgatt	1620
cacaaattag aaaaagataa cttactccca aacccaaaag ataaagtaac catttatgaa	1680
aacctagacg tcacaaatga atttcacaca gccaatcttt ccaccaaaaga taatggaaat	1740
ttatgtgatt atgggaccca caatatattg gatttgaana agtatgtcac tgatattcaa	1800
ccctcagagc aaaatgaatc aaacattaat aagcttagaa agaaagtaaa ccggaagaca	1860
gaaataatth ctggaatgaa ccacatgtat gaagataatg ataaagatgt ggtgcatggc	1920
ctaaaaaaag gtaatthttt tttcaaaacc caagaggata aagaacctat ctctgaaaac	1980
atagaagtht ccaaagagct tcaaatccca gctctttcta cttagagataa tgaaaatcaa	2040
tgtgactata ggacccagaa tgtgttggtt ttgcaaaagc agatcaccaa tatgtacccc	2100
gttcagcaaa atgaatcaaa agttaaataag aagcttaggc agaaagtaaa tcggaagaca	2160
gaaataatth ctgaagttaa tcatttagat aatgacaaaa gtatagaata cacagthtaa	2220
agtcaactac tctthttaac gcaaaaagat aaggaataa tccccgaaa cctagaagac	2280
ccaagtgagt ttgaaacacc tgctctttct accaaagata gtggaaacct gtatgattct	2340
gagattcaaa atgtthttgg ggtgaaacat ggccatgata tgcaacctgc ttgtcaaat	2400
gattcaaaaa taggtaagaa gcctagacta aatgtatgtc aaaagtcaga aataatctct	2460
gaaaccaacc aaatatatga gaatgataac aaaggtgtac atgacctaga aaaagataac	2520
ttctctctc taaccccaaa ggataaagaa acaatthctg aaaatctaca agtcacaaat	2580
gaatttcaaa cagttgatct tctcatcaaa gataatggaa atthtatgta ttatgacacc	2640
cagaatata tggagttgaa aaagtatgtt actgatagga aatctgctga gcaaatgaa	2700
tcaaaaaata ataaagctcag gaataaagtg aattggaaga cagaataat thctgaaatg	2760
aaccagatat atgaggataa tgataaagat gcacatgtcc aagaaagcta tacaaaagat	2820
cttgatttha aagtaataa atcaaaaca aaactgaaat gccaaagcat tatcaataaa	2880
cactatatgg aagtaacag taatgaaaag gaaagttgtg atcaaattht agattctac	2940
aaagtagtta aaaaacgtaa gaaagaatca tcatgcaagg caaagaacat tttgacaaa	3000
gctaagaaca aactgtctc acagttaaca gaatcttcac agacatctat ctcttagaa	3060
tctgatttha aacatattac tagtgaagca gattctgatc caggaaacct agttgaaacta	3120
tgtaagactc agaagcaaaag cactaccact ttgaataaaa aagatctccc tttgtggaa	3180
gaaataaaag aaggagagtg tcaggthtaa aagthaaata aaatgacatc taagtcaaa	3240
aaaaggaaag cctccataga tcctctcca gagagccatg aagthaatgga aagaactth	3300
gacagcgttc agggaaagtc tactgtatct gaacaagctg ataaagaaaa caatthggag	3360
aatgagaaaa tggtaaaaa taagccagac thttacaca aggcatthtag atctthgtct	3420
gagatacatt cacctaacat acaagattct tcctthgaca gtgtctgtga aggtthtagta	3480
ctthtgagc tthctctg taaaaatgtg ataaataaaag aaaatthtgc ctthgagtg	3540
tccccagcct thcaagthaa tgatgatgag catgagaaga tgaacaagat gaaatthaaa	3600
gtcaaccgga gaacccaaaa atcaggaata ggtgatagac cattacagga ctthtcaaat	3660
accagthttg thcaaaaaa cactgctgaa tctgaaaaata agtcagaaga tctatcttca	3720

-continued

---

gaacggacaa gcagaagaag aaggtgtact cctttctatt ttaaagagcc aagcctcaga 3780

gacaagatga gaagatga 3798

&lt;210&gt; SEQ ID NO 20

&lt;211&gt; LENGTH: 1265

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 20

```

Met Glu Cys Pro Val Met Glu Thr Gly Ser Leu Phe Thr Ser Gly Ile
 1          5          10          15
Lys Arg His Leu Lys Asp Lys Arg Ile Ser Lys Thr Thr Lys Leu Asn
 20          25          30
Val Ser Leu Ala Ser Lys Ile Lys Thr Lys Ile Leu Asn Asn Ser Ser
 35          40          45
Ile Phe Lys Ile Ser Leu Lys His Asn Asn Arg Ala Leu Ala Gln Ala
 50          55          60
Leu Ser Arg Glu Lys Glu Asn Ser Arg Arg Ile Thr Thr Glu Lys Met
 65          70          75          80
Leu Leu Gln Lys Glu Val Glu Lys Leu Asn Phe Glu Asn Thr Phe Leu
 85          90          95
Arg Leu Lys Leu Asn Asn Leu Asn Lys Lys Leu Ile Asp Ile Glu Ala
 100         105         110
Leu Met Asn Asn Asn Leu Ile Thr Ala Ile Glu Met Ser Ser Leu Ser
 115         120         125
Glu Phe His Gln Ser Ser Phe Leu Leu Ser Ala Ser Lys Lys Lys Arg
 130         135         140
Ile Ser Lys Gln Cys Lys Leu Met Arg Leu Pro Phe Ala Arg Val Pro
 145         150         155         160
Leu Thr Ser Asn Asp Asp Glu Asp Glu Asp Lys Glu Lys Met Gln Cys
 165         170         175
Asp Asn Asn Ile Lys Ser Lys Thr Leu Pro Asp Ile Pro Ser Ser Gly
 180         185         190
Arg Thr Thr Gln Pro Leu Ser Thr Gln Asp Asn Ser Gly Val Leu Phe
 195         200         205
Leu Lys Glu Asn Asn Gln His Val Tyr Gly Leu Asp Asp Ser Glu His
 210         215         220
Ile Ser Ser Ile Val Asp Val Pro Pro Arg Glu Ser His Ser His Ser
 225         230         235         240
Asp Gln Ser Ser Lys Thr Ser Leu Met Ser Glu Met Arg Asn Ala Gln
 245         250         255
Ser Ile Gly Arg Arg Trp Glu Lys Pro Ser Pro Ser Asn Val Thr Glu
 260         265         270
Arg Lys Lys Arg Gly Ser Ser Trp Glu Ser Asn Asn Leu Ser Ala Asp
 275         280         285
Thr Pro Cys Ala Thr Val Leu Asp Lys Gln His Ile Ser Ser Pro Glu
 290         295         300
Leu Asn Cys Asn Asn Glu Ile Asn Gly His Thr Asn Glu Thr Asn Thr
 305         310         315         320
Glu Met Gln Arg Asn Lys Gln Asp Leu Pro Gly Leu Ser Ser Glu Ser
 325         330         335
Ala Arg Glu Pro Asn Ala Glu Cys Met Asn Gln Ile Glu Asp Asn Asp

```

-continued

340					345					350					
Asp	Phe	Gln	Leu	Gln	Lys	Thr	Val	Tyr	Asp	Ala	Asp	Met	Asp	Leu	Thr
		355							360				365		
Ala	Ser	Glu	Val	Ser	Lys	Ile	Val	Thr	Val	Ser	Thr	Gly	Ile	Lys	Lys
		370				375						380			
Lys	Ser	Asn	Lys	Lys	Thr	Asn	Glu	His	Gly	Met	Lys	Thr	Phe	Arg	Lys
		385			390					395					400
Val	Lys	Asp	Ser	Ser	Ser	Glu	Lys	Lys	Arg	Glu	Arg	Ser	Lys	Arg	Gln
				405					410					415	
Phe	Lys	Asn	Ser	Ser	Asp	Val	Asp	Ile	Gly	Glu	Lys	Ile	Glu	Asn	Arg
				420				425						430	
Thr	Glu	Arg	Ser	Asp	Val	Leu	Asp	Gly	Lys	Arg	Gly	Ala	Glu	Asp	Pro
		435					440					445			
Gly	Leu	Phe	Phe	Asn	Asn	Glu	Gln	Leu	Ala	Gln	Met	Asn	Glu	Gln	Leu
		450				455					460				
Ala	Gln	Val	Asn	Glu	Leu	Lys	Lys	Met	Thr	Leu	Gln	Thr	Gly	Phe	Glu
		465			470					475					480
Gln	Gly	Asp	Arg	Glu	Asn	Val	Leu	Cys	Asn	Lys	Lys	Glu	Lys	Arg	Val
				485					490					495	
Thr	Asn	Glu	Gln	Glu	Glu	Thr	Tyr	Ser	Leu	Ser	Gln	Ser	Ser	Gly	Lys
			500					505					510		
Phe	His	Gln	Glu	Ser	Lys	Phe	Asp	Lys	Gly	Gln	Asn	Ser	Leu	Thr	Cys
		515					520					525			
Asn	Lys	Ser	Lys	Ala	Ser	Arg	Gln	Thr	Phe	Val	Ile	His	Lys	Leu	Glu
		530				535					540				
Lys	Asp	Asn	Leu	Leu	Pro	Asn	Gln	Lys	Asp	Lys	Val	Thr	Ile	Tyr	Glu
		545			550					555					560
Asn	Leu	Asp	Val	Thr	Asn	Glu	Phe	His	Thr	Ala	Asn	Leu	Ser	Thr	Lys
				565					570					575	
Asp	Asn	Gly	Asn	Leu	Cys	Asp	Tyr	Gly	Thr	His	Asn	Ile	Leu	Asp	Leu
			580					585					590		
Lys	Lys	Tyr	Val	Thr	Asp	Ile	Gln	Pro	Ser	Glu	Gln	Asn	Glu	Ser	Asn
		595					600					605			
Ile	Asn	Lys	Leu	Arg	Lys	Lys	Val	Asn	Arg	Lys	Thr	Glu	Ile	Ile	Ser
		610				615						620			
Gly	Met	Asn	His	Met	Tyr	Glu	Asp	Asn	Asp	Lys	Asp	Val	Val	His	Gly
		625			630					635					640
Leu	Lys	Lys	Gly	Asn	Phe	Phe	Phe	Lys	Thr	Gln	Glu	Asp	Lys	Glu	Pro
				645					650					655	
Ile	Ser	Glu	Ser	Ile	Glu	Val	Ser	Lys	Glu	Leu	Gln	Ile	Pro	Ala	Leu
			660					665					670		
Ser	Thr	Arg	Asp	Asn	Glu	Asn	Gln	Cys	Asp	Tyr	Arg	Thr	Gln	Asn	Val
		675					680						685		
Leu	Gly	Leu	Gln	Lys	Gln	Ile	Thr	Asn	Met	Tyr	Pro	Val	Gln	Gln	Asn
		690				695					700				
Glu	Ser	Lys	Val	Asn	Lys	Lys	Leu	Arg	Gln	Lys	Val	Asn	Arg	Lys	Thr
				710						715					720
Glu	Ile	Ile	Ser	Glu	Val	Asn	His	Leu	Asp	Asn	Asp	Lys	Ser	Ile	Glu
				725					730					735	
Tyr	Thr	Val	Lys	Ser	His	Ser	Leu	Phe	Leu	Thr	Gln	Lys	Asp	Lys	Glu
			740					745						750	

-continued

---

Ile Ile Pro Gly Asn Leu Glu Asp Pro Ser Glu Phe Glu Thr Pro Ala  
 755 760 765

Leu Ser Thr Lys Asp Ser Gly Asn Leu Tyr Asp Ser Glu Ile Gln Asn  
 770 775 780

Val Leu Gly Val Lys His Gly His Asp Met Gln Pro Ala Cys Gln Asn  
 785 790 795 800

Asp Ser Lys Ile Gly Lys Lys Pro Arg Leu Asn Val Cys Gln Lys Ser  
 805 810 815

Glu Ile Ile Pro Glu Thr Asn Gln Ile Tyr Glu Asn Asp Asn Lys Gly  
 820 825 830

Val His Asp Leu Glu Lys Asp Asn Phe Phe Ser Leu Thr Pro Lys Asp  
 835 840 845

Lys Glu Thr Ile Ser Glu Asn Leu Gln Val Thr Asn Glu Phe Gln Thr  
 850 855 860

Val Asp Leu Leu Ile Lys Asp Asn Gly Asn Leu Cys Asp Tyr Asp Thr  
 865 870 875 880

Gln Asn Ile Leu Glu Leu Lys Lys Tyr Val Thr Asp Arg Lys Ser Ala  
 885 890 895

Glu Gln Asn Glu Ser Lys Ile Asn Lys Leu Arg Asn Lys Val Asn Trp  
 900 905 910

Lys Thr Glu Ile Ile Ser Glu Met Asn Gln Ile Tyr Glu Asp Asn Asp  
 915 920 925

Lys Asp Ala His Val Gln Glu Ser Tyr Thr Lys Asp Leu Asp Phe Lys  
 930 935 940

Val Asn Lys Ser Lys Gln Lys Leu Glu Cys Gln Asp Ile Ile Asn Lys  
 945 950 955 960

His Tyr Met Glu Val Asn Ser Asn Glu Lys Glu Ser Cys Asp Gln Ile  
 965 970 975

Leu Asp Ser Tyr Lys Val Val Lys Lys Arg Lys Lys Glu Ser Ser Cys  
 980 985 990

Lys Ala Lys Asn Ile Leu Thr Lys Ala Lys Asn Lys Leu Ala Ser Gln  
 995 1000 1005

Leu Thr Glu Ser Ser Gln Thr Ser Ile Ser Leu Glu Ser Asp Leu  
 1010 1015 1020

Lys His Ile Thr Ser Glu Ala Asp Ser Asp Pro Gly Asn Pro Val  
 1025 1030 1035

Glu Leu Cys Lys Thr Gln Lys Gln Ser Thr Thr Thr Leu Asn Lys  
 1040 1045 1050

Lys Asp Leu Pro Phe Val Glu Glu Ile Lys Glu Gly Glu Cys Gln  
 1055 1060 1065

Val Lys Lys Val Asn Lys Met Thr Ser Lys Ser Lys Lys Arg Lys  
 1070 1075 1080

Thr Ser Ile Asp Pro Ser Pro Glu Ser His Glu Val Met Glu Arg  
 1085 1090 1095

Ile Leu Asp Ser Val Gln Gly Lys Ser Thr Val Ser Glu Gln Ala  
 1100 1105 1110

Asp Lys Glu Asn Asn Leu Glu Asn Glu Lys Met Val Lys Asn Lys  
 1115 1120 1125

Pro Asp Phe Tyr Thr Lys Ala Phe Arg Ser Leu Ser Glu Ile His  
 1130 1135 1140

-continued

---

Ser Pro Asn Ile Gln Asp Ser Ser Phe Asp Ser Val Arg Glu Gly  
 1145 1150 1155

Leu Val Pro Leu Ser Val Ser Ser Gly Lys Asn Val Ile Ile Lys  
 1160 1165 1170

Glu Asn Phe Ala Leu Glu Cys Ser Pro Ala Phe Gln Val Ser Asp  
 1175 1180 1185

Asp Glu His Glu Lys Met Asn Lys Met Lys Phe Lys Val Asn Arg  
 1190 1195 1200

Arg Thr Gln Lys Ser Gly Ile Gly Asp Arg Pro Leu Gln Asp Leu  
 1205 1210 1215

Ser Asn Thr Ser Phe Val Ser Asn Asn Thr Ala Glu Ser Glu Asn  
 1220 1225 1230

Lys Ser Glu Asp Leu Ser Ser Glu Arg Thr Ser Arg Arg Arg Arg  
 1235 1240 1245

Cys Thr Pro Phe Tyr Phe Lys Glu Pro Ser Leu Arg Asp Lys Met  
 1250 1255 1260

Arg Arg  
 1265

<210> SEQ ID NO 21  
 <211> LENGTH: 45  
 <212> TYPE: PRT  
 <213> ORGANISM: yeast

<400> SEQUENCE: 21

Met Glu Ser Leu Lys Lys Lys Phe Leu Lys Gln Asn Arg Glu Ile Ile  
 1 5 10 15

Lys Ile Asn Thr Gln Leu Ser Ile Lys Ile Arg Glu Ser Glu Asn Glu  
 20 25 30

Ile Gln Asp Leu Ile Gln Glu Asn Phe Thr Leu Lys Ser  
 35 40 45

<210> SEQ ID NO 22  
 <211> LENGTH: 45  
 <212> TYPE: PRT  
 <213> ORGANISM: yeast

<400> SEQUENCE: 22

Val Glu Asp Leu Lys Lys Lys Gln Ile Arg Gln Tyr Lys Glu Ile Ile  
 1 5 10 15

Arg Ile Ser Lys Ala Gln Ser Ile Arg Ile Lys Glu Leu Gln Leu Glu  
 20 25 30

Asn Glu Arg Leu Leu Ser Glu Asn Ile Asp Leu Arg Thr  
 35 40 45

<210> SEQ ID NO 23  
 <211> LENGTH: 45  
 <212> TYPE: PRT  
 <213> ORGANISM: yeast

<400> SEQUENCE: 23

Val Glu Asn Ile Arg Gln Ser Tyr Ser Arg Gln Asn Ser Leu Leu Ala  
 1 5 10 15

Lys Asp Asn Ser Ile Leu Lys Ile Lys Val Asn Ser Leu Glu Lys Lys  
 20 25 30

Ile Ser Gln Leu Val Gln Glu Asn Val Thr Leu Arg Ser



-continued

---

20 25

<210> SEQ ID NO 29  
 <211> LENGTH: 28  
 <212> TYPE: PRT  
 <213> ORGANISM: Neurospora crassa

<400> SEQUENCE: 29

Glu Thr Ser Arg Pro Ser Arg Arg Ala Arg Ala Ala Ile Ser Tyr Thr  
 1 5 10 15  
 Glu Pro Asn Leu Arg Asp Lys Met Arg Arg Pro Thr  
 20 25

<210> SEQ ID NO 30  
 <211> LENGTH: 27  
 <212> TYPE: PRT  
 <213> ORGANISM: Dactylicapnos macrocapnos

<400> SEQUENCE: 30

Asn Ser Ala Arg Pro Ser Arg Ser Cys Arg Pro Thr Ser Leu Val Glu  
 1 5 10 15  
 Pro Ser Leu Lys Asn Lys Leu Arg Asn Gly Ser  
 20 25

<210> SEQ ID NO 31  
 <211> LENGTH: 28  
 <212> TYPE: PRT  
 <213> ORGANISM: Caenorhabditis elegans

<400> SEQUENCE: 31

Thr Val Arg Arg Gln Arg Ser Ala Lys Met Asn Ile Lys Ser Leu Lys  
 1 5 10 15  
 Glu Pro Ser Gly Lys Asp Lys Leu Arg Arg Pro Gly  
 20 25

<210> SEQ ID NO 32  
 <211> LENGTH: 29  
 <212> TYPE: PRT  
 <213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 32

Thr Val Gly Arg Pro Ser Arg Gln Ala Ala Glu Lys Ile Lys Ser Tyr  
 1 5 10 15  
 Lys Glu Pro Ser Leu Lys Glu Lys Met Arg Gly Gly Phe  
 20 25

<210> SEQ ID NO 33  
 <211> LENGTH: 29  
 <212> TYPE: PRT  
 <213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 33

Ser Val Gly Arg Pro Ser Arg His Ala Ala Glu Lys Val Gln Ser Tyr  
 1 5 10 15  
 Arg Glu Val Ser Leu Arg Val Lys Met Arg Arg Lys Cys  
 20 25

<210> SEQ ID NO 34  
 <211> LENGTH: 28  
 <212> TYPE: PRT



-continued

&lt;213&gt; ORGANISM: mouse

&lt;400&gt; SEQUENCE: 34

Ala Val Ala Leu Thr Lys Arg Arg Cys Ser Thr Ile Lys Ser Tyr Lys  
 1 5 10 15

Glu Pro Thr Leu Ala Ser Lys Leu Arg Arg Gly Asp  
 20 25

&lt;210&gt; SEQ ID NO 35

&lt;211&gt; LENGTH: 25

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: mouse

&lt;400&gt; SEQUENCE: 35

His Pro Met Arg Arg Lys Arg Gln Cys Val Pro Leu Asn Leu Thr Glu  
 1 5 10 15

Pro Ser Leu Arg Ser Lys Met Arg Arg  
 20 25

&lt;210&gt; SEQ ID NO 36

&lt;211&gt; LENGTH: 28

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 36

Ala Val Ala Leu Pro Lys Arg Arg Cys Thr Ala Ser Val Asn Tyr Lys  
 1 5 10 15

Glu Pro Thr Leu Ala Ser Lys Leu Arg Arg Gly Asp  
 20 25

&lt;210&gt; SEQ ID NO 37

&lt;211&gt; LENGTH: 26

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 37

Ser Glu Arg Thr Ser Arg Arg Arg Arg Cys Thr Pro Phe Tyr Phe Lys  
 1 5 10 15

Glu Pro Ser Leu Arg Asp Lys Met Arg Arg  
 20 25

&lt;210&gt; SEQ ID NO 38

&lt;211&gt; LENGTH: 21

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: ?TriplEx

&lt;400&gt; SEQUENCE: 38

ctcgggaagc ggcattgt g

21

&lt;210&gt; SEQ ID NO 39

&lt;211&gt; LENGTH: 22

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 39

cctggctgaa tcagtttg tg

22

&lt;210&gt; SEQ ID NO 40

---

-continued

---

<211> LENGTH: 23  
<212> TYPE: DNA  
<213> ORGANISM: Artificial  
<220> FEATURE:  
<223> OTHER INFORMATION: hSgo1  
  
<400> SEQUENCE: 40  
  
aagucuacug auaaugucuu att 23

<210> SEQ ID NO 41  
<211> LENGTH: 23  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: hSgo2  
  
<400> SEQUENCE: 41  
  
aagcacuacc acuuugaaua att 23

<210> SEQ ID NO 42  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: hSgo1  
  
<400> SEQUENCE: 42  
  
gugagccucu gugaaucaat t 21

<210> SEQ ID NO 43  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: hSgo2  
  
<400> SEQUENCE: 43  
  
gcucucauga acaauaacut t 21

<210> SEQ ID NO 44  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: siRNA,Target1  
  
<400> SEQUENCE: 44  
  
gagugaucac gauuucuaat t 21

<210> SEQ ID NO 45  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: siRNA,Target2  
  
<400> SEQUENCE: 45  
  
aacgggcauu ugaauaugaa a 21

---

- 1. A DNA encoding a following protein (a) or (b):
  - (a) a protein consisting of an amino acid sequence shown in SEQ ID NO: 2,
  - (b) a protein comprising an amino acid sequence where one or several amino acids are deleted, replaced or added in an amino acid sequence shown in SEQ ID NO: 2, and having a regulatory activity of chromosome segregation.
- 2. A DNA consisting of a base sequence shown in SEQ ID NO: 1 or a complementary sequence thereof.
- 3. A DNA containing part or whole of a base sequence shown in SEQ ID NO: 1 or a complementary sequence thereof, and encoding a protein that has a regulatory activity of chromosome segregation.
- 4. A DNA hybridizing with the DNA according to claim 2 under stringent conditions and encoding a protein that has a regulatory activity of chromosome segregation.
- 5. A protein consisting of an amino acid sequence shown in SEQ ID NO: 2.
- 6. A protein consisting of an amino acid sequence where one or several amino acids are deleted, replaced or added in an amino acid sequence shown in SEQ ID NO: 2, and having a regulatory activity of chromosome segregation.
- 7. A DNA encoding a following protein (a) or (b):
  - (a) a protein consisting of an amino acid sequence shown in SEQ ID NO: 4,
  - (b) a protein consisting of an amino acid sequence where one or several amino acids are deleted, replaced or added in an amino acid sequence shown in SEQ ID NO: 4, and having a regulatory activity of chromosome segregation.
- 8. A DNA consisting of a base sequence shown in SEQ ID NO: 3 or a complementary sequence thereof.
- 9. A DNA containing part or whole of a base sequence shown in SEQ ID NO: 3 or a complementary sequence thereof, and encoding a protein that has a regulatory activity of chromosome segregation.
- 10. A DNA hybridizing with the DNA according to claim 8 under stringent conditions and encoding a protein that has a regulatory activity of chromosome segregation.
- 11. A protein consisting of an amino acid sequence shown in SEQ ID NO: 4.
- 12. A protein consisting of an amino acid sequence where one or several amino acids are deleted, replaced or added in an amino acid sequence shown in SEQ ID NO: 4, and having a regulatory activity of chromosome segregation.
- 13. A DNA encoding a following protein (a) or (b):
  - (a) a protein consisting of an amino acid sequence shown in SEQ ID NO: 6,
  - (b) a protein consisting of an amino acid sequence where one or several amino acids are deleted, replaced or added in an amino acid sequence shown in SEQ ID NO: 6, and having a regulatory activity of chromosome segregation.

- 14. A DNA consisting of a base sequence shown in SEQ ID NO: 5 or a complementary sequence thereof.
- 15. A DNA containing part or whole of a base sequence shown in SEQ ID NO: 5 or a complementary sequence thereof, and encoding a protein that has a regulatory activity of chromosome segregation.
- 16. A DNA hybridizing with the DNA according to claim 14 under stringent conditions and encoding a protein that has a regulatory activity of chromosome segregation.
- 17. A protein consisting of an amino acid sequence shown in SEQ ID NO: 6.
- 18. A protein consisting of an amino acid sequence where one or several amino acids are deleted, replaced or added in an amino acid sequence shown in SEQ ID NO: 6, and having a regulatory activity of chromosome segregation.
- 19. A DNA encoding a following protein (a) or (b) that has a regulatory activity of chromosome segregation:
  - (a) a protein consisting of an amino acid sequence shown in SEQ ID NO: 8, 10, 12, 14, 16, 18 or 20,
  - (b) a protein consisting of an amino acid sequence where one or several amino acids are deleted, replaced or added in an amino acid sequence shown in SEQ ID NO: 8, 10, 12, 14, 16, 18 or 20.
- 20. A DNA consisting of a base sequence shown in SEQ ID NO: 7, 9, 11, 13, 15, 17 or 19 or a complementary sequence thereof, and encoding a protein that has a regulatory activity of chromosome segregation.
- 21. A DNA containing part or whole of a base sequence shown in SEQ ID NO: 7, 9, 11, 13, 15, 17 or 19 or a complementary sequence thereof, and encoding a protein that has a regulatory activity of chromosome segregation.
- 22. A DNA hybridizing with the DNA according to claim 7, 9, 11, 13, 15, 17 or 19 under stringent conditions and encoding a protein that has a regulatory activity of chromosome segregation.
- 23. A protein consisting of an amino acid sequence shown in SEQ ID NO: 8, 10, 12, 14, 16, 18 or 20, and having a regulatory activity of chromosome segregation.
- 24. A protein consisting of an amino acid sequence where one or several amino acids are deleted, replaced or added in an amino amino acid sequence shown in SEQ ID NO: 8, 10, 12, 14, 16, 18 or 20, and having a regulatory activity of chromosome segregation.
- 25. A fusion protein in which the protein according to claim 5, 6, 11, 12, 23 or 24 is bound with a marker protein and/or a peptide tag.
- 26. An antibody specifically binding to the protein according to claim 5, 6, 11, 12, 23 or 24.
- 27. The antibody according to claim 26, which is a monoclonal antibody.

\* \* \* \* \*