



US 20110054190A1

(19) **United States**

(12) **Patent Application Publication**
Kobayashi et al.

(10) **Pub. No.: US 2011/0054190 A1**

(43) **Pub. Date: Mar. 3, 2011**

(54) **CATALYST AND REACTION PROCESS**

(30) **Foreign Application Priority Data**

(75) Inventors: **Shu Kobayashi**, Tokyo (JP);
Magno Agostinho, Serra de Santo
Antonio (PT); **Uwe Schneider**,
Tokyo (JP); **Miyuki Yamaguchi**,
Tokyo (JP)

Jan. 28, 2008 (JP) 2008-015906

Publication Classification

(73) Assignee: **JAPAN SCIENCE AND
TECHNOLOGY AGENCY**,
Kawaguchi-shi (JP)

(51) **Int. Cl.**
C07D 207/02 (2006.01)
C07C 67/00 (2006.01)
C07F 3/00 (2006.01)

(52) **U.S. Cl.** **548/540**; 560/54; 564/82

(21) Appl. No.: **12/865,053**

(57) **ABSTRACT**

(22) PCT Filed: **Jan. 28, 2009**

Disclosed is a technology for enabling an efficient asymmetric Michael addition reaction which does not require a large amount of a malonic ester, while having a short reaction time. Specifically disclosed is a catalyst which is composed of MX^2 (wherein M is Be, Mg, Ca, Sr, Ba or Ra and X is an arbitrary group) and a compound represented by general formula [I]. [In the formula, R^7 , R^8 , R^9 and R^{10} each represents a substituted cyclic group or an unsubstituted cyclic group.]

(86) PCT No.: **PCT/JP2009/051322**

§ 371 (c)(1),
(2), (4) Date: **Nov. 18, 2010**

CATALYST AND REACTION PROCESS

APPLICABLE FIELD IN THE INDUSTRY

[0001] The present invention relates to a catalyst and a reaction method. The present invention relates, for example, to a technology of highly enantioselective addition reaction to an enone using a malonic ester as a nucleophile.

BACKGROUND ART

[0002] The reaction between a malonic ester and an enone is shown in the following documents.

[0003] Non-patent document 1: Park, S.-Y.; Morimoto, H.; Matsunaga, S.; Shibasaki, M. *Tetrahedron Lett.* 2007, 48, 2815-2818.

[0004] Non-patent document 2: Chen, C.; Zhu, S.-F.; Wu, X.-Y.; Zhou, Q.-L. *Tetrahedron: Asymmetry* 2006, 17, 2761-2767.

[0005] Non-patent document 3: Kumaraswamy, G.; Jena, N.; Sastry, M. N. V.; Rao, G. V.; Ankamma, K. *J. Mol. Catal. A* 2005, 230, 59-67.

[0006] Non-patent document 4: Velmathi, S.; Swarnalaxshmi, S.; Narasimhan, S. *Tetrahedron: Asymmetry* 2003, 14, 113-117.

[0007] Non-patent document 5: Annamalai, V.; DiMauro, E. F.; Carroll, P. J.; Kozlowski, M. C. *J. Org. Chem.* 2003, 68, 1973-1981.

[0008] Non-patent document 6: Xu, Y.; Otori, K.; Ohsshima, T.; Shibasaki, M. *Tetrahedron* 2002, 58, 2585-2588.

[0009] Non-patent document 7: Kumaraswamy, G.; Sastry, M. N. V.; Jena, N. *Tetrahedron Lett.* 2001, 42, 8515-8517.

[0010] Non-patent document 8: Kim, Y. S.; Matsunaga, S.; Das, J.; Sekine, A.; Ohsshima, T.; Shibasaki, M. *J. Am. Chem. Soc.* 2000, 122, 6506-6507.

[0011] Non-patent document 9: End, N.; Macko, L.; Zehnder, M.; Pfaltz, A. *Chem. Eur. J.* 1998, 4, 818-824.

[0012] Non-patent document 10: Manickam, G.; Sundararajan, G. *Tetrahedron: Asymmetry* 1997, 8, 2271-2278.

[0013] Non-patent document 11: Sasai, H.; Arai, T.; Satow, Y.; Houk, K. N.; Shibasaki, M. *J. Am. Chem. Soc.* 1995, 117, 6194-8.

[0014] Non-patent document 12: Wang, Z.; Wang, Q.; Zhang, Y.; Bao, W. *Tetrahedron Lett.* 2005, 46, 4657-4660.

[0015] Non-patent document 13: Ooi, T.; Ohara, D.; Fukumoto, K.; Maruoka, K. *Org. Lett.* 2005, 7, 3195-3197.

[0016] Non-patent document 14: Dere, R. T.; Pal, R. R.; Patil, P. S.; Salunkhe, M. M. *Tetrahedron Lett.* 2003, 44, 5351-5353.

[0017] Non-patent document 15: Kim, D. Y.; Huh, S. C.; Kim, S. M. *Tetrahedron Lett.* 2001, 42, 6299-6301.

[0018] Non-patent document 16: Wang, J.; Li, H.; Zu, L.; Jiang, W.; Xie, H.; Duan, W.; Wang, W. *J. Am. Chem. Soc.* 2006, 128, 12652-12653.

[0019] Non-patent document 17: Knudsen, K. R.; Mitchell, C. E. T.; Ley, S. V. *Chem. Commun.* 2006, 66-68.

[0020] Non-patent document 18: Halland, N.; Aburel, P. S.; Jorgensen, K. A. *Angew. Chem., Int. Ed.* 2003, 42, 661-665.

[0021] Non-patent document 19: Yamaguchi, M.; Shiraishi, T.; Hiram, M. *J. Org. Chem.* 1996, 61, 3520-30.

[0022] Non-patent document 20: Yamaguchi, M.; Shiraishi, T.; Hiram, M. *Angew. Chem., Int. Ed.* 1993, 32, 1176-8.

DISCLOSURE OF THE INVENTION


Problems to be Solved by the Invention

[0023] Conventionally, an asymmetric Michael addition reaction between a malonic ester and an enone requires a large volume of a malonic ester so as to gain a high yield. Yet, a reaction time thereof is long. For this, the prior art is poor in efficiency.

[0024] Thus, a task that the present invention is to solve, that is, an object of the present invention is to provide, for example, a technology for enabling an efficient asymmetric Michael addition reaction that does not require a large amount of a malonic ester, while having a short reaction time.

Means for Solving the Problem

[0025] The foregoing problems are solved by a catalyst configured using MX_2 (wherein M is Be, Mg, Ca, Sr, Ba or Ra and X is an arbitrary group) and a compound represented by the following general formula [I].



General formula [I]

[0026] $[R^7, R^8, R^9,$ and R^{10} each represents a substituted cyclic group or an unsubstituted cyclic group. There are two cases for R^9 and R^{10} , i.e. the case that they form a ring and the case that they do not form a ring.]

[0027] And, the foregoing problems are solved by the above-mentioned catalyst that is characterized in that the foregoing X is an alkoxide group. Among others, the foregoing problems are solved by the above-mentioned catalyst that is characterized in that the foregoing MX_2 is $M(OR^5)_2$ (M is Mg, Ca, Sr or Ba. R^5 is an alkyl group). More preferably, the foregoing problems are solved by the above-mentioned catalyst that is characterized in that the foregoing MX_2 is $M(OR^5)_2$ (M is Ca, Sr or Ba. R^5 is an alkyl group having a carbon number of 1 to 10). In particular, the foregoing problems are solved by the above-mentioned catalyst that is characterized in that the foregoing MX_2 is $Sr(OR^5)_2$ (R^5 is an alkyl group having a carbon number of 1 to 10).

[0028] Further, the foregoing problems are solved by the above-mentioned catalyst that is characterized in that the foregoing X is an amide group. In particular, the foregoing problems are solved by the above-mentioned catalyst that is characterized in that the foregoing X is hexamethyldisilazide (HMDS).


[0029] Further, the foregoing problems are solved by the above-mentioned catalyst that is characterized in that the foregoing cyclic group of the foregoing general formula [I] is an aromatic group.

[0030] Further, the foregoing problems are solved by the above-mentioned catalyst that is characterized in that the compound represented by the foregoing general formula [I] and M of the foregoing compound MX_2 are coordinate-bonded to each other.

[0031] The above-mentioned catalyst is a catalyst that is used for a reaction between a compound represented by the following general formula [II] and a compound represented by the following general formula [III].



General formula [II]



General formula [III]

[0032] Each of the foregoing R^1 , R^2 , R^3 , and R^4 is an arbitrary substituent. Preferably, it is an H group or a hydrocarbon group.

[0033] Each of the foregoing W^1 , W^2 , and W^3 is an electron-withdrawing group. Preferably, it is an ester group or a carbonyl group.


[0034] The compound represented by the foregoing general formula [II] is, particularly, a dicarboxylate ester. Among others, it is a malonic ester.

[0035] The compound represented by the foregoing general formula [III] is, particularly, an enone.

[0036] Further, the foregoing problems are solved by a reaction method that is characterized in reacting the compound represented by the foregoing general formula [II] with the compound represented by the foregoing general formula [III] in the presence of the foregoing catalyst.

[0037] A molecular sieve is preferably added to a solution of the foregoing reaction. Further, an aromatic hydrocarbon solvent is preferably used as a solvent of the foregoing reaction.

[0038] And, a compound represented by the following general formula [IV] is obtained with the above-mentioned reaction.



General formula [IV]

AN ADVANTAGEOUS EFFECT OF THE INVENTION

[0039] A large amount of the compound of the foregoing general formula [II] such as a malonic ester is not required in the asymmetric Michael addition reaction between the compound of the foregoing general formula [II] (for example, a dicarboxylate ester such as a malonic ester) and the compound of the foregoing general formula [III] (for example, an enone).

[0040] And, the reaction time was shortened.

[0041] That is, the compound of the foregoing general formula [IV] was efficiently obtained.

BEST MODE FOR CARRYING OUT THE INVENTION

[0042] The present invention relates to a catalyst. In particular, the present invention relates a catalyst that is used for the reaction between the compound represented by the

foregoing general formula [II] and the compound represented by the foregoing general formula [III]. Each of the foregoing R^1 , R^2 , R^3 , and R^4 of the foregoing general formula [II] and general formula [III] is an arbitrary substituent. In particular, it is an H group or a hydrocarbon group. Each of the foregoing W^1 , W^2 , and W^3 of the foregoing general formula [II] and general formula [III] is an electron-withdrawing group. For example, it is an electron-withdrawing group such as an ester group, a carboxyl group, a carbonyl group, a nitrile group, a nitro group, and a hydroxyl group. The particularly preferable electron-withdrawing group is an ester group and a carbonyl group. The preferable compound of the foregoing general formula [II] is a dicarboxylate ester. Among others, it is a malonic ester. In particular, it is a malonic ester represented by $R^a\text{OOCCH}_2\text{COOR}^b$ (each of R^a and R^b is a hydrocarbon group. In particular, it is a hydrocarbon group having a carbon number of 1 to 10. For example, it is an alkyl group having a carbon number of 1 to 6.) The preferable compound of the foregoing general formula [III] is an enone.

[0043] The catalyst of the present invention is configured using a compound A and a compound B. The compound A is a compound represented by MX_2 (M is a member selected from a group of alkaline earth metals. X is an arbitrary group.) The compound B is a compound represented by the foregoing general formula [I]. The alkaline earth metal is Be, Mg, Ca, Sr, Ba, or Ra. The preferable alkaline earth metal is Mg, Ca, Sr, or Ba. The particularly preferable alkaline earth metal is Ca, Sr, or Ba. Among others, it is Sr. Any group is acceptable as far as the group X bonded with the alkaline earth metal is concerned. The preferable group, out of the group X, is an alkoxide group. For example, it is an alkoxide group having a carbon number of 1 to 10. More preferably, it is an alkoxide group having a carbon number of 1 to 6. For example, it is a propoxide group such as i-propoxide group, or a butoxide group such as a tert-butoxide group. An amide group is also a preferable group instead of the foregoing alkoxide group. For example, hexamethyldisilazide (HMDS) is a particularly preferable group similarly to the propoxide group. That is, isopropanol is generated within a reaction system when strontium isopropoxide is used as the foregoing MX_2 . On the other hand, hexamethyldisilazane (base) is generated within a reaction system when strontium hexamethyldisilazide is used as the foregoing MX_2 . Thus, the reaction progresses without the base added. Therefore, the above reaction is applicable to other reaction requiring the base. Further, using hexamethyldisilazide allowed the reaction to progress at a high yield and highly enantioselectively, similarly to the case of using isopropoxide. Each of R^7 , R^8 , R^9 , and R^{10} of the foregoing general formula [I] is a cyclic group. There are two cases for this cyclic group, i.e. the case of having a substituent and the case of not having a substituent. The preferable cyclic group is an aromatic group. For example, it is a phenyl group. Or it is a phenyl group having a substituent. Additionally, with regard to R^9 and R^{10} , there is the case that a ring is formed by R^9 and R^{10} . Needless to say, there is case that no ring is formed. And, the catalyst of the present invention is configured using the foregoing compound A and the foregoing compound B. For example, mixing the foregoing compound A and the foregoing compound B allows the catalyst of the present invention to be configured. For example, the catalyst (the catalysts of the present invention) assuming a structure in which the alkaline earth metal has been coordinate-bonded to an asymmetric ligand represented by the foregoing general

formula [I] is configured. A preferable mixture ratio of the foregoing compound A and the foregoing compound B is A:B=1:1 to 2 (mole ratio).

[0044] The present invention also relates to a reaction method. In particular, the present invention relates a reaction method of obtaining the compound represented by the foregoing general formula [IV]. That is, the method of the present invention is a method of reacting the compound represented by the foregoing general formula [II] with the compound represented by the foregoing general formula [III] in the presence of the foregoing catalyst of the present invention. An amount of the catalyst is 0.01 to 20 parts by mass to 100 parts by mass of a substrate. In particular, it is 0.5 to 10 parts by mass. The reaction is conducted at temperature of 0° C. to room temperature. Molecular sieve is preferably added to a solution of this reaction. Further, an aromatic hydrocarbon solvent such as toluene, xylene, and benzene, is preferably used as a solvent.

[0045] Hereinafter, the present invention will be explained more specifically.

[0046] As the compound A represented by MX₂, calcium isopropoxide (Ca(O-i-Pr)₂), strontium isopropoxide (Sr(O-i-Pr)₂), barium butoxide (Ba(O-t-Bu)₂), magnesium butoxide (Mg(O-t-Bu)₂) were used.

[0047] Ca(O-i-Pr)₂ was procured from Sigma-Aldrich Company. Sr(O-i-Pr)₂ and Ba(O-t-Bu)₂ were procured from JAPAN PURE CHEMICAL CO., LTD. Mg(O-t-Bu)₂ was procured from Alfa Aesar Company.

[0048] An asymmetric ligand III represented by the general formula [I] in accordance with the present invention was synthesized with the method described in the document (Evans, D. A.; Nelson, S. G. J. Am. Chem. Soc. 1997, 119, 6452 to 6453). An asymmetric ligand I, being a comparative example, was synthesized with the method described in the document (Lowenthal, R. E.; Abiko, A.; Masamune, S. Tetrahedron Lett. 1990, 31, 6005-8). An asymmetric ligand II, being a comparative example, was synthesized with the method described in the document (Hilgraf, R.; Pfaltz, A. Adv. Synth. Catal. 2005, 347, 61-77).


[0049] A malonic ester 1a (where R is Me), a malonic ester 1b (where R is Et), a malonic ester 1c (where R is n-Pr), a malonic ester 1e (where R is n-Bu), and a malonic ester 1f (where R is Bn (benzyl group)) shown in Table 2 were procured from TOKYO CHEMICAL INDUSTRY CO., LTD. (TCI). A malonic ester 1d (where R is i-Pr) was procured from Wako Pure Chemical industries, LTD.

[0050] α,β -unsaturated carbonyl compounds 2a, 2b, 2c, 2d, 2e, 2f, 2g, 2h, 2i, 2j, 2k, 2l, 2m, 2n, 2o, 2p, 2q, 2t, and 2u shown in Table 3 were procured from TCI, Sigma-Aldrich Company, Alfa Aesar Company, Acros Company, and Wako Pure Chemical industries, LTD. An O-unsaturated carbonyl compound 2r shown in Table 3 was synthesized with the method described in the document (Bhagat, S.; Sharma, R.; Sawant, D. M.; Sharma, L.; Chakraborti, A. K. J. Mol. Catal. A: Chem. 2006, 244, 20-24). An α,β -unsaturated carbonyl compound 2s shown in Table 3 was synthesized with the method described in the document (Evans, D. A.; Borg, G; Scheidt, K. A. Angew. Chem., Int. Ed. 2002, 41, 3188-3191). Additionally, the above-mentioned α,β -unsaturated carbonyl compounds 2b to 2u are equivalent to entries 1 to 20 of Table 3, respectively.

[0051] Molecular sieves (powder) were procured from Aldrich Company, and activated (200° C., <1 mmHg, 16 hours) for use.

[0052] Toluene was procured from Wako Pure Chemical industries, LTD. And, this toluene (anhydride solvent) was distilled in the presence of benzophenone and sodium.


[Asymmetric ligand III]



[0053] (1R,2R)-1,2-Diphenylethylene-1,2-bis(2,5-dimethylphenyl)sulfonamide:

[0054] ¹H NMR (600.2 MHz, THF-D₈, TMS): δ =7.37 (s, 2H; Ar), 7.06-6.81 (m, 12H; Ar, NH), 6.72-6.67 (m, 4H; Ar), 4.43 (m, 2H, CH), 2.40 (s, 6H, CH₃), 2.14 (s, 6H, CH₃); ¹³C {¹H} NMR (150.9 MHz, THF-D₈, TMS): δ =140.0, 138.2, 136.1, 134.6, 133.2, 132.7, 130.3, 128.7, 128.2, 127.9, 63.3, 20.6, 19.7; $[\alpha]_D^{21}$ =+40.27 (c=1.0 in CHCl₃), for the SS enantiomer of III Evans reported $[\alpha]_D^{25}$ =-42 (c=0.96 in CHCl₃).


[Complex A]



[0055] The complex A was prepared by stirring Sr(O-i-Pr)₂ (0.15 mmol) in one equivalent of the asymmetric ligand III, and deuterated THF (0.75 mL) for two hours.

[0056] ¹H NMR (600.2 MHz, THF-D₈, TMS): δ =7.40-6.60 (br m, 16H; Ar), 4.40 (br m, 2H; CH), 3.86-3.79 (m, 2H; CH free i-PrOH), 3.43 (d, 3J_{HH}=3.6 Hz, 2H; OH free i-PrOH), 2.53 (br s, 6H; CH₃), 1.44 (br s, 6H; CH₃), 1.06 (d, J_{HH}=6.1 Hz, 12H; CH₃ free i-PrOH); ¹³C {¹H} NMR (150.9 MHz, THF-D₈, TMS): δ =145.6, 144.8, 134.9, 134.2, 131.6, 130.8, 129.7, 127.7, 125.9, 69.4, 63.6 (free i-PrOH), 25.9 (free i-PrOH), 21.1, 20.3.

[Complex B]




[0057] The complex B was prepared by adding one equivalent of dimethyl malonate (1a, 17 μ L) to THF (H=D) containing the complex A.

[0058] And, NMR thereof was observed in a solution state.

[0059] ^{13}C $\{^1\text{H}\}$ NMR (150.9 MHz, THF- D_8 , TMS) selected data: $\delta=174.6$ (COO, coordinated malonate), 68.8 (NCH, ligand), 64.6 (CH_2 , malonate), 63.6 (free *i*-PrOH), 49.7 (OCH_3 , malonate), 25.9 (free *i*-PrOH), 20.6 (CH_3 , ligand), 19.8 (CH_3 , ligand).


[0060] [A method of Preparing a Crude Material]



3-(5-Methylfuran-2-yl)-1-phenylprop-2-en-1-one

[0061] It was prepared with the method described in the document (Bhagat, S.; Sharma, R.; Sawant, D. M.; Sharma, L.; Chakraborti, A. K. J. Mol. Catal. A: Chem. 2006, 244, 20-24).

[0062] (Yield 63%), Yellow solid, Mp 57-61 $^\circ$ C.: IR[cm^{-1}] (KBr): 1580, 1520, 1367, 1016: ^1H NMR (600.2 MHz, CDCl_3 , TMS): $\delta=8.05$ -8.01 (m, 2H), 7.58-7.46 (m, 4H), 7.38 (d, $J_{\text{HH}}=15.2$ Hz, 1H), 6.62 (d, $J_{\text{HH}}=3.4$ Hz, 1H), 6.13 (d, $J_{\text{HH}}=3.4$ Hz, 1H), 2.39 (s, 3H); ^{13}C $\{^1\text{H}\}$ NMR (150.9 MHz, CDCl_3 , TMS): $\delta=189.8$, 155.9, 150.3, 138.4, 132.5, 130.8, 128.5, 128.4, 118.3, 117.5, 109.4, 109.3, 14.0: ESI-HRMS (m/z) calcd. for $\text{C}_{14}\text{H}_{13}\text{O}_2$ ((M+H) $^+$): 213.0916, found: 213.0924, calcd. for $\text{C}_{14}\text{H}_{12}\text{O}_2\text{Na}$ ((M+Na) $^+$): 235.0735, found: 235.0732.



3-Phenyl-1-(1H-pyrrol-1-yl)prop-2-en-1-one

[0063] It was prepared with the method described in the document (Evans, D. A.; Borg, G.; Scheidt, K. A. Angew. Chem., Int. Ed. 2002, 41, 3188-3191).

[0064] White solid, Mp 101-105 $^\circ$ C.: IR [cm^{-1}] (KBr): 1689, 1624, 1468, 1352: ^1H NMR (495.1 MHz, CDCl_3 , TMS): $\delta=7.99$ (d, $J_{\text{HH}}=15.5$ Hz, 1H), 7.64-7.60 (m, 2H), 7.48-7.41 (m, 5H), 7.14 (d, $J_{\text{HH}}=15.5$ Hz, 1H), 6.36 (appearance of t, $J_{\text{HH}}=2.4$ Hz, 2H); ^{13}C $\{^1\text{H}\}$ NMR (150.9 MHz, CDCl_3 , TMS): $\delta=162.9$, 147.5, 134.2, 130.9, 129.0, 128.4, 119.3, 115.7, 113.4.

[0065] [A General Manipulation of the Catalytic Asymmetric Michael Reaction]

[0066] A flask with a capacity of 30 mL was heated and dried. A toluene (1.0 mL) suspension of $\text{Sr}(\text{O-}i\text{-Pr})_2$ (0.015 mmol), the foregoing ligand III (0.018 mmol), and molecular sieves MS 4A (100 mg) were poured into this flask. And it was stirred for two hours at room temperature.

[0067] Thereafter, a toluene (1.0 mL) solution of diethyl malonate (0.36 mmol) and a toluene (1.0 mL) solution of chalcone (0.30 mmol) were sequentially added.


[0068] After confirming the finishing of the reaction by use of TLC, a saturated ammonium chloride aqueous solution NH_4Cl (10 mL) was added. And an organic phase was separated by adding methylene chloride (CH_2Cl_2 , 10 mL), and was extracted from a water phase with methylene chloride CH_2Cl_2 (15 mL \times 3).

[0069] The organic phase was collected, and dried over anhydrous sodium sulfate.

[0070] After filtering and concentration under reduced pressure, the crude product was refined with a preparative thin-layer chromatography (hexane/ethyl acetate=4/1). With this, the target compound was obtained.

[0071] Enantioselectivity was determined by an HPLC analysis of the target compound.

[0072] Additionally, the above-mentioned reaction formula is shown in the following Table 1 to Table 3.



3aa


[0073] Dimethyl

2-(3-oxo-1,3-diphenylpropyl)malonate (table 2, entry 1)

[0074] It was synthesized in accordance with the above-mentioned manipulation.

[0075] Yield 65%, White solid, Mp 77-80 $^\circ$ C.: IR [cm^{-1}] (KBr): 1730, 1680, 1239, 1157: ^1H NMR (600.2 MHz, CDCl_3 , TMS): $\delta=7.91$ -7.88 (m, 2H; Ar), 7.55-7.50 (m, 1H; Ar), 7.44-7.39 (m, 2H; Ar), 7.27-7.22 (m, 4H; Ar), 7.20-7.15 (m, 1H; Ar), 4.22-4.16 (m, 1H; CH), 3.88-3.85 (m, 1H; CH), 3.72 (s, 3H, CH_3), 3.56-3.46 (m, 5H; CH_2 , CH_3), ^{13}C $\{^1\text{H}\}$ NMR (150.9 MHz, CDCl_3 , TMS): $\delta=197.5$, 168.7, 168.1, 140.4, 136.8, 133.1, 128.5, 128.5, 128.1 (from intensity corresponds to 2 peaks), 127.2, 57.3, 52.6, 42.3, 40.7: HPLC (Chiralpak AS-H, hexane/*i*-propanol=19/1, flow rate 0.5 mL/min, $\lambda=254$ nm): $t_{\text{major}}=38.7$ min, $t_{\text{minor}}=46.3$ min, ee=94%: $[\alpha]_D^{21}=+27.37$ (c=2.0 in CHCl_3), literature value reported by Shibasaki⁷ for the S enantiomer $[\alpha]_D^{24}=+25.64$

($c=2.0$ in CHCl_3 , 77% ee); ESI-HRMS (m/z) calcd. for $\text{C}_{20}\text{H}_{20}\text{O}_5\text{Na}$ ($(\text{M}+\text{Na})^+$): 363.1208, found: 363.1282.



3ba


[0076] Diethyl

2-(3-oxo-1,3-diphenylpropyl)malonate (table 2, entry 2)

[0077] It was synthesized in accordance with the above-mentioned manipulation.

[0078] Yield 97%, White solid, Mp 62-66° C.: IR [cm^{-1}] (KBr): 1731, 1685, 1288, 1241: ^1H NMR (600.2 MHz, CDCl_3 , TMS): $\delta=7.90$ -7.86 (m, 2H; Ar), 7.52-7.49 (m, 1H; Ar), 7.42-7.38 (m, 2H; Ar), 7.28-7.21 (m, 4H; Ar), 7.17-7.13 (m, 1H; Ar), 4.24-4.14 (m, 3H; CH, OCH_2), 3.94 (q, $3J_{\text{HH}}=7.1$ Hz, 2H; OCH_2), 3.83 (d, $3J_{\text{HH}}=9.6$ Hz, 1H; CH), ABM spin system ($A=B=M=H$, $\delta_A=3.54$, $\delta_B=3.46$, $^2J_{AB}=16.6$, $^3J_{AM}=4.4$, $^3J_{BM}=9.1$ Hz, 2H; CH_2), 1.23 (t, $^3J_{\text{HH}}=7.1$ Hz, 3H; CH_3), 1.00 (t, $^3J_{\text{HH}}=7.1$ Hz, 3H; CH_3); ^{13}C { ^1H } NMR (150.9 MHz, CDCl_3 , TMS): $\delta=197.5$, 168.4, 167.1, 140.5, 136.8, 133.0, 128.5, 128.4, 128.2, 128.1, 127.1, 61.6, 61.3, 57.6, 42.6, 40.8, 14.0, 13.8: HPLC (Chiralpak AS-H, hexane/*i*-propanol=19/1, flow rate 0.5 mL/min, $\lambda=254$ nm): $t_{\text{major}}=28.1$ min, $t_{\text{minor}}=31.8$ min, ee=97%

[0079] [α] $^{22}_D=+19.39$ ($c=1.0$ in CHCl_3), [α] $^{19}_D=+6.35$ ($c=2.5$ in benzene), literature value reported by Koga⁹ for the S enantiomer [α] $^{25}_D=+5.4$ ($c=2.61$ in benzene, 82% ee); ESI-HRMS (m/z) calcd. for $\text{C}_{22}\text{H}_{24}\text{O}_5\text{Na}$ ($(\text{M}+\text{Na})^+$): 391.1521, found: 391.1502.



3ca


[0080] Dipropyl

2-(3-oxo-1,3-diphenylpropyl)malonate (table 2, entry 3)

[0081] It was synthesized in accordance with the above-mentioned manipulation.

[0082] Yield 92%, White solid, Mp 55-58° C.: IR [cm^{-1}] (KBr): 1725, 1686, 1293, 1241, 1168: ^1H NMR (600.2 MHz, CDCl_3 , TMS): $\delta=7.89$ -7.87 (m, 2H; Ar), 7.52-7.49 (m, 1H; Ar), 7.42-7.38 (m, 2H; Ar), 7.28-7.21 (m, 4H; Ar), 7.17-7.13 (m, 1H; Ar), 4.19 (t d, $^3J_{\text{HH}}=4.5$, $^3J_{\text{HH}}=9.4$ Hz, 1H; CH),

4.14-4.05 (m, 2H; OCH_2), 3.87-3.83 (m, 3H; CH, OCH_2), ABM spin system ($A=B=M=H$, $\delta_A=3.54$, $\delta_B=3.47$, $^2J_{AB}=16.7$, $^3J_{AM}=4.6$, $^3J_{BM}=9.1$ Hz, 2H; CH_2), 1.63 (appearance of sext, $^3J_{\text{HH}}=7.1$ Hz, 2H; CH_2CH_3), 1.47-1.37 (m, 2H; CH_2CH_3), 0.90 (t, $^3J_{\text{HH}}=7.3$ Hz, 3H; CH_3), 0.77 (t, $3J_{\text{HH}}=7.5$ Hz, 3H; CH_3); ^{13}C { ^1H } NMR (150.9 MHz, CDCl_3 , TMS): $\delta=197.6$, 168.5, 167.9, 140.6, 136.9, 133.0, 128.5, 128.4, 128.2, 128.1, 127.1, 67.2, 66.9, 57.6, 42.6, 40.8, 21.8, 21.6, 10.3, 10.2: HPLC (Chiralpak AS-H, hexane/*i*-propanol=100/1, flow rate 0.5 mL/min, $\lambda=254$ nm): $t_{\text{major}}=47.2$ min, $t_{\text{minor}}=52.0$ min, ee=99%; [α] $^{21}_D=+24.29$ ($c=1.0$ in CHCl_3); ESI-HRMS (m/z) calcd. for $\text{C}_{24}\text{H}_{28}\text{O}_5\text{Na}$ ($(\text{M}+\text{Na})^+$): 419.1834, found: 419.1865.



3da


[0083] Diisopropyl

2-(3-oxo-1,3-diphenylpropyl)malonate (table 2, entry 7)

[0084] It was synthesized in accordance with the above-mentioned manipulation.

[0085] Yield 83%, White solid, Mp 69-71° C.: IR [cm^{-1}] (KBr): 1725, 1685, 1283, 1239, 1106: ^1H NMR (600.2 MHz, CDCl_3 , TMS): $\delta=7.90$ -7.86 (m, 2H; Ar), 7.52-7.48 (m, 1H; Ar), 7.40 (br t, $^3J_{\text{HH}}=7.7$ Hz, 2H; Ar), 7.27-7.19 (m, 4H; Ar), 7.16-7.12 (m, 1H; Ar), 5.07 (sept, $^3J_{\text{HH}}=6.2$ Hz, 1H; CH), 4.79 (sept, $^3J_{\text{HH}}=6.3$ Hz, 1H; CH), 4.16 (t d, $^3J_{\text{HH}}=9.7$, $^3J_{\text{HH}}=4.1$ Hz, 1H; CH), 3.78 (d, $^3J_{\text{HH}}=9.7$ Hz, 1H; CH), ABM spin system ($A=B=M=H$, $\delta_A=3.53$, $\delta_B=3.43$, $^2J_{AB}=16.5$, $^3J_{AM}=4.1$, $^3J_{BM}=9.7$ Hz, 2H; CH_2), 1.23 (d, $^3J_{\text{HH}}=6.3$ Hz, 6H; CH_3), 1.04 (d, $^3J_{\text{HH}}=6.2$ Hz, 3H; CH_3), 0.96 (d, $^3J_{\text{HH}}=6.2$ Hz, 3H; CH_3); ^{13}C { ^1H } NMR (150.9 MHz, CDCl_3 , TMS): $\delta=197.6$, 167.9, 167.2, 140.5, 136.9, 133.0, 128.5, 128.4, 128.3, 128.1, 127.0, 69.2, 68.8, 57.9, 42.9, 40.7, 21.7, 21.5, 21.3, 21.3: HPLC (Chiracel OD-H, hexane/*i*-propanol=9/1, flow rate 0.5 mL/min, $\lambda=254$ nm): $t_{\text{major}}=12.4$ min, $t_{\text{minor}}=13.7$ min, ee=89%; [α] $^{22}_D=+21.27$ ($c=1.0$ in CHCl_3); ESI-HRMS (m/z) calcd. for $\text{C}_{24}\text{H}_{28}\text{O}_5\text{Na}$ ($(\text{M}+\text{Na})^+$): 419.1834, found: 419.1898.

3ea




[0086] Dibutyl

2-(3-oxo-1,3-diphenylpropyl)malonate8 (table 2, entry 8)

[0087] It was synthesized in accordance with the above-mentioned manipulation.

[0088] Yield 85%, Colorless liquid; IR [cm^{-1}] (neat): 1733, 1687, 1254, 1223, 1158; ^1H NMR (600.2 MHz, CDCl_3 ,

TMS): δ =7.90-7.87 (m, 2H; Ar), 7.52-7.48 (m, 1H; Ar), 7.41-7.38 (m, 2H; Ar), 7.28-7.20 (m, 4H; Ar), 7.17-7.13 (m, 1H; Ar), 4.21-4.09 (m, 3H; CH, OCH₂), 3.92-3.84 (m, 3H; CH, OCH₂), ABM spin system (A=B=M=H, δ_A =3.54, δ_B =3.46, $^2J_{AB}$ =16.7, $^3J_{AM}$ =4.5, $^3J_{BM}$ =9.3 Hz, 2H; CH₂), 1.61-1.55 (m, 2H; CH₂), 1.41-1.30 (m, 4H; CH₂), 1.22-1.15 (m, 2H; CH₂), 0.89 (t, $^3J_{HH}$ =7.3 Hz, 3H; CH₃), 0.82 (t, $^3J_{HH}$ =7.3 Hz, 3H; CH₃); ¹³C {¹H} NMR (150.9 MHz, CDCl₃, TMS): δ =197.5, 168.5, 167.8, 140.6, 136.9, 133.0, 128.5, 128.4, 128.2, 128.1, 127.1, 65.5, 65.2, 57.6, 42.6, 40.8, 30.5, 30.3, 19.0, 18.9, 13.6, 13.6; HPLC (Chiralpak AS-H, hexane/i-propanol=100/1, flow rate 1.0 mL/min, λ =254 nm): t_{major} =28.2 min, t_{minor} =29.9 min, ee=96%; $[\alpha]_D^{22}$ =+19.65 (c=1.0 in CHCl₃); ESI-HRMS (m/z) calcd. for C₂₆H₃₂O₅Na ((M+Na)⁺): 447.2147, found: 447.2145, calcd. for C₂₆H₃₃O₅ ((M+H)⁺): 425.2328, found: 425.2316.




3fa

[0089] Dibenzyl

2-(3-oxo-1,3-diphenylpropyl)malonate 11 (table 2, entry 9)

[0090] It was synthesized in accordance with the above-mentioned manipulation.

[0091] Yield 85%, White solid, Mp 89-92° C.; IR [cm⁻¹] (KBr): 1735, 1682, 1230, 1154; ¹H NMR (600.2 MHz, CDCl₃, TMS): δ =7.82-7.77 (m, 2H; Ar), 7.52-7.47 (m, 1H; Ar), 7.39-7.35 (m, 2H; Ar), 7.29-7.13 (m, 13H; Ar), 7.07-7.04 (m, 2H; Ar), 5.18-5.09 (m, 2H; OCH₂), 4.90 (s, 2H; OCH₂), 4.25-4.19 (m, 1H; CH), 3.97-3.92 (m, 1H; CH), 3.47-3.42 (m, 2H; CH₂); ¹³C {¹H} NMR (150.9 MHz, CDCl₃, TMS): δ =197.3, 168.0, 167.5, 140.3, 136.7, 135.1, 135.0, 133.0, 128.5, 128.4, 128.3, 128.1, 128.0, 127.2, 67.3, 67.1, 57.5, 42.2, 40.7; HPLC (Chiralpak AS-H, hexane/i-propanol=19/1, flow rate 0.5 mL/min, λ =254 nm): t_{major} =56.3 min, t_{minor} =63.7 min, ee=84%; $[\alpha]_D^{21}$ =+17.55 (c=1.0 in CHCl₃); ESI-HRMS (m/z) calcd. for C₃₂H₂₈O₅Na ((M+Na)⁺): 515.1834, found: 515.1847.



3cb


[0092] Dipropyl

2-(1-(2-chlorophenyl)-3-oxo-3-phenylpropyl)malonate (table 3, entry 1)

[0093] It was synthesized in accordance with the above-mentioned manipulation.

[0094] Yield 76%, Colorless liquid; IR [cm⁻¹] (neat): 1730, 1687, 1266, 1227, 1160, 738; ¹H NMR (600.2 MHz, CDCl₃, TMS): δ =7.94-7.90 (m, 2H; Ar), 7.54-7.49 (m, 1H; Ar), 7.41 (appearance of br t, $^3J_{HH}$ =7.8 Hz, 2H; Ar), 7.34-7.30 (m, 2H; Ar), 7.16-7.09 (m, 2H; Ar), 4.66 (t, $^3J_{HH}$ =8.8, $^3J_{HH}$ =4.3 Hz, 1H; CH), 4.13-4.00 (m, 3H; CH, OCH₂), 3.94 (t, $^3J_{HH}$ =6.7 Hz, 2H; OCH₂), ABM spin system (A=B=M=H, δ_A =3.71, δ_B =3.62, $^2J_{AB}$ =17.2, $^3J_{AM}$ =9.1, $^3J_{BM}$ =4.3 Hz, 2H; CH₂), 1.63-1.45 (m, 4H; CH₂CH₃), 0.86 (t, $^3J_{HH}$ =7.3 Hz, 3H; CH₃), 0.83 (t, $^3J_{HH}$ =7.5 Hz, 3H; CH₃); ¹³C {¹H} NMR (150.9 MHz, CDCl₃, TMS): δ =197.5, 168.4, 167.9, 137.9, 136.8, 134.1, 133.1, 130.1, 129.4, 128.5, 128.2, 128.1, 126.8, 67.1, 67.1, 55.2, 40.3, 37.4, 21.8, 21.7, 10.2, 10.2; HPLC (Chiralcel OJ-H, hexane/i-propanol=9/1, flow rate 0.5 mL/min, λ =254 nm): t_{major} =20.5 min, t_{minor} =26.6 min, ee=92%; $[\alpha]_D^{22}$ =+41.20 (c=1.0 in CHCl₃), ESI-HRMS (m/z) calcd. for C₂₄H₂₇ClO₅Na ((M+Na)⁺): 453.1445, found: 453.1437.


3cc

**[0095]** Dipropyl

2-(1-(4-chlorophenyl)-3-oxo-3-phenylpropyl)malonate (table 3, entry 2)

[0096] It was synthesized in accordance with the above-mentioned manipulation.

[0097] Yield 93%, White solid, Mp 91-94° C.; IR [cm⁻¹] (KBr): 1727, 1686, 1239; ¹H NMR (600.2 MHz, CDCl₃, TMS): δ =7.88 (br d, $^3J_{HH}$ =7.2 Hz, 2H; Ar), 7.52 (appearance of br t, $^3J_{HH}$ =7.3 Hz, 1H; Ar), 7.41 (t, $^3J_{HH}$ =7.8 Hz, 2H; Ar), 7.23-7.19 (m, 4H; Ar), 4.19-4.05 (m, 3H; CH, OCH₂), 3.88 (t, $^3J_{HH}$ =6.6 Hz, 2H; OCH₂), 3.82 (d, $^3J_{HH}$ =9.5 Hz, 1H; CH), ABM spin system (A=B=M=H, δ_A =3.53, δ_B =3.45, $^2J_{AB}$ =17.0, $^3J_{AM}$ =4.2, $^3J_{BM}$ =9.5 Hz, 2H; CH₂), 1.64 (appearance of sext, $^3J_{HH}$ =7.2 Hz, 2H; CH₂CH₃), 1.50-1.41 (m, 2H; CH₂CH₃), 0.90 (t, $^3J_{HH}$ =7.3 Hz, 3H; CH₃), 0.79 (t, $^3J_{HH}$ =7.5 Hz, 3H; CH₃); ¹³C {¹H} NMR (150.9 MHz, CDCl₃, TMS): δ =197.3, 168.2, 167.7, 139.1, 136.7, 133.2, 132.9, 129.7, 128.6, 128.5, 128.0, 67.3, 67.1, 57.4, 42.4, 40.2, 21.8, 21.7, 10.3, 10.2; HPLC (Chiralcel OJ-H, hexane/i-propanol=19/1, flow rate 0.3 mL/min, λ =254 nm): t_{major} =47.8 min, t_{minor} =54.8 min, ee=97%; $[\alpha]_D^{22}$ =+24.09 (c=1.0 in CHCl₃); ESI-HRMS (m/z) calcd. for C₂₄H₂₇ClO₅Na ((M+Na)⁺): 453.1445, found: 453.1486.




3cd

[0098] Dipropyl

2-(1-(4-fluorophenyl)-3-oxo-3-phenylpropyl)malonate (table 3, entry 3)

[0099] It was synthesized in accordance with the above-mentioned manipulation.

[0100] Yield 92%, White solid, Mp 35-38° C.; IR [cm⁻¹] (neat): 1733, 1687, 1510, 1226, 1161, 739; ¹H NMR (600.2 MHz, CDCl₃, TMS): δ=7.91-7.87 (m, 2H; Ar), 7.52 (appearance of br t, ³J_{HH}=7.5 Hz, 1H; Ar), 7.42 (t, ³J_{HH}=7.8 Hz, 2H; Ar), 7.27-7.23 (m, 2H; Ar), 6.92 (appearance of br t, ³J_{HH}=8.7 Hz, 2H; Ar), 4.20-4.06 (m, 3H; CH, OCH₂), 3.87 (t, ³J_{HH}=6.8 Hz, 2H; OCH₂), 3.82 (d, ³J_{HH}=9.6 Hz, 1H; CH), ABM spin system (A=B=M=H, δ_A=3.53, δ_B=3.44, ²J_{AB}=16.6, ³J_{AM}=4.3, ³J_{BM}=9.6 Hz, 2H; CH₂), 1.64 (appearance of sext, ³J_{HH}=7.0 Hz, 2H; CH₂CH₃), 1.50-1.40 (m, 2H; CH₂CH₃), 0.91 (t, ³J_{HH}=7.5 Hz, 3H; CH₃), 0.79 (t, ³J_{HH}=7.5 Hz, 3H; CH₃); ¹³C {¹H} NMR (150.9 MHz, CDCl₃, TMS): δ=197.4, 168.3, 167.8, 161.8 (d, J_{CF}=245.4 Hz), 136.8, 136.3, 133.1, 129.9 (d, J_{CF}=7.9 Hz), 128.6, 128.1, 115.2 (d, J_{CF}=21.2 Hz), 67.3, 67.0, 57.6, 42.6, 40.1, 21.8, 21.7, 10.3, 10.2; HPLC (Chiracel OJ-H, hexane/*i*-propanol=19/1, flow rate 0.3 mL/min, λ=254 nm): t_{major}=50.7 min, t_{minor}=63.1 min, ee=98%; [α]_D²¹=+25.21 (c=1.0 in CHCl₃); ESI-HRMS (m/z) calcd. for C₂₄H₂₇FO₅Na ((M+Na)⁺): 437.1740, found: 437.1728.



3ce


[0101] Dipropyl

2-(1-(4-methoxyphenyl)-3-oxo-3-phenylpropyl)malonate (table 3, entry 4)

[0102] It was synthesized in accordance with the above-mentioned manipulation.

[0103] Yield 80%, White solid, Mp 41-44° C.; IR [cm⁻¹] (KBr): 1729, 1678, 1511, 1243, 1162; ¹H NMR (600.2 MHz, CDCl₃, TMS): δ=7.89 (br d, ³J_{HH}=7.1 Hz, 2H; Ar), 7.50

(appearance of br t, ³J_{HH}=7.5 Hz, 1H; Ar), 7.40 (t, ³J_{HH}=7.8 Hz, 2H; Ar), 7.17 (d, ³J_{HH}=8.7 Hz, 2H; Ar), 6.76 (d, ³J_{HH}=8.5 Hz, 2H; Ar), 4.16-4.05 (m, 3H; CH, OCH₂), 3.86 (t, ³J_{HH}=6.6 Hz, 2H; OCH₂), 3.82 (d, ³J_{HH}=9.6 Hz, 1H; CH), 3.72 (s, 3H; OCH₃), ABM spin system (A=B=M=H, δ_A=3.51, δ_B=3.42, ²J_{AB}=16.5, ³J_{AM}=4.2, ³J_{BM}=9.5 Hz, 2H; CH₂), 1.64 (appearance of next, ³J_{HH}=7.1 Hz, 2H; CH₂CH₃), 1.50-1.40 (m, 2H; CH₂CH₃), 0.91 (t, ³J_{HH}=7.5 Hz, 3H; CH₃), 0.79 (t, ³J_{HH}=7.5 Hz, 3H; CH₃); ¹³C {¹H} NMR (150.9 MHz, CDCl₃, TMS): δ=197.7, 168.5, 167.9, 158.6, 136.9, 133.0, 132.5, 129.2, 128.5, 128.1, 113.8, 67.2, 66.9, 57.8, 55.1, 42.8, 40.2, 21.8, 21.7, 10.3, 10.2; HPLC (Chiracel OJ-H, hexane/*i*-propanol=9/1, flow rate 1.5 mL/min, λ=254 nm): t_{major}=17.6 min, t_{minor}=11.0 min, ee>99%; [α]_D²¹=+17.51 (c=1.0 in CHCl₃); ESI-HRMS (m/z) calcd. for C₂₅H₃₀O₆Na ((M+Na)⁺): 449.1940, found: 449.1939.




3cf

[0104] Dipropyl

2-(1-(4-nitrophenyl)-3-oxo-3-phenylpropyl)malonate (table 3, entry 5)

[0105] It was synthesized in accordance with the above-mentioned manipulation.

[0106] Yield 98%, White solid, Mp 82-86° C.; IR [cm⁻¹] (KBr): 1726, 1686, 1519, 1346, 1239, 1151; ¹H NMR (600.2 MHz, CDCl₃, TMS): δ=8.12-8.09 (m, 2H; Ar), 7.90-7.87 (m, 2H; Ar), 7.56-7.49 (m, 3H; Ar), 7.45-7.41 (m, 2H; Ar), 4.31 (t d, ³J_{HH}=4.3, ³J_{HH}=9.4 Hz, 1H; CH), 4.17-4.07 (m, 2H; OCH₂), 3.93-3.86 (m, 3H; CH, OCH₂), ABM spin system (A=B=M=H, δ_A=3.61, δ_B=3.55, ²J_{AB}=17.3, ³J_{AM}=4.2, ³J_{BM}=9.5 Hz, 2H; CH₂), 1.64 (appearance of sext, ³J_{HH}=7.0 Hz, 2H; CH₂CH₃), 1.52-1.43 (m, 2H; CH₂CH₃), 0.91 (t, ³J_{HH}=7.3 Hz, 3H; CH₃), 0.80 (t, ³J_{HH}=7.4 Hz, 3H; CH₃); ¹³C {¹H} NMR (150.9 MHz, CDCl₃, TMS): δ=196.8, 167.9, 167.4, 148.6, 147.0, 136.5, 133.4, 129.4, 128.7, 128.0, 123.6, 67.5, 67.3, 56.9, 42.0, 40.4, 21.8, 21.7, 10.2, 10.2; HPLC (Chiracel OJ-H, hexane/*i*-propanol=9/1, flow rate 1.0 mL/min, λ=254 nm): t_{major}=28.5 min, t_{minor}=41.2 min, ee=96%; [α]_D²²=+31.37 (c=1.0 in CHCl₃); ESI-HRMS (m/z) calcd. for C₂₄H₂₇NO₇Na ((M+Na)⁺): 464.1685, found: 464.1644.




3cg

[0107] Dipropyl 2-(1-(3-nitrophenyl)-3-oxo-3-phenylpropyl)malonate (table 3, entry 6)

[0108] It was synthesized in accordance with the above-mentioned manipulation.

[0109] Yield 94%, Colorless liquid; IR [cm^{-1}] (neat): 1731, 1686, 1638, 1532, 1352, 1265, 738; ^1H NMR (600.2 MHz, CDCl_3 , TMS): δ =8.18 (t, $J_{\text{HH}}=1.9$ Hz, 1H; Ar), 8.04 (d d, $J_{\text{HH}}=1.8$ Hz, $J_{\text{HH}}=8.2$ Hz, 1H; Ar), 7.89 (br d, $J_{\text{HH}}=7.2$ Hz, 2H; Ar), 7.71 (br d, $J_{\text{HH}}=7.8$ Hz, 1H; Ar), 7.54 ((br t, $J_{\text{HH}}=7.3$ Hz, 1H; Ar), 7.46-7.41 (m, 3H; Ar), 4.32 (t d, $^3J_{\text{HH}}=4.3$, $^3J_{\text{HH}}=9.3$ Hz, 1H; CH), 4.17-4.08 (m, 2H; OCH_2), 3.92-3.88 (m, 3H; CH, OCH_2), ABM spin system (A=B=M=H, $\delta_A=3.63$, $\delta_B=3.57$, $^2J_{\text{AB}}=17.4$, $^3J_{\text{AM}}=4.4$, $^3J_{\text{BM}}=9.4$ Hz, 2H; CH_2), 1.64 (appearance of sext, $^3J_{\text{HH}}=7.1$ Hz, 2H; CH_2CH_3), 1.52-1.41 (m, 2H; CH_2CH_3), 0.91 (t, $^3J_{\text{HH}}=7.5$ Hz, 3H; CH_3), 0.79 (t, $^3J_{\text{HH}}=7.4$ Hz, 3H; CH_3); ^{13}C { ^1H } NMR (150.9 MHz, CDCl_3 , TMS): δ =196.9, 168.0, 167.5, 148.2, 143.1, 136.5, 135.3, 133.4, 129.3, 128.7, 128.0, 123.1, 122.2, 67.5, 67.2, 57.0, 42.1, 40.2, 21.8, 21.7, 10.2, 10.1; HPLC (Chiralpak AD-H, hexane/*i*-propanol=9/1, flow rate 1.0 mL/min, λ =254 nm): $t_{\text{major}}=48.9$ min, $t_{\text{minor}}=32.1$ min, ee=94%; $[\alpha]_D^{23}=+31.81$ (c=1.0 in CHCl_3); ESI-HRMS (m/z) calcd. for $\text{C}_{24}\text{H}_{27}\text{NO}_7\text{Na}$ ((M+Na) $^+$): 464.1685, found: 464.1658.



3ci


[0113] Dipropyl

2-(3-(4-fluorophenyl)-1-(4-methoxyphenyl)-3-oxopropyl)malonate (table 3, entry 8)

[0114] It was synthesized in accordance with the above-mentioned manipulation.

[0115] Yield 81%, White solid, Mp 60-63° C.; IR [cm^{-1}] (neat): 1731, 1686, 1599, 1514, 1266, 1251, 1157, 739; ^1H NMR (600.2 MHz, CDCl_3 , TMS): δ =7.94-7.91 (m, 2H; Ar), 7.18-7.15 (m, 2H; Ar), 7.07 (appearance of br t, $^3J_{\text{HH}}=8.6$ Hz, 2H; Ar), 6.78-6.75 (m, 2H; Ar), 4.15-4.05 (m, 3H; CH, OCH_2), 3.87 (t, $^3J_{\text{HH}}=6.7$ Hz, 2H; OCH_2), 3.82 (d, $^3J_{\text{HH}}=9.6$ Hz, 1H; CH), ABM spin system (A=B=M=H, $\delta_A=3.53$, $\delta_B=3.40$, $^2J_{\text{AB}}=16.7$, $^3J_{\text{AM}}=4.3$, $^3J_{\text{BM}}=9.6$ Hz, 2H; CH_2), 1.68-1.61 (m, 2H; CH_2CH_3), 1.50-1.40 (m, 2H; CH_2CH_3), 0.91 (t, $^3J_{\text{HH}}=7.5$ Hz, 3H; CH_3), 0.79 (t, $^3J_{\text{HH}}=7.4$ Hz, 3H; CH_3); ^{13}C { ^1H } NMR (150.9 MHz, CDCl_3 , TMS): δ =196.2, 168.6, 167.9, 165.7 (d, $J_{\text{CF}}=254.6$ Hz), 158.6, 133.4, 132.3, 130.8 (d, $J_{\text{CF}}=9.2$ Hz), 129.2, 115.6 (d, $J_{\text{CF}}=21.8$ Hz), 113.9, 67.2, 67.0, 57.8, 55.1, 42.7, 40.3, 21.9, 21.7, 10.3, 10.2; HPLC (Chiralcel OD-H, hexane/*i*-propanol=19/1, flow rate 0.5 mL/min, λ =254 nm): $t_{\text{major}}=33.0$ min, $t_{\text{minor}}=29.4$ min, ee>99%; $[\alpha]_D^{21}=+16.28$ (c=1.0 in CHCl_3); ESI-HRMS (m/z) calcd. for $\text{C}_{25}\text{H}_{29}\text{FO}_6\text{Na}$ ((M+Na) $^+$): 467.1846, found: 467.1852.

3ch




[0110] Dipropyl

2-(1,3-bis(4-fluorophenyl)-3-oxopropyl)malonate (table 3, entry 7)

[0111] It was synthesized in accordance with the above-mentioned manipulation.

[0112] Yield 91%, Colorless liquid; IR [cm^{-1}] (neat): 1732, 1686, 1600, 1510, 1228, 1157; ^1H NMR (600.2 MHz, CDCl_3 , TMS): δ =7.94-7.91 (m, 2H; Ar), 7.26-7.22 (m, 2H; Ar), 7.11-7.06 (m, 2H; Ar), 6.95-6.90 (m, 2H; Ar), 4.18-4.06 (m, 3H; CH, OCH_2), 3.86 (t, $^3J_{\text{HH}}=6.6$ Hz, 2H; OCH_2), 3.81 (d, $^3J_{\text{HH}}=9.6$ Hz, 1H; CH), 3.72 (s, 3H; OCH_3), ABM spin system (A=B=M=H, $\delta_A=3.50$, $\delta_B=3.38$, $^2J_{\text{AB}}=16.4$, $^3J_{\text{AM}}=4.2$, $^3J_{\text{BM}}=9.6$ Hz, 2H; CH_2), 1.64 (appearance of sext, $^3J_{\text{HH}}=7.1$ Hz, 2H; CH_2CH_3), 1.50-1.40 (m, 2H; CH_2CH_3), 0.91 (t, $^3J_{\text{HH}}=7.5$ Hz, 3H; CH_3), 0.79 (t, $^3J_{\text{HH}}=7.4$ Hz, 3H; CH_3); ^{13}C { ^1H } NMR (150.9 MHz, CDCl_3 , TMS): δ =195.9, 168.4, 167.8, 165.8 (d, $J_{\text{CF}}=254.7$ Hz), 161.7 (d, $J_{\text{CF}}=245.7$ Hz), 136.1, 133.2, 130.8 (d, $J_{\text{CF}}=9.0$ Hz), 129.9 (d, $J_{\text{CF}}=7.9$ Hz), 115.7 (d, $J_{\text{CF}}=21.8$ Hz), 115.3 (d, $J_{\text{CF}}=21.3$ Hz), 67.3, 67.1, 57.6, 42.6, 40.2, 21.9, 21.7, 10.3, 10.2; HPLC (Chiralpak AS-H, hexane/*i*-propanol=9/1, flow rate 0.5 mL/min, λ =254 nm): $t_{\text{major}}=20.3$ min, $t_{\text{minor}}=23.4$ min, ee=96%; $[\alpha]_D^{22}=+21.56$ (c=1.0 in CHCl_3); ESI-HRMS (m/z) calcd. for $\text{C}_{24}\text{H}_{26}\text{F}_2\text{O}_5\text{Na}$ ((M+Na) $^+$): 455.1646, found: 455.1668.



3cj


[0116] Dipropyl

2-(1-(3,4-dimethoxyphenyl)-3-(4-fluorophenyl)-3-oxopropyl)malonate (table 3, entry 9)

[0117] It was synthesized in accordance with the above-mentioned manipulation.

[0118] Yield 61%, Colorless liquid; IR [cm^{-1}] (neat): 1728, 1686, 1598, 1519, 1265, 746, 705; ^1H NMR (600.2 MHz, CDCl_3 , TMS): δ =7.95-7.91 (m, 2H; Ar), 7.11-7.06 (m, 2H;

Ar), 6.79-6.72 (m, 3H; Ar), 4.16-4.05 (m, 3H; CH, OCH₂), 3.88 (t, ³J_{HH}=6.6 Hz, 2H; OCH₂), 3.84 (d, ³J_{HH}=9.6 Hz, 1H; CH), 3.82 (s, 3H; OCH₃), 3.80 (s, 3H; OCH₃), ABM spin system (A=B=M=H, δ_A=3.50, δ_B=3.39, ²J_{AB}=16.3, ³J_{AM}=4.3, ³J_{BM}=9.4 Hz, 2H; CH₂), 1.64 (appearance of sext, ³J_{HH}=7.1 Hz, 2H; CH₂CH₃), 1.51-1.41 (m, 2H; CH₂CH₃), 0.91 (t, ³J_{HH}=7.4 Hz, 3H; CH₃), 0.80 (t, ³J_{HH}=7.4 Hz, 3H; CH₃); ¹³C {¹H} NMR (150.9 MHz, CDCl₃, TMS): δ=196.3, 168.5, 167.9, 165.7 (d, J_{CF}=254.8 Hz), 148.7, 148.1, 133.4, 132.9, 130.8 (d, J_{CF}=9.2 Hz), 120.0, 115.6 (d, J_{CF}=21.8 Hz), 111.9, 111.2, 67.2, 67.0, 57.7, 55.9, 55.8, 42.7, 40.7, 21.9, 21.7, 10.3, 10.2; HPLC (Chiralpak AS-H, hexane/i-propanol=9/1, flow rate 1.0 mL/min, λ=254 nm): t_{major}=16.4 min, t_{minor}=14.0 min, ee=96%; [α]_D²²=+17.02 (c=0.8 in CHCl₃); ESI-HRMS (m/z) calcd. for C₂₆H₃₁FO₇Na ((M+Na)⁺): 497.1951, found: 497.1966.



3cl


[0122] Dipropyl

2-(1-(2-chlorophenyl)-3-(4-fluorophenyl)-3-oxopropyl)malonate (table 3, entry 11)

[0123] It was synthesized in accordance with the above-mentioned manipulation.

[0124] Yield 80%, Colorless liquid; IR [cm⁻¹] (neat): 1732, 1686, 1599, 1265, 1232, 1156, 738, 705; ¹H NMR (600.2 MHz, CDCl₃, TMS): δ=7.97-7.94 (m, 2H; Ar), 7.34-7.28 (m, 2H; Ar), 7.16-7.06 (m, 4H; Ar), 4.64 (t d, ³J_{HH}=5.5, ³J_{HH}=8.3 Hz, 1H; CH), 4.12-4.01 (m, 3H; CH, OCH₂), 3.94 (t, ³J_{HH}=6.6 Hz, 2H; OCH₂), ABM spin system (A=B=M=H, δ_A=3.64, δ_B=3.60, ²J_{AB}=16.7, ³J_{AM}=8.1, ³J_{BM}=5.1 Hz, 2H; CH₂), 1.60 (appearance of sext, ³J_{HH}=7.1 Hz, 2H; CH₂CH₃), 1.55-1.45 (m, 2H; CH₂CH₃), 0.87 (t, ³J_{HH}=7.3 Hz, 3H; CH₃), 0.83 (t, ³J_{HH}=7.4 Hz, 3H; CH₃); ¹³C {¹H} NMR (150.9 MHz, CDCl₃, TMS): δ=196.0, 168.4, 167.9, 165.7 (d, J_{CF}=254.8 Hz), 137.7, 134.1, 133.3, 130.8 (d, J_{CF}=9.3 Hz), 130.1, 129.3, 128.3, 126.8, 115.6 (d, J_{CF}=21.8 Hz), 67.2, 67.1, 55.2, 40.4, 37.5, 21.8, 21.7, 10.2, 10.2; HPLC (Chiralpak AD-H, hexane/i-propanol=9/1, flow rate 1.0 mL/min, λ=254 nm): t_{major}=39.9 min, t_{minor}=16.8 min, ee=93%; [α]_D²²=+35.13 (c=1.0 in CHCl₃); ESI-HRMS (m/z) calcd. for C₂₄H₂₆ClFO₅Na ((M+Na)⁺): 471.1350, found: 471.1335.

3ck


**[0119]** Dipropyl

2-(1-(4-chlorophenyl)-3-(4-fluorophenyl)-3-oxopropyl)malonate (table 3, entry 10)

[0120] It was synthesized in accordance with the above-mentioned manipulation.

[0121] Yield 97%, White solid, Mp 69-71° C.; IR [cm⁻¹] (KBr): 1730, 1685, 1600, 1300, 1238, 1157; ¹H NMR (600.2 MHz, CDCl₃, TMS): δ=7.94-7.90 (m, 2H; Ar), 7.21 (br s, 4H; Ar), 7.11-7.07 (m, 2H; Ar), 4.17-4.05 (m, 3H; CH, OCH₂), 3.88 (t, ³J_{HH}=6.7 Hz, 2H; OCH₂), 3.82 (d, ³J_{HH}=9.6 Hz, 1H; CH), ABM spin system (A=B=M=H, δ_A=3.53, δ_B=3.41, ²J_{AB}=16.6, ³J_{AM}=4.2, ³J_{BM}=9.6 Hz, 2H; CH₂), 1.64 (appearance of sext, ³J_{HH}=7.1 Hz, 2H; CH₂CH₃), 1.51-1.41 (m, 2H; CH₂CH₃), 0.91 (t, ³J_{HH}=7.3 Hz, 3H; CH₃), 0.78 (t, ³J_{HH}=7.5 Hz, 3H; CH₃); ¹³C {¹H} NMR (150.9 MHz, CDCl₃, TMS): δ=195.7, 168.3, 167.7, 165.8 (d, J_{CF}=255.0 Hz), 139.0, 133.2, 133.0, 130.8 (d, J_{CF}=9.1 Hz), 129.7, 128.6, 115.7 (d, J_{CF}=21.9 Hz), 67.3, 67.1, 57.3, 42.3, 21.8, 21.7, 10.2, 10.2; HPLC (Chiralcel OD-H, hexane/i-propanol=19/1, flow rate 0.5 mL/min, λ=254 nm): t_{major}=22.8 min, t_{minor}=20.1 min, ee=97%; [α]_D²⁰=+20.30 (c=1.0 in CHCl₃); ESI-HRMS (m/z) calcd. for C₂₄H₂₆ClFO₅Na ((M+Na)⁺): 471.1350, found: 471.1350.

3cm


**[0125]** Dipropyl

2-(1-(4-fluorophenyl)-3-oxo-3-p-tolylpropyl)malonate (table 3, entry 12)

[0126] It was synthesized in accordance with the above-mentioned manipulation.

[0127] Yield 90%, White solid, Mp 64-67° C.; IR [cm⁻¹] (KBr): 1727, 1673, 1606, 1513, 1296, 1242, 836; ¹H NMR (600.2 MHz, CDCl₃, TMS): δ=7.78 (d, ³J_{HH}=8.2 Hz, 2H; Ar), 7.27-7.20 (m, 4H; Ar), 6.93-6.89 (m, 2H; Ar), 4.20-4.05 (m, 3H; CH, OCH₂), 3.86 (t, ³J_{HH}=6.6 Hz, 2H; OCH₂), 3.82 (d, ³J_{HH}=9.6 Hz, 1H; CH), ABM spin system (A=B=M=H,

$\delta_A=3.49$, $\delta_B=3.41$, $^2J_{AB}=16.6$, $^3J_{AM}=4.2$, $^3J_{BM}=9.5$ Hz, 2H; CH₂), 2.37 (s, 3H; CH₃), 1.64 (appearance of sext, $^3J_{HH}=7.1$ Hz, 2H; CH₂CH₃), 1.49-1.40 (m, 2H; CH₂CH₃), 0.91 (t, $^3J_{HH}=7.5$ Hz, 3H; CH₃), 0.79 (t, $^3J_{HH}=7.4$ Hz, 3H; CH₃); ¹³C {¹H} NMR (150.9 MHz, CDCl₃, TMS): $\delta=197.0$, 168.3, 167.8, 161.8 (d, $J_{CF}=245.3$ Hz), 143.9, 136.3, 134.3, 129.9 (d, $J_{CF}=7.6$ Hz), 129.2, 128.2, 115.2 (d, $J_{CF}=21.3$ Hz), 67.2, 67.0, 57.6, 42.4, 40.2, 21.8, 21.6, 21.5, 10.2, 10.1; HPLC (Chiralpak AS-H, hexane/i-propanol=9/1, flow rate 0.5 mL/min, $\lambda=254$ nm): $t_{major}=30.8$ min, $t_{minor}=33.5$ min, ee=98%; $[\alpha]_D^{21}=+24.25$ (c=1.0 in CHCl₃); ESI-HRMS (m/z) calcd. for C₂₅H₂₉FO₅Na ((M+Na)⁺): 451.1897, found: 451.1859.




3cn

[0128] Dipropyl

2-(3-(4-chlorophenyl)-3-oxo-1-phenylpropyl)malonate (table 3, entry 13)

[0129] It was synthesized in accordance with the above-mentioned manipulation.

[0130] Yield 98%, Colorless liquid; IR [cm⁻¹] (neat): 1731, 1687, 1589, 1265, 743, 703; ¹H NMR (600.2 MHz, CDCl₃, TMS): $\delta=7.84$ -7.81 (m, 2H; Ar), 7.39-7.36 (m, 2H; Ar), 7.26-7.21 (m, 4H; Ar), 7.17-7.14 (m, 1H; Ar), 4.18-4.05 (m, 3H; CH, OCH₂), 3.87-3.84 (m, 3H; CH, OCH₂), ABM spin system (A=B=M=H, $\delta_A=3.53$, $\delta_B=3.42$, $^2J_{AB}=16.6$, $^3J_{AM}=4.4$, $^3J_{BM}=9.4$ Hz, 2H; CH₂), 1.63 (appearance of sext, $^3J_{HH}=7.0$ Hz, 2H; CH₂CH₃), 1.47-1.38 (m, 2H; CH₂CH₃), 0.90 (t, $^3J_{HH}=7.5$ Hz, 3H; CH₃), 0.77 (t, $^3J_{HH}=7.4$ Hz, 3H; CH₃); ¹³C {¹H} NMR (150.9 MHz, CDCl₃, TMS): $\delta=196.5$, 168.5, 167.8, 140.3, 139.4, 135.2, 129.6, 128.8, 128.5, 128.2, 127.2, 67.2, 67.0, 57.5, 42.6, 40.9, 21.8, 21.6, 10.3, 10.2; HPLC (2 \times Chiralcel OJ-H, hexane/i-propanol=9/1, flow rate 0.5 mL/min, $\lambda=254$ nm): $t_{major}=76.2$ min, $t_{minor}=68.8$ min, ee=99%; $[\alpha]_D^{21}=+17.53$ (c=1.0 in CHCl₃); ESI-HRMS (m/z) calcd. for C₂₄H₂₇ClO₅Na ((M+Na)⁺): 453.1445, found: 453.1418.



3co


[0131] Dipropyl

2-(3-(4-fluorophenyl)-3-oxo-1-phenylpropyl)malonate (table 3, entry 14)

[0132] It was synthesized in accordance with the above-mentioned manipulation.

[0133] Yield 92%, Colorless liquid; IR [cm⁻¹] (neat): 1731, 1686, 1599, 1265, 1157, 745, 703; ¹H NMR (600.2 MHz, CDCl₃, TMS): $\delta=7.94$ -7.90 (m, 2H; Ar), 7.27-7.21 (m, 4H; Ar), 7.18-7.14 (m, 1H; Ar), 7.09-7.05 (m, 2H; Ar), 4.19-4.05 (m, 3H; CH, OCH₂), 3.87-3.83 (m, 3H; CH, OCH₂), ABM spin system (A=B=M=H, $\delta_A=3.53$, $\delta_B=3.43$, $^2J_{AB}=16.5$, $^3J_{AM}=4.3$, $^3J_{BM}=9.5$ Hz, 2H; CH₂), 1.63 (appearance of sext, $^3J_{HH}=7.0$ Hz, 2H; CH₂CH₃), 1.47-1.38 (m, 2H; CH₂CH₃), 0.90 (t, $^3J_{HH}=7.3$ Hz, 3H; CH₃), 0.77 (t, $^3J_{HH}=7.3$ Hz, 3H; CH₃); ¹³C {¹H} NMR (150.9 MHz, CDCl₃, TMS): $\delta=196.1$, 168.5, 167.8, 165.7 (d, $J_{CF}=254.5$ Hz), 140.4, 133.3, 130.8 (d, $J_{CF}=9.2$ Hz), 128.5, 128.2, 127.2, 115.6 (d, $J_{CF}=21.9$ Hz), 67.2, 67.0, 57.6, 42.5, 40.9, 21.8, 21.7, 10.3, 10.2; HPLC (Chiralcel OJ-H, hexane/i-propanol=19/1, flow rate 0.5 mL/min, $\lambda=254$ nm): $t_{major}=41.0$ min, $t_{minor}=56.7$ min, ee=99%; $[\alpha]_D^{22}=+21.69$ (c=1.0 in CHCl₃); ESI-HRMS (m/z) calcd. for C₂₄H₂₇FO₅Na ((M+Na)⁺): 437.1740, found: 437.1729.


3cp

**[0134]** Dipropyl

2-(3-(4-methoxyphenyl)-3-oxo-1-phenylpropyl)malonate (table 3, entry 15)

[0135] It was synthesized in accordance with the above-mentioned manipulation.

[0136] Yield 85%, White solid, Mp 63-65 $^{\circ}$ C.; IR [cm⁻¹] (neat): 1731, 1677, 1602, 1265, 1171, 739, 702; ¹H NMR (600.2 MHz, CDCl₃, TMS): $\delta=7.89$ -7.86 (m, 2H; Ar), 7.27-7.20 (m, 4H; Ar), 7.16-7.12 (m, 1H; Ar), 6.89-6.86 (m, 2H; Ar), 4.17 (t d, $^3J_{HH}=4.6$, $^3J_{HH}=9.5$ Hz, 1H; CH), 4.14-4.05 (m, 2H; OCH₂), 3.87-3.83 (m, 3H; CH, OCH₂), 3.82 (s, 3H; OCH₃), ABM spin system (A=B=M=H, $\delta_A=3.45$, $\delta_B=3.40$, $^2J_{AB}=16.3$, $^3J_{AM}=4.5$, $^3J_{BM}=9.4$ Hz, 2H; CH₂), 1.63 (appearance of sext, $^3J_{HH}=7.1$ Hz, 2H; CH₂CH₃), 1.47-1.37 (m, 2H; CH₂CH₃), 0.90 (t, $^3J_{HH}=7.5$ Hz, 3H; CH₃), 0.77 (t, $^3J_{HH}=7.4$ Hz, 3H; CH₃); ¹³C {¹H} NMR (150.9 MHz, CDCl₃, TMS): $\delta=196.1$, 168.5, 167.9, 163.4, 140.6, 130.4, 130.0, 128.4, 128.2, 127.1, 113.7, 67.2, 66.9, 57.7, 55.4, 42.3, 41.0, 21.8, 21.6, 10.3, 10.2; HPLC (Chiralcel OD-H, hexane/i-propanol=19/1, flow rate 0.5 mL/min, $\lambda=254$ nm): $t_{major}=40.4$ min, $t_{minor}=47.3$ min, ee=99%; $[\alpha]_D^{20}=+20.73$ (c=1.0 in CHCl₃); ESI-HRMS (m/z) calcd. for C₂₅H₃₀O₆Na ((M+Na)⁺): 449.1940, found: 449.1944.




3cq

[0137] Dipropyl

2-(3-oxo-1,3-di(thiophen-2-yl)propyl)malonate
(table 3, entry 16)

[0138] It was synthesized in accordance with the above-mentioned manipulation.

[0139] Yield 73%, Beige liquid; IR [cm^{-1}] (neat): 1734, 1663, 1416, 1265, 736, 704; ^1H NMR (600.2 MHz, CDCl_3 , TMS): $\delta=7.75\text{--}7.73$ (m, 1H; Ar), 7.61-7.58 (m, 1H; Ar), 7.11-7.08 (m, 2H; Ar), 6.93-6.91 (m, 1H; Ar), 6.86-6.83 (m, 1H; Ar), 4.53-4.48 (m, 1H; CH), 4.14-4.05 (m, 2H; OCH_2), 3.99-3.93 (m, 2H; OCH_2), 3.90 (d, $^3J_{\text{HH}}=8.4$ Hz, 1H; CH), 3.48 (d, $^3J_{\text{HH}}=6.8$ Hz, 2H; CH_2), 1.64 (appearance of sext, $^3J_{\text{HH}}=7.1$ Hz, 2H; CH_2CH_3), 1.56-1.48 (m, 2H; CH_2CH_3), 0.91 (t, $^3J_{\text{HH}}=7.3$ Hz, 3H; CH_3), 0.84 (t, $^3J_{\text{HH}}=7.3$ Hz, 3H; CH_3); ^{13}C { ^1H } NMR (150.9 MHz, CDCl_3 , TMS): $\delta=190.1$, 168.1, 167.7, 144.1, 143.3, 133.8, 132.1, 128.1, 126.6, 125.9, 124.2, 67.3, 67.2, 57.8, 43.8, 36.3, 21.8, 21.7, 10.3, 10.2; HPLC (Chiralpak AS-H, hexane/*i*-propanol=9/1, flow rate 1.0 mL/min, $\lambda=254$ nm): $t_{\text{major}}=12.7$ min, $t_{\text{minor}}=17.3$ min, ee=97%; $[\alpha]_D^{21}=+22.21$ (c=1.0 in CHCl_3); ESI-HRMS (m/z) calcd. for $\text{C}_{20}\text{H}_{24}\text{O}_5\text{S}_2\text{Na}$ ((M+Na) $^+$): 431.0963, found: 431.0922.



3cr


[0140] Dipropyl

2-(1-(5-methylfuran-2-yl)-3-oxo-3-phenylpropyl)
malonate (table 3, entry 17)

[0141] It was synthesized in accordance with the above-mentioned manipulation.

[0142] Yield 71%, Yellow liquid; IR [cm^{-1}] (neat): 1735, 1686, 1265, 750; ^1H NMR (600.2 MHz, CDCl_3 , TMS): $\delta=7.95$ (br d, $J_{\text{HH}}=7.4$ Hz, 2H; Ar), 7.58-7.51 (m, 1H; Ar), 7.47-7.41 (m, 2H; Ar), 5.95 (br s, 1H; Ar), 5.76 (br s, 1H; Ar), 4.29-4.23 (m, 1H; CH), 4.14-3.90 (m, 5H; CH, OCH_2 , OCH_2), 3.58-3.41 (m, 2H; CH_2), 2.17 (s, 3H; CH_3), 1.68-1.53 (m, 4H; CH_2CH_3 , CH_2CH_3), 0.95-0.85 (m, 6H; CH_3 , CH_3); ^{13}C { ^1H } NMR (150.9 MHz, CDCl_3 , TMS): $\delta=197.5$, 168.2, 168.0, 151.6, 151.0, 136.8, 133.1, 128.5, 128.1, 107.7, 106.1,

67.1, 67.1, 55.3, 39.8, 34.4, 21.8, 21.8, 13.5, 10.3, 10.3; HPLC (Chiralpak AD-H, hexane/*i*-propanol=19/1, flow rate 0.5 mL/min, $\lambda=254$ nm): $t_{\text{major}}=40.8$ min, $t_{\text{minor}}=37.2$ min, ee=96%; $[\alpha]_D^{21}=+9.82$ (c=1.0 in CHCl_3); ESI-HRMS (m/z) calcd. for $\text{C}_{23}\text{H}_{28}\text{O}_6\text{Na}$ ((M+Na) $^+$): 423.1784, found: 423.1777.




3cs

[0143] Dipropyl

2-(3-oxo-1-phenyl-3-(1H-pyrrol-1-yl)propyl)malonate (table 3, entry 18)

[0144] It was synthesized in accordance with the above-mentioned manipulation.

[0145] Yield 93%, White solid, Mp 88-91° C.; IR [cm^{-1}] (KBr): 1716, 1471, 1280, 1229, 1172, 748; ^1H NMR (600.2 MHz, CDCl_3 , TMS): $\delta=7.30\text{--}7.25$ (m, 6H; Ar), 7.22-7.18 (m, 1H; Ar), 6.26-6.23 (m, 2H; Ar), 4.16-4.06 (m, 3H; CH, OCH_2), 3.88-3.84 (m, 3H; CH, OCH_2), ABM spin system ($A=B=M=H$, $\delta_A=3.45$, $\delta_B=3.29$, $^2J_{AB}=16.3$, $^3J_{AM}=4.2$, $^3J_{BM}=9.7$ Hz, 2H; CH_2), 1.64 (appearance of sext, $^3J_{\text{HH}}=7.2$ Hz, 2H; CH_2CH_3), 1.48-1.39 (m, 2H; CH_2CH_3), 0.90 (t, $^3J_{\text{HH}}=7.5$ Hz, 3H; CH_3), 0.78 (t, $^3J_{\text{HH}}=7.5$ Hz, 3H; CH_3); ^{13}C NMR (150.9 MHz, CDCl_3 , TMS): $\delta=168.3$, 168.2, 167.6, 139.6, 128.6, 128.1, 127.5, 119.1, 113.1, 67.3, 67.1, 57.2, 41.1, 38.7, 21.8, 21.6, 10.2, 10.2; HPLC (Chiralpak AS-H, hexane/*i*-propanol=19/1, flow rate 0.5 mL/min, $\lambda=254$ nm): $t_{\text{major}}=24.4$ min, $t_{\text{minor}}=31.2$ min, ee=99%; $[\alpha]_D^{20}=+16.38$ (c=1.0 in CHCl_3); ESI-HRMS (m/z) calcd. for $\text{C}_{22}\text{H}_{27}\text{NO}_5\text{Na}$ ((M+Na) $^+$): 408.1787, found: 408.1755.



3ct


[0146] Dipropyl

2-(3-oxo-1,5-diphenylpent-4-enyl)malonate (table 3, entry 19)

[0147] It was synthesized in accordance with the above-mentioned manipulation.

[0148] Yield 75%, White solid, Mp 69-73° C.; IR [cm^{-1}] (KBr): 1731, 1646, 1227, 1163, 705; ^1H NMR (600.2 MHz, CDCl_3 , TMS): $\delta=7.51\text{--}7.46$ (m, 3H; Ar), 7.37-7.34 (m, 3H; Ar), 7.29-7.23 (m, 4H; Ar), 7.19-7.14 (m, 1H; Ar), 7.14 (d,

$^3J_{HH}=16.4$, 1H; Ar), 4.16-4.07 (m, 3H; CH, OCH₂), 3.87-3.81 (m, 3H; CH, OCH₂), ABM spin system (A=B=M=H, $\delta_A=3.20$, $\delta_B=3.16$, $^2J_{AB}=16.1$, $^3J_{AM}=4.6$, $^3J_{BM}=9.2$ Hz, 2H; CH₂), 1.65 (appearance of sext, $^3J_{HH}=7.1$ Hz, 2H; CH₂CH₃), 1.48-1.38 (m, 2H; CH₂CH₃), 0.92 (t, $^3J_{HH}=7.5$ Hz, 3H; CH₃), 0.77 (t, $^3J_{HH}=7.3$ Hz, 3H; CH₃); ^{13}C { ^1H } NMR (150.9 MHz, CDCl₃, TMS): $\delta=197.4$, 168.4, 167.8, 142.8, 140.4, 134.5, 130.4, 128.9, 128.4, 128.3, 128.2, 127.2, 126.0, 67.2, 66.9, 57.6, 44.8, 41.0, 21.9, 21.6, 10.3, 10.2; HPLC (Chiralpak AS-H, hexane/*i*-propanol=9/1, flow rate 1.0 mL/min, $\lambda=254$ nm): $t_{major}=11.8$ min, $t_{minor}=14.4$ min, ee=86%; $[\alpha]_D^{22}=+14.56$ (c=1.0 in CHCl₃); ESI-HRMS (m/z) calcd. for C₂₆H₃₀O₅Na ((M+Na)⁺): 445.1991, found: 445.1975.



3cu

[0149] Dipropyl

2-(5-oxo-1,5-diphenylpent-1-en-3-yl)malonate (table 3, entry 20)

[0150] It was synthesized in accordance with the above-mentioned manipulation.


[0151] Yield 46%, White solid, Mp 42-45° C.; IR [cm⁻¹] (KBr): 1728, 1682, 1234, 754, 692; ^1H NMR (600.2 MHz, CDCl₃, TMS): $\delta=7.98$ -7.94 (m, 2H; Ar), 7.56-7.52 (m, 1H; Ar), 7.47-7.43 (m, 2H; Ar), 7.29-7.22 (m, 4H; Ar), 7.20-7.16 (m, 1H; Ar), 6.46 (d, $^3J_{HH}=15.8$ Hz, 1H; Ar), 6.25 (dd, $^3J_{HH}=15.8$, $^3J_{HH}=9.0$ Hz, 1H; Ar), 4.15-4.00 (m, 4H; OCH₂, OCH₂), 3.78 (d, $^3J_{HH}=7.5$ Hz, 1H; CH), 3.73-3.67 (m, 1H; CH), ABM spin system (A=B=M=H, $\delta_A=3.40$, $\delta_B=3.27$, $^2J_{AB}=16.8$, $^3J_{AM}=4.9$, $^3J_{BM}=7.9$ Hz, 2H; CH₂), 1.68-1.55 (m, 4H; CH₂CH₃, CH₂CH₃), 0.92 (t, $^3J_{HH}=7.5$ Hz, 3H; CH₃), 0.87 (t, $^3J_{HH}=7.5$ Hz, 3H; CH₃); ^{13}C { ^1H } NMR (124.5 MHz, CDCl₃, TMS): $\delta=198.0$, 168.4, 168.3, 137.0, 136.9, 133.1, 132.6, 128.6, 128.5, 128.4, 128.2, 127.5, 126.4, 67.1, 67.0, 55.7, 41.3, 38.8, 21.9, 21.8, 10.3, 10.3; HPLC (Chiralpak AS-H, hexane/*i*-propanol=40/1, flow rate 0.5 mL/min, $\lambda=254$ nm): $t_{major}=32.5$ min, $t_{minor}=38.0$ min, ee=97%; $[\alpha]_D^{21}=+1.20$ (c=1.0 in CHCl₃); ESI-HRMS (m/z) calcd. for C₂₆H₃₀O₅Na ((M+Na)⁺): 445.1991, found: 445.1988, C₂₆H₃₁O₅ ((M+H)⁺): 423.2171, found: 423.2150.

TABLE 1


Effect of metal sources and chiral ligands.^a

Reaction scheme: 1b (diethyl malonate) + 2a (chalcone) $\xrightarrow[\text{Solv., 0.1 M}]{\text{Metal (x mol \%), Ligand (1.2 x mol \%), MS 4A (100 mg)}}$ 3ba (chiral product)

entry	metal (xmol %)	ligand	solv.	time (h)	yield (%) ^b	ee (%) ^c
1 ^d	Ca(O- <i>i</i> -Pr) ₂ (10%)	I	THF	24	47	4
2 ^d	Ca(O- <i>i</i> -Pr) ₂ (10%)	II	THF	24	47	49
3 ^d	Ca(O- <i>i</i> -Pr) ₂ (10%)	III	THF	24	89	52
4 ^e	Ca(O- <i>i</i> -Pr) ₂ (5%)	III	Tol.	18	58	65
5 ^e	Sr(O- <i>i</i> -Pr) ₂ (5%)	III	Tol.	18	91	97
6 ^e	Ba(O- <i>i</i> -Pr) ₂ (5%)	III	Tol.	18	80	76
7 ^e	Ba(O- <i>t</i> -Bu) ₂ (5%)	III	Tol.	18	82	70



I



II R = 4-tolyl
III R = 2,5-dimethylbenzene

TABLE 2

Conjugate addition reactions of malonates 1a-f to 2a.^a

Reaction scheme showing the conjugate addition of malonates 1a-f to enone 2a. The reaction conditions are Sr(O-i-Pr)₂ (x mol %), Ligand III (1.2 equiv), MS 4A (100 mg), 25° C., Toluene, 0.1 M. The product is 3aa-fa.

entry	Sr(O-i-Pr) ₂ (xmol %)	R	time (h)	yield (%) ^b	ee (%) ^c (configuration)
1	5	Me	24	65	94 (R) ^d
2	5	Et	18	91	97 (R) ^d
3	5	n-Pr	17	92	99
4	2.5	n-Pr	7	90	99
5 ^e	1	n-Pr	9	70	97
6 ^f	0.5	n-Pr	24	72	97
7	5	i-Pr	21	83	89
8	5	n-Bu	3	85	96
9	5	Bn	18	85	84

TABLE 3

Conjugate addition reactions of 1c to enones 2b-u.^a

Reaction scheme showing the conjugate addition of malonate 1c to enones 2b-u. The reaction conditions are Sr(O-i-Pr)₂ (5 mol %), Ligand III (6 mol %), MS 4A (100 mg), 7 h, 25° C., Toluene, 0.1 M. The product is 3cb-cu.

entry	R ¹	R ²	adduct	yield (%) ^b	ee (%) ^c
1	2-ClC ₆ H ₄	Ph	3cb	76	92
2	4-ClC ₆ H ₄	Ph	3cc	93	97
3	4-FC ₆ H ₄	Ph	3cd	92	98
4	4-MeOC ₆ H ₄	Ph	3ce	80	>99
5	4-NO ₂ C ₆ H ₄	Ph	3cf	98	96
6	3-NO ₂ C ₆ H ₄	Ph	3cg	94	94
7	4-FC ₆ H ₄	4-FC ₆ H ₄	3ch	91	96
8	4-MeOC ₆ H ₄	4-FC ₆ H ₄	3ci	81	>99
9	3,4-di-MeOC ₆ H ₃	4-FC ₆ H ₄	3cj	61	96
10	4-ClC ₆ H ₄	4-FC ₆ H ₄	3ck	97	97
11	2-ClC ₆ H ₄	4-FC ₆ H ₄	3cl	80	93
12	4-FC ₆ H ₄	4-MeC ₆ H ₄	3cm	90	98
13	Ph	4-ClC ₆ H ₄	3cn	98	99
14	Ph	4-FC ₆ H ₄	3co	92	99
15	Ph	4-MeOC ₆ H ₄	3cp	85	99
16	2-thienyl	2-thienyl	3cq	73	97
17 ^d	5-methylfuran-2-yl	Ph	3cr	71	96
18 ^e	Ph	1-pyrrolyl	3cs	90	>99
19 ^f	Ph	—CH=CHPh	3ct	97	86
20 ^f	—CH=CH—Ph	Ph	3cu	62	97


^aSee footnote in Table 1.^bIsolated yields.^cDetermined by chiral HPLC analysis.^dReaction time 48 h.^eReaction time 24 h.^f2.2 equivalents of malonate 1c were used.

[0152] [Michael Addition to Chalcone of a Malonic Ester]

[0153] 0.015 mmol of strontium hexamethyldisilazide (Sr(HMDS)₂), 0.015 mmol of ligand, and 100 mg of molecular sieve 4A were suspended in 1 mL of toluene, and stirred for two hours at room temperature under an argon atmosphere. Thereafter, a toluene solution (1 mL) of 0.36 mmol of a malonic acid di-n-propyl ester, and a toluene solution of chalcone (0.3 mmol) were added. After the finishing of the reaction (the finishing of the reaction was confirmed by the thin-layer chromatography), a saturated ammonium chloride aqueous solution was added to the reaction solution. In addition, an organic phase was separated with dichloromethane. And a water phase was extracted with dichloromethane. The organic phase was collected and dried over anhydrous sodium sulfate. Thereafter, sodium sulfate was filtered off, and the solvent was removed by the distillation under reduced pressure. And the crude refined product was refined with the preparative thin-layer chromatography.

[0154] An enantiometric excess of the target product (refined product) obtained in such a manner was determined with a high performance liquid chromatography.

[0155] Additionally, Sr(HMDS)₂ was synthesized with the method (Inorg. Chem., 1991, 30, 96-101) reported by Wasterhausen. Ligand was synthesized with method (J. Am. Chem. Soc., 1997, 119, 6452-6453) reported by Evans. Chalcones were procured from TOKYO CHEMICAL INDUSTRY CO., LTD. and Wako Pure Chemical industries, LTD.



[0156] Dipropyl

[0157] 2-((1-oxo-1,2,3,4-tetrahydronaphthalen-2-yl)(phenyl)methyl)malonate (table 5): Colorless oil; IR [cm⁻¹] (neat): 1747, 1731, 1682, 1600, 1455, 1266, 1221, 1155, 1059, 743, 702; NMR (600.2 MHz, CDCl₃, TMS): δ=8.01-7.97 (m, 1H; Ar), 7.41-7.37 (m, 1H; Ar), 7.32-7.11 (m, 7H; Ar), 4.98 (d, ³J_{HH}=12.0 Hz, 1H; CH^c), 4.14-4.00 (m, 2H; OCH₂), 3.85-3.75 (m, 3H; CH^d, OCH₂), 3.11 (dt, ³J_{HH}=12.9, ³J_{HH}=4.1 Hz, 1H; CH^c), 2.99-2.92 (m, 1H; CH^a), 2.86-2.81 (m, 1H; CH^a), 2.15-2.10 (m, 1H; CH^b), 1.86 (appearance of d, ³J_{HH}=12.9, J_{HH}=4.2 Hz, 1H; CH^b), 1.60 (appearance of sext, ³J_{HH}=7.1 Hz, 2H; CH₂CH₃), 1.40-1.32 (m, 2H; CH₂CH₃), 0.89 (t, ³J_{HH}=7.3 Hz, 3H; CH₃), 0.74 (t, ³J_{HH}=7.5 Hz, 3H; CH₃); ¹³C {¹H} NMR (150.9 MHz, CDCl₃, TMS): δ=198.8, 169.2, 168.7, 143.6, 139.3, 133.3, 133.2, 129.5, 128.4, 128.2, 127.3, 127.0, 126.5, 67.0, 66.7, 54.9, 49.6, 47.7, 29.5, 27.8, 21.8, 21.6, 10.2, 10.2; HPLC (Chiralpak AS-H, hexane/i-propanol=40/1, flow rate 0.5 mL/min, λ=254 nm): t_{major}=16.5 min, t_{minor}=19.6 min, ee=96%, t_{major}=23.3 min, t_{minor}=26.3 min; [α]_D²¹=59.52 (c=1.0 in CHCl₃); ESI-HRMS (m/z) calcd. for C₂₆H₃₀O₅Na [(M+Na)⁺]: 445.1991, found: 445.2042.

TABLE 4

Strontiums	Catalyst amount (mole %)	Yield (%) ^[a]	Enantiometric excess (%) ^[b]
1 Sr(O-i-Pr) ₂	5	92	99
2 Sr(HMDS) ₂	5	97	99
3 Sr(HMDS) ₂	3	99	96
4 Sr(HMDS) ₂	2	96	96

^[a]Isolated yield

^[b]Determined with chiral HPLC analysis

TABLE 5

Strontiums	Compound 1 (y equivalents)	Yield (%) ^[a]	Diastereomer ratio	Enantiometric excess (%) ^[b]
1 Sr(O-i-Pr) ₂	2.5	50	97:3	80
2 Sr(O-i-Pr) ₂	5	92	97:3	66
3 Sr(O-i-Pr) ₂	1.2	40	97:3	88
4 Sr(HMDS) ₂	1.2	38	97:3	95
5 Sr(HMDS) ₂	1.2	26	98:2	95
6 Sr(HMDS) ₂	2.5	50	97:3	94
7 Sr(HMDS) ₂	1.2	86	98:2	60


^[a]Isolated yield^[b]Determined with chiral HPLC analysis

TABLE 6

Entry	Substituent R ¹	Substituent R ²	Reaction time (time)	Yield (%) ^[a]	Enantioselectivity (%) ^[b]
1	2-Cl-C ₆ H ₄	Ph	7	82	95
2	4-Cl-C ₆ H ₄	Ph	7	93	97
3	2-Cl-C ₆ H ₄	4-F-C ₆ H ₄	7	92	93
4	Ph	-CH=CHPh	7	95	90
5	4-F-C ₆ H ₄	4-F-C ₆ H ₄	7	94	96
6	3-NO ₂ -C ₆ H ₄	Ph	7	86	96
7	4-NO ₂ -C ₆ H ₄	Ph	7	81	97

^[a]Isolated yield^[b]Determined with chiral HPLC analysis

1. A reaction method, comprising reacting a compound represented by the following general formula [II] with a compound represented by the following general formula [III] in the presence of a catalyst configured using MX₂, wherein M is Be, Mg, Ca, Sr, Ba or Ra, and X is an arbitrary group, and a compound represented by the following general formula [I]



general formula


wherein R⁷, R⁸, R⁹, and R¹⁰ each represents a substituted cyclic group or an unsubstituted cyclic group and wherein optionally R⁹ and R¹⁰ form a ring



general formula [III]

-continued

general formula [III]



Wherein each of R¹, R², R³, and R⁴ is an arbitrary substituent, and each of W¹, W², and W³ is an electron-withdrawing group.

2. A reaction method according to claim 1, wherein said MX₂ is M(OR⁵)₂ wherein M is Mg, Ca, Sr or Ba and R⁵ is an alkyl group.

3. A reaction method according to claim 1, wherein said MX₂ is M(OR⁵)₂ and M is Ca, Sr or Ba and R⁵ is an alkyl group having a carbon number of 1 to 10.

4. A reaction method according to claim 1, wherein said MX₂ is Sr(OR⁵)₂ and R⁵ is an alkyl group having a carbon number of 1 to 10.

5. A reaction method according to claim 1, wherein said X is an amide group.

6. A reaction method according to claim 1, wherein said X is hexamethyldisilazide.

7. A reaction method according to claim 1, wherein said cyclic group is an aromatic group.

8. A reaction method according to claim 1, wherein the compound represented by said general formula [I] and M of said compound MX_2 are coordinate-bonded to each other.

9. A reaction method according to claim 1, wherein each of said R^1 , R^2 , R^3 , and R^4 is an H group or a hydrocarbon group.

10. A reaction method according to claim 1, wherein said electron-withdrawing group is an ester group or a carbonyl group.


11. A reaction method according to claim 1, wherein the compound represented by said general formula [II] is a dicarboxylate ester.

12. A reaction method according to claim 1, wherein the compound represented by said general formula [III] is a malonic ester.

13. A reaction method according to claim 1, wherein the compound represented by said general formula [III] is an enone.


14. A reaction method according to claim 1, wherein an aromatic hydrocarbon solvent is used as a solvent of said reaction.

15. A reaction method according to claim 1, said reaction method used for obtaining a compound represented by the following general formula [IV]



general formula [IV]

16. A catalyst for a reaction between a compound represented by the following general formula [II] and a compound represented by the following general formula [III], said catalyst configured using MX_2 , wherein M is Be, Mg, Ca, Sr, Ba or Ra, and X is an arbitrary group, and a compound represented by the following general formula [I]



general formula [I]


herein R^7 , R^8 , R^9 , and R^{10} each represents a substituted cyclic group or an unsubstituted cyclic group, and optionally R^9 and R^{10} form a ring,



general formula [II]


-continued

general formula [III]



wherein each of R^1 , R^2 , R^3 , and R^4 is an arbitrary substituent, and each of W^1 , W^2 , and W^3 is an electron-withdrawing group.

17. A catalyst configured using MX_2 , wherein M is Ca, Sr, Ba or Ra, and X is an arbitrary group, and a compound represented by the following general formula [I]



general formula [I]

wherein R^7 , R^8 , R^9 , and R^{10} each represents a substituted cyclic group or an unsubstituted cyclic group, and optionally R^9 and R^{10} form a ring.

18. A catalyst according to claim 16, wherein said MX_2 is $M(OR^5)_2$, wherein M is Ca, Sr or Ba, and R^5 is an alkyl group.

19. A catalyst according to claim 16, wherein said MX_2 is $M(OR^5)_2$, wherein M is Ca, Sr or Ba, and R^5 is an alkyl group having a carbon number of 1 to 10.

20. A catalyst according to claim 16, wherein said MX_2 is $Sr(OR^5)_2$, wherein R^5 is an alkyl group having a carbon number of 1 to 10.

21. A catalyst according to claim 16, wherein said X is an amide group.

22. A catalyst according to claim 16, wherein said X is hexamethyldisilazide.

23. A catalyst according to claim 16, wherein said cyclic group is an aromatic group.

24. A catalyst according to claim 16, wherein the compound represented by said general formula [I] and M of said compound MX_2 are coordinate-bonded to each other.

25. A catalyst according to claim 16, wherein each of said R^1 , R^2 , R^3 , and R^4 is an H group or a hydrocarbon group.

26. A catalyst according to claim 16, wherein said electron-withdrawing group is an ester group or a carbonyl group.

27. A catalyst according to claim 16, wherein the compound represented by said general formula [II] is a dicarboxylate ester.

28. A catalyst according to claim 16, wherein the compound represented by said general formula [II] is a malonic ester.

29. A catalyst according to claim 16, wherein the compound represented by said general formula [III] is an enone.

30. A catalyst according to claim 17, wherein said MX_2 is $M(OR^5)_2$, wherein M is Ca, Sr or Ba, and R^5 is an alkyl group.

31. A catalyst according to claim 17, wherein said MX_2 is $M(OR^5)_2$, wherein M is Ca, Sr or Ba, and R^5 is an alkyl group having a carbon number of 1 to 10.

32. A catalyst according to claim 17, wherein said MX_2 is $\text{Sr}(\text{OR}^5)_2$, and R^5 is an alkyl group having a carbon number of 1 to 10.

33. A catalyst according to claim 17, wherein said X is an amide group.

34. A catalyst according to claim 17, wherein said X is hexamethyldisilazide.

35. A catalyst according to claim 17, wherein said cyclic group is an aromatic group.

36. A catalyst according to claim 17, wherein the compound represented by said general formula [I] and M of said compound MX_2 are coordinate-bonded to each other.

* * * * *