



(11) **EP 1 323 440 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

- (45) Date of publication and mention of the grant of the patent: **09.05.2012 Bulletin 2012/19**
- (21) Application number: **01963506.9**
- (22) Date of filing: **07.09.2001**
- (51) Int Cl.: **A61M 1/28** (2006.01) **A61K 9/08** (2006.01)
A61K 47/20 (2006.01)
- (86) International application number: **PCT/JP2001/007772**
- (87) International publication number: **WO 2002/022188 (21.03.2002 Gazette 2002/12)**

(54) **PERITONEAL DIALYSATES**

PERITONEAL-DIALYSATE

DIALYSATS PERITONEAUX

- (84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE TR
- (30) Priority: **13.09.2000 JP 2000277810**
16.02.2001 JP 2001040718
20.06.2001 JP 2001186642
- (43) Date of publication of application:
02.07.2003 Bulletin 2003/27
- (73) Proprietor: **Japan Science and Technology Agency**
Kawaguchi-shi
Saitama (JP)
- (72) Inventors:
• **SAKAI, Asahi**
Sakura-shi,
Chiba 285-0867 (JP)
• **NAKAYAMA, Masaaki**
Tokyo 150-0013 (JP)
- (74) Representative: **Waldren, Robin Michael et al**
Marks & Clerk LLP
90 Long Acre
London
WC2E 9RA (GB)
- (56) References cited:
WO-A-01/89478 WO-A-02/053094
WO-A1-00/10606 WO-A1-00/20052
JP-A- 7 136 255 JP-A- 8 337 590
JP-B2- 5 058 604
- **HALABY S F ET AL: "ABSORPTION OF SODIUM BISULFITE FROM PERITONEAL DIALYSIS SOLUTIONS" JOURNAL OF PHARMACEUTICAL SCIENCES, AMERICAN PHARMACEUTICAL ASSOCIATION. WASHINGTON, US, vol. 54, no. 1, January 1965 (1965-01), pages 52-55, XP002946328 ISSN: 0022-3549**
 - **LANG K ET AL: "SULFIT IN INFUSIONSLOESUNGEN" ZEITSCHRIFT FUER ERNAHRUNGSWISSENSCHAFT, STEINKOPF VERLAG, DARMSTADT, DE, vol. 18, no. 1, 1979, pages 37-41, XP002946330 ISSN: 0044-264X**
 - **HOWELL S B ET AL: "INTRAPERITONEAL CIS-DIAMMINIDICHLOROPLATINUM WITH SYSTEMIC THIOSULFATE PROTECTION" CANCER RESEARCH, AMERICAN ASSOCIATION FOR CANCER RESEARCH, BALTIMORE, MD, US, vol. 43, March 1983 (1983-03), pages 1426-1431, XP002946327 ISSN: 0008-5472**
 - **WIESLANDER A P ET AL: "HEAT STERILIZATION OF GLUCOSE-CONTAINING FLUIDS FOR PERITONEAL DIALYSIS: BIOLOGICAL CONSEQUENCES OF CHEMICAL ALTERATIONS" PERITONEAL DIALYSIS INTERNATIONAL, PERGAMON PRESS, NEW YORK, NY, US, vol. 15, no. 7, SUPPL, 1995, pages S52-S60, XP002946331 ISSN: 0896-8608**
 - **FUMINORI KATO ET AL: "Immunosuppressive effects of 3,4-dideoxyglucosone-3-ene, an intermediate in the maillard reaction" JOURNAL OF AGRICULTURAL AND FOOD CHEMISTRY, AMERICAN CHEMICAL SOCIETY. WASHINGTON, US, vol. 42, no. 9, 1994, pages 2068-2073, XP002966230 ISSN: 0021-8561**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 323 440 B1

- **TORBJORN LINDEN ET AL: "3-Deoxyglucosone, a promoter of advanced glycation end products in fluids for peritoneal dialysis" PERITONEAL DIALYSIS INTERNATIONAL, PERGAMON PRESS, NEW YORK, NY, US, vol. 18, no. 3, May 1998 (1998-05), pages 290-293, XP002966246 ISSN: 0896-8608**

Remarks:

The file contains technical information submitted after the application was filed and not included in this specification

Description

FIELD OF THE INVENTION

5 **[0001]** The present invention relates to the dialysate for peritoneal dialysis, that is, therapy for end stage renal disease, more specifically, the dialysate for peritoneal dialysis to suppress protein cross linking due to sugar osmotic agents, such as glucose and the like.


BACKGROUND OF THE INVENTION

10 **[0002]** The peritoneal dialysis has been applied as an effective therapy for end stage renal disease patients. The dialysis is proceeded by infusing dialysate into peritoneal cavity through a catheter, which is implanted in the patient's peritoneal cavity, and storing it for a certain period, thence withdrawing the dialysate out through the catheter. This procedure is repeated a few times every day.

15 **[0003]** This dialysis has a few advantages over hemodialysis in physiological point of view, as it purifies blood continuously through the patient's peritoneum, while hemodialysis purifies blood through an artificial membrane intermittently. Also peritoneal dialysis enables the patients' social activity, so that the peritoneal dialysis has been widely applied.

20 **[0004]** In hemodialysis, removal of excess liquid is achieved by raising the pressure of blood line over that of the dialysate line. However the same means can not be applied to peritoneal dialysis, therefore an osmotic agent is added into the dialysate so as to raise the osmotic pressure of the dialysate over that of plasma. The dialysate is infused into the peritoneal cavity to contact to peritoneum for removing excess liquid from the patient's body. For this purpose, glucose has been used as an osmotic agent. Glucose had been recognized to be safe and physiological, and to cause no problem on metabolism after it is absorbed into the body. However, adverse effects were recognized as serious problems, such as disfunctioning of peritoneum, due to the absorption of large quantity of glucose into the patient body and the reaction with amino acids, peptide and protein, followed by the formation of AGEs, progress of collagen synthesis, and cross linking of protein. Consequently it causes peritoneum sclerosis and leads to cease of the therapy.

25 **[0005]** The cross linking reaction of protein molecules with glucose is assumed to take place as follows;




Carbonyl residue(s) containing sugars, such as glucose and the like, react with amino acids, peptides or protein(II), and through Schiff base (III) and Amadori compounds(IV), lead to advanced glycation end products (AGE) ; the cross linking between protein molecules

[0006] Also it has been reported that glucose is converted through dialysate sterilization process under high pressure and temperature to the following compounds, that is called glucose degradation products (GDP); glyoxal, methylglyoxal, 3-deoxyglucosone.

[0007] The above-described GDPs are more reactive substances for AGE formation compared to sugars per se. When dialysate contains the GDPs, the protein cross linking reaction is accelerated by a few dozen to a few thousands times as fast as when the dialysate contains glucose alone.

[0008] An example of protein cross linkage formed in AGEs by the reaction between carbonyl residues of sugar or GDP and amino residues of lysine or arginine that constitutes protein molecules is shown as formula(2).



[0009] As one of the means for the solution of this problem, the modification of heat sterilization conditions for the dialysate has been proposed, but it can not prohibit the protein cross linking completely, and cross-linking reaction of protein by glucose itself may not be disregarded. So that effective prohibition means of protein cross linking are required for peritoneal dialysis.

[0010] The inventors of the present invention has proposed the technology wherein the patient's plasma protein, which is migrated out to dialysate, is recovered from peritoneal dialysis effluent and after the concentration, it is used as an osmotic agent for substituting a part or all of glucose. (Japanese Patent Application Hei 8-150930 and Hei 9-302388). However as long as glucose is used as a portion of osmotic agents, the problem of protein cross linking may not be solved completely.

[0011] Reference in this technical field include:

WO 01 89478 A (OLSSON LARS FRIDE ;WIESLANDER ANDERS (SE); GAMBRO LUNDIA AB (SE);) 29 November 2001 (2001-11-29)

WO 02 053094 A (DEPPISCH REINHOLD ;WIESLANDER ANDERS (SE); GAMBRO LUNDIA AB (SE);) 11 July 2002 (2002-07-11)

HALABY S F ET AL: 'ABSORPTION OF SODIUM BISULFITE FROM PERITONEAL DIALYSIS SOLUTIONS' JOURNAL OF PHARMACEUTICAL SCIENCES, AMERICAN PHARMACEUTICAL ASSOCIATION. WASHINGTON, US, vol. 54, no. 1, January 1965 (1965-01), pages 52-55, XP002946328 ISSN: 0022-3549

LANG K ET AL: 'SULFIT IN INFUSIONSLOESUNGEN' ZEITSCHRIFT FUER ERNAEHRUNGSWISSENSCHAFT, STEINKOPF VERLAG, DARMSTADT, DE, vol. 18, no. 1, 1979, pages 37-41, XP002946330 ISSN: 0044-264X

WIESLANDER A P ET AL: 'HEAT STERILIZATION OF GLUCOSE-CONTAINING FLUIDS FOR PERITONEAL DIALYSIS: BIOLOGICAL CONSEQUENCES OF CHEMICAL ALTERATIONS' PERITONEAL DIALYSIS INTERNATIONAL, PERGAMON PRESS, NEW YORK, NY, US, vol. 15, no. 7, SUPPL, 1995, pages S52-S60,

XP002946331 ISSN: 0896-8608

None of these documents describes a dialysate solution containing a salt of thiosulfate or dithionite.

5 HOWELL S B ET AL: 'INTRAPERITONEAL CIS-DIAMMINIDICHLOROPLATINIUM WITH SYSTEMIC THIOSULFATE PROTECTION' CANCER RESEARCH, AMERICAN ASSOCIATION FOR CANCER RESEARCH, BALTIMORE, MD, US, vol. 43, March 1983 (1983-03), pages 1426-1431, XP002946327 ISSN: 0008-5472,

mentions thiosulfate, but not as additive in a dialysate.

10

DESCRIPTION OF THE INVENTION

[0012] The inventors of the present invention understood protein cross linking with glucose or GDPs takes place through the afore-described mechanism and assumed that the protein cross linking reaction may be suppressed by addition of the inhibitors of the reaction, and searched a variety of chemical compounds which they assumed to be inhibitors. Resultantly, they found a number of effective compounds in suppression of the protein cross linking. Especially they found that physiologically safe salt of thiosulfate or dithionite reductive salts of sulfur oxy-acid are effective.

15

[0013] Also they found effective compounds that may split cross linkage of protein, even after cross linkage is formed.

20

[0014] The present invention is defined by the appended claims.

MOST PREFERRABLE EMBODIMENT FOR CARRYING OUT THE INVENTION

[0015] In the present invention, sugars are used as osmotic agents for the dialysate of peritoneal dialysis. The example of the sugars are monosaccharides such as glucose, mannose, and the like, disaccharides such as sucrose, fructose and the like, or oligomer and polymers such as dextran, dextrin and the like. Also the dialysate may contain amino acids, peptides and protein in addition to sugars.

25

[0016] The electrolytes may be mixture of sodium chloride, magnesium chlorides, calcium chloride, sodium lactate, sodium bicarbonate and the like. The composition and the concentration of the electrolytes are favorably close to those of the serum. When the concentration of calcium and magnesium in the patients serum differs from the ordinal value due to drug dosage for complication, the adjusted dialysate in calcium and magnesium concentration are preferably used.

30

[0017] In the present invention, the inhibitors of protein cross linking, as additives to the aqueous solution of electrolytes and sugars, may be used if they are physiologically safe and inhibit AGE reaction at any stage.

[0018] Also the splitters of protein cross linkage may be used if they split protein cross linkage under physiological conditions and their reaction products are not toxic.

35

[0019] Generally speaking, reductants may be effective suppresser of cross linking

[0020] The reductants having lower redox potential may be more effective suppresser. Especially those having lower redox potential compared to that of saline solution (+160mV~+180mV) are effective.

40

[0021] More specifically, examples of effective suppressers of cross linking, which fall into reductants, anti-oxidants or other compounds than the above described, may be mercaptan, sulfides, hydrosulfides, salts of reductive sulfur oxy-acid, thiourea and their derivatives, hydroxyl or carboxyl residue(s) containing, cyclic compounds, flavonoids, nitrogen containing hetero cyclic compounds, hydrazil compounds, uronic acid(s) containing mucopolysaccharides. While examples of splitters of protein cross linkage may be thiazole derivatives

[0022] Examples of mercaptan, i.e. those having mercapto (-SH) residue(s), may be cysteine, acetylcysteine, mercaptoethanol, glutathione, dithioerythrytol, N-acetylmercaptosuccinic anhydride and the like.

45

[0023] Examples of sulfides and hydrosulfides may be sodium sulfide, sodium hydrosulfide and the like.

[0024] Examples of salts of reductive sulfur oxy-acid may be sodium, potassium, or other physiologically safe salts of sulfite, bisulfite, thio-sulfate, metabisulfite (disulfite) or dithionite. These salts may be acidic salts (for example acid bisulfite) as well as neutral salts

[0025] Examples of thiourea and its derivatives are thiourea, dimethyl thiourea and the like.

50

[0026] Examples of hydroxyl and/or carboxyl residue(s) containing cyclic compounds may be acetyl salicylic acid, ascorbic acid, or their sodium salts or other physiologically safe salts.

[0027] Examples of flavonoids may be hetero-cyclic compounds which contain more than two hydroxyl residues, such as quercetin dihydrate, catechin, epicatechin, or their hydrates,

55

[0028] Examples of nitrogen containing hetero cyclic compounds may be thiazole, thiazoline, thiazolidine, triazole, tetrazole, indole, imidazole, pyridine, and pyrimidine cycles.

[0029] Examples of thiazole containing cyclic compounds may be N-(2-thiazolyl) sulfanilamide, N-phenacyl thiazolium bromide. The latter compound may reacts with the cross linked protein to split the linkage.

[0030] Example of indole containing cyclic compounds may be N-acetyltryptophan.

[0031] Example of triazol containing cyclic compounds may be 4-(1, 2, 3, 4-thiazolo-5-rylamino) phenol

[0032] Example of thiazoline containing cyclic compounds which comprise may be 2-mercaptothiazoline.

[0033] Example of thiazolidine containing cyclic compounds may be 2-Oxothiazolidine-4- carboxylic acid

[0034] Example of hydrazil compounds may be aminoguanidine hydrochloride.

[0035] Example of uronic acid(s) containing mucopolysaccharides may be heparin

[0036] The dose of the protein cross linking suppressors may vary in the range of 0.1 ~ 200weight %, preferably 1~100weight %. When the dose is too low, the suppression effect of protein cross linking would be insufficient, while the dose is too high, it may cause poor dissolving in dialysate.

[0037] The concentration of sugar osmotic agents may be adequate in the range of 5~300mOsm/l in addition to that of electrolytes. When the concentration of sugar osmotic agents is too low, adequate ultra filtration may not be achieved, while if it is too high, it causes adverse effects such as worsening of diabetes mellitus.

[0038] Additionally, oncotic agents other than sugar osmotic agents may be added into dialysate as much as they indicate significant oncotic pressure. Examples of the oncotic agents may be polymers such as albumin, globulin and the like.

[0039] These oncotic agents are not restricted to be chemical reagents nor medical intravenous solution, but a mixture of serum protein which is recovered from the patient's own effluent of peritoneal dialysis

[0040] The recovery method is described in the Japanese Patent Application number Hei 8-150930 and Hei 9-302388 filed by the inventor of the present invention, that is, concentrating the effluent through a semi permeable membrane and diluting the condensate with water or electrolyte solution, followed by repeating these procedures so as to refine the protein, or alternately adding acid into the concentrate, followed by de-acidification by membrane dialysis with water to deposit the protein at the iso-electric point, thence separating the deposit from supernatant and re-dissolving it into fresh dialysate as oncotic agent, The dialysate may be supplemented by additional oncotic agents too.

[0041] By substituting a part of sugar osmotic agents with non-sugar oncotic agents, dose of the sugar can be decreased, so that protein cross linking due to sugar can be suppressed. In the dialysate containing oncotic agents, the concentration of oncotic agents may be preferable in the range of 0.1~30g/dl.

[0042] The dialysate of peritoneal dialysis in the present invention is prepared by addition of a sodium, potassium or other physiologically safe salt of thiosulfate or dithionite, functioning as protein cross linking suppressor(s) and/or protein cross linkage splitter(s) into the solution of electrolytes and sugar osmotic agent. Conventional dialysate is sterilized by heating at 110°C or higher temperature. Also pH of the dialysate is lowered down to around 5.0-5.8.

[0043] When the cross linking suppressor(s) are added and the dialysate is heat sterilized, some instable suppressor (s) or the splitter(s) are decomposed. In such a case, it is preferable that the dialysate is sterilized separately, thence the suppressor(s) or the splitter(s) are added before use. If heat labile compound(s) are added into the dialysate, it is desirable not to sterilize by heating it but to sterilize it by filtering it through nano-filter(s).

[0044] The sugar osmotic agents which is used in the present invention may be glucose and the like that have reductive terminal residue(s), and tends to decompose easily through heat sterilization. In autoclave, if the sugar osmotic agents and heat stable suppressor(s) such as bisulfite, sulfite, sulfide, hydrosulfide and the like are sterilized together, the decomposition reaction of glucose may be suppressed, and consequently the concentration of glucose degradation products could be decreased below a detectable level.

[EXAMPLES]

[0045] The advantages of the present invention are explained by citing the results of actual verification examples. In the verification tests, the degree of protein cross linking by AGE formation were estimated by the determination of fluorescent intensity, based on the phenomenon that AGEs emits fluorescence, (Reference: Lee K.W. et al : "A systematic approach to evaluate the modification of lens protein by glycation-induced cross linking" Biochim.Biophys Acta: 1453(1) 141-151 Jan.6,1999)

(Estimation of suppression effect on protein cross linking by fluorescent intensity)

[Example 1] (Not the invention)

[0046] For estimating the suppression effect on protein cross linking sensitively, glyoxal was used. This chemical is one of glucose degradation products and much greater promoter to accelerate AGE formation than glucose does. Glyoxal (20mM/l) was added into phosphate buffer solution, in which human serum albumin (50mg/ml) was dissolved. As a test specimen, mercapto compound, listed in Table 1, was added into the solution at 20mM/l, then incubated at 37°C as long as for two weeks. Fluorescent intensity (FI) of the cross linked albumin in the incubated solution was determined at 440nm (excitation: at 370 nm) by fluorescence meter (Nihon Bunkou Co.). The suppression effect was estimated as follows; dividing the increase in the fluorescent intensity through the incubation in the case of "test specimen added ",

EP 1 323 440 B1

by that of "no test specimen added (control 1)". The result is shown in Table 1.

[Control 1]

5 **[0047]** In Control test 1, the fluorescent intensity was determined in the same manner as those in Example 1

[Table 1]

10

15

20

	Specimen(Cross linking Suppressors)	Increase in FI %
Example1 (a)	N-Acetylcysteine	7
(b)	2-Mercaptoethanol	42
(c)	Dithio erythrytol	31
(d)	Glutathione	25
(e)	S-Acetylmercaptosuccinic anhydride	50
(f)	Dimethylcysteine (Penicillamine)	83
(g)	Thiodiglycol	91
Control 1	Control (no specimen added)	100

[Example 2] (not the invention)

25

[0048] Methylglyoxal in place of glyoxal as GDP(Glucose Degradation Products) in Example 1 was used under the similar conditions as Example 1. Sodium bisulfite and sodium sulfite were tested as protein cross linking suppressors. Fluorescent intensity was determined in the same manners as in Example 1. The results are shown in Table 2.

30

[Control 2]

[0049] No protein cross linking suppressor, but methylglyoxal alone was added to human serum albumin. Fluorescent intensity was determined in the same manners as Example 2. The results are shown in Table 2.

35

[Table 2]

40

45

50

55

	Specimen(Cross linking suppressors)	Increase in FI (%)
Example 2 (a)	Sodium bisulfite	46
(b)	Sodium sulfite	61
Example 3 (a)	Acetyl salicylic acid	53
(b)	Ascorbic acid	86
Example 4 (a)	Quercetin dihydrate	19
(b)	Catechin hydrate	64
(c)	Epicatechin	72
Example 5	Heparin	96
Example 6	Amino guanidine hydrochloride	31
Example 7	N-Phenacyl thiazolium bromide	28
Example 8 (a)	N-(2-thiazolyl)-sulfanylamide	45
(b)	2-mercapto-4-methyl-thiazole acetic acid	82
(c)	4-(1,2,3,4-thia triazo)phenol hydrate	92
(d)	2-Oxo-thiazolidine-4-carboxylic acid	8

EP 1 323 440 B1

(continued)

	Specimen(Cross linking suppressors)	Increase in FI (%)
5	(e) 2-Mercaptothiazoline	48
	Control 2 Control (No suppressor added)	100

[Example 3] (not the invention)

10 **[0050]** The cross linking suppressors in Example 2 were replaced by acetylsalicylic acid and ascorbic acid and other conditions were similar to those in Example 2. The increase in fluorescent intensity was determined to estimate relative ratio to those of Control 2. The results are shown in Table 2.

15 [Example4] (not the invention)

20 **[0051]** The cross linking suppressors in Example 2 were replaced by quercetin dihydrate, catechin hydrate or epicatechin which were dissolved into dimethylsulfoxide, thence phosphate buffer solution was added at the ratio 1:3. The increase in fluorescent intensity was determined to estimate relative ratio to those of Control 2. The results are shown in Table 2.

[Example 5] (not the invention)

25 **[0052]** The cross linking suppressors in Example 2 were replaced by heparin, and other conditions were similar to those in Example 2. The increase in fluorescent intensity was determined to estimate relative ratio to those of Control 2. The results are shown in Table 2.

[Example 6] (not the invention)

30 **[0053]** The cross linking suppressors in Example 2 were replaced by amino guanidine, and other conditions were similar to those in Example 2. The increase in fluorescent intensity was determined to estimate relative ratio to those of Control 2. The results are shown in Table 2.

[Example 7] (not the invention)

35 **[0054]** The cross linking suppressors in Example 2 were replaced by N-phenacylthiazolium bromide which was dissolved into mixture of methanol and phosphate buffer solution at the ratio of 1:1 at 20mM/l.

[0055] The increase in fluorescent intensity was determined to estimate relative ratio to those of Control 2. The results are shown in Table 2.

40 [Example 8] (not the invention)

45 **[0056]** The cross linking suppressors in Example 7 were replaced by N-(2-thiazolyl)-sulfanilamide, 2-mercapto-4-methyl-thiazole acetic acid, 4-(1, 2, 3, 4-thia triazo) phenol hydrate, 2-Oxo-thiazolidine-4-carboxylic acid, or 2-Mercaptothiazoline.

[0057] Each compound was dissolved at the concentration of 20mM/l respectively in the mixture of dimethylsulfoxide and phosphate buffer solution at the ratio of 1:3. The increase in fluorescent intensity was determined to estimate relative ratio to those of Control 2. The results are shown in Table 2.

50 [Example 9, Control 3]

[0058] Collagen IV, methylglyoxal and one of the chemicals listed in Table 3 was dissolved into phosphate buffer solution, and incubated at 37°C as long as for 7 days. The increase in fluorescent intensity was determined to estimate relative ratio to those of Control 3, wherein methylglyoxal alone was added to collagen IV

55 **[0059]** * The test results are shown together with their redox potential in Table 3.

EP 1 323 440 B1

[Table 3]

	Specimen(Cross linking suppressors)	Increase in FI %	Redox potential (mV)	
5	Example 9 (a)	Sodium sulfide (Na ₂ S)	19	-616
	(b)	Sodium hydrosulfide (NaHS)	10	- 620
10	(c)	Sodium thiosulfate (Na ₂ S ₂ O ₃)	67	+40
	(d)	Sodium metabisulfite (Na ₂ S ₂ O ₅)	0	+17
	(e)	Sodium hydrosulfite(*)	0	- 79
	(f)	Sodium hydrosulfite (Na ₂ S ₂ O ₄)	0	- 360
15	(g)	Dimethylthiourea	2	- 40
	(h)	Sodium bisulfite (NaHSO ₃)	0	-127
	Control 3	No suppressor added	100	+282
20	* : Mixture of HSO ₃ and Na ₂ S ₂ O ₅ (Sigma Aldrich)			

[0060] The compounds in Table 3 are all known to be reductants having lower redox potential than that of saline solution; +160 ~ +180mV.

[0061] It is apparent that the reductants having lower redox potential than that of saline solution are effective suppressors of protein cross linking.

(Quantitative analysis of GDPs by redox potential titration)

[Example 10, Control 4] (not the invention)

[0062] The solution of sodium chloride and glucose was heated at 125°C for 45minutes in an autoclave and GDP was analyzed by redox potential titration method with 0.1M/l standard sodium thiosulfate solution. The results are shown in Table 4.

[Table 4]

	Additives	Additive concentration (μM/l)	GDP concentration (μM/l)	
35	Example 10 (a)	Sodium bisulfite	100	Not detected
40	(b)	Sodium hydrosulfide	50	Not detected
	(c)	Lactic acid	35,000	185
	(d)	Sodium bicarbonate	15,000	860
45	Control 4	No additive	-	560

(Effects of protein cross linkage splitter)

[Example11] (not the invention)

[0063] Human serum albumin was dissolved in phosphate buffer solution at 50mg/ml, wherein 3-deoxyglucosone was added at 50mM/l, thence the solution was incubated at 37°C for 7days, followed by the determination of fluorescent intensity.

[0064] Residual 3-deoxyglucosone was removed by membrane dialysis with phosphate buffer solution. N-phenacyl thiazolium bromide was added at 20mM/l concentration into the albumin solution (mixture of ethanol : phosphate buffer solution 1;1), Thence another 7days incubation was proceeded at 37°C. The fluorescent intensity after the second incubation (14th day) was compared with 7th day value. The decrease in FI is shown in Table 5.

[Example12] (not the invention)

[0065] Epicatechin and catechin dihydrate were dissolved into a mixture of dimethylsulfoxide (DMSO): phosphate buffer solution (1:3) at 20mM/l in place of N-phenacyl thiazolium bromide in Example 11 and used for the incubation for 7days in the same manner as in Example 9. After 14day incubation, the intensity of fluorescence (IF) was determined and the decrease in IF from that of after 7 days incubation was estimated. The results are shown in Table 5.

[Table 5]

	Additives added after 7days	Decrease %
Example 11	N-Phenacyl thiazolium bromide	- 21%
Example 12(a)	Epicatechin	- 6%
Example 12(b)	Catechin dihydrate	- 4%

[0066] It is apparent in Table 5 that the cross linkage in the albumin formed by 3-deoxyglucosone was split by N-phenacyl thiazolium bromide.

(Determination of suppression effect on AGE formation in peritoneum tissue)

[Control 5]

[0067] Intra-peritoneal infusion of 15ml of solution (A) listed in Table 6 was performed into peritoneal cavity of 7 weeks old rats (weight:300 ± 15g) every day for 5 days, thence the parietal peritoneum tissue was taken, frozen and sliced. The slice was stained with anti-AGE antibody and examined through a microscope. The observation results are shown in Table 7.

[Table 6]

NaCl	97 mM/l
Na ₂ CO ₃ ·CH (OH) CH ₃	17.5 mM/l
MgCl ₂	0.75 mM/l
CaCl ₂	1.75 mM/l
Aqueous solution/methanol ratio: 9/1	

[Control 6]

[0068] The solution (B) prepared by the addition of methylglyoxal into the mixed solution (A) in Control 5, was infused in the same manner as Control 5. The rat peritoneum tissue was stained with anti-AGE antibody and examined in the same way. The observation results are shown in Table 7.

[Example 13] (not the invention)

[0069] The solution (c) that is prepared by addition of electrolytes in Table 6, methylglyoxal (20mM/l) and N-phenacyl thiazolium bromide (20mM/l) into the mixed solution of water and methanol (1/9), in place of the solution (B) in Control 6, was infused in the same manner as Control 6. The rat peritoneum tissue was stained with anti-AGE antibody and examined in the same way. The observation result is shown in Table 7.

[Example 14] (not the invention)

[0070] The infusion was performed in the same manner as those in Control 6 for 5 days, thence the rat peritoneal cavity was rinsed, For another 5 days the rat was intraperitoneally infused 15ml of the solution (D), which was prepared by removing methylglyoxal from the solution (C), every day. The peritoneum tissue slice was submitted to anti-AGE antibody staining test. The result is shown in Table 7.

[Table 7]

	Solution	Additives	Anti-AGE antibody staining test	
5	Control 5	(A)	None	○
	Control 6	(B)	Methylglyoxal	×
10	Example 13	(C) *	Methylglyoxal + N-Phenacyl Thiazolium bromide	○
	Example 14	(D) *	After 5 ay infusion of the solution(B), N-Phenacyl thiazolium bromide alone infused for another 5 days	○
15	○ : Not stained with anti-AGE antibody × : Stained with anti-AGE antibody * Solution; electrolytes in the mixture of water/methanol: 9/1			

[0071] It is apparent in Table 7, that the solution (A), which does not contain GDPs, did not form AGE in peritoneum, while in the case of the solution (B) which contains GDP, the formation of AGE was clearly observed in the peritoneum tissue. In contrast, the solution (C), which was added N-phenacylthiazolium bromide, did not form AGE.

[0072] While, in the case of the second 5 days infusion with the solution (D) which contains N-phenacyl thiazolium bromide, the AGE, formed on the peritoneum tissue through the first 5days infusion, disappeared. It indicates that protein cross linkage may be split by N-phenacyl thiazolium bromide.

INDUSTRIAL APPLICABILITY

[0073] Peritoneal dialysis is favorable therapy for end stage renal disease as it is physiological dialysis and it allows the patients enjoy social activities. However, the removal of excess liquid is proceeded by hyper-osmolarity of the dialysate, so that the osmotic agent, glucose and its degradation products cause cross linking of the protein that consists of peritoneum, resulting in encapsulated peritoneum sclerosis, which prevents continuation of this therapy.

[0074] The present invention can supply the dialysate to solve this problem, that is, protecting the peritoneum by adding a sodium, potassium or other physiologically safe salt of thiosulfate or dithionite protein cross linking suppressors into the dialysate, or healing the peritoneum, which is suffering from sclerosis to some extent, by adding protein cross linkage splitters into the dialysate to allow the patients keep on the therapy for long period without suffering from peritoneum sclerosis, and to retain ultra filtration capacity. So that the present invention supplies an effective means to enable the therapy keeping on without problem

Claims

1. Dialysate of peritoneal dialysis comprising an aqueous solution of electrolytes and sugar osmotic agents and a sodium, potassium or other physiologically safe salt of thiosulfate or dithionite.
2. A dialysate of peritoneal dialysis according to claim 1, which contains a non-sugar oncotic agent as an osmotic agent.
3. A preparation method of a dialysate of peritoneal dialysis according to claim 1 or claim 2, which method comprises sterilizing dialysate under high temperature and pressure, and thereafter mixing the dialysate with the physiologically safe salt which is sterilized separately.
4. A preparation method of a dialysate of peritoneal dialysis according to claim 1 or claim 2, which method comprises sterilizing sugar osmotic agent and the physiologically safe salt under high temperature and pressure, and thereafter mixing with electrolytes which is sterilized separately.

Patentansprüche

1. Dialysat einer Peritonealdialyse mit einer wässrigen Lösung von Elektrolyten und osmotisch aktiven zuckerartigen Substanzen und einem Natrium-, Kalium- oder einem anderen physiologisch sicheren Salz von Thiosulfat oder

Dithionit.

2. Dialysat einer Peritonealdialyse nach Anspruch 1, das eine nicht zuckerartige, onkotisch aktive Substanz als osmotisch aktive Substanz enthält.

5 3. Herstellungsverfahren für ein Dialysat einer Peritonealdialyse nach Anspruch 1 oder Anspruch 2, wobei das Verfahren aufweist: Sterilisieren von Dialysat unter hoher Temperatur und hohem Druck, und danach Vermischen des Dialysats mit dem physiologisch sicheren Salz, das getrennt sterilisiert wird.

10 4. Herstellungsverfahren für ein Dialysat einer Peritonealdialyse nach Anspruch 1 oder Anspruch 2, wobei das Verfahren aufweist: Sterilisieren einer zuckerartigen osmotisch aktiven Substanz und des physiologisch sicheren Salzes unter hoher Temperatur und hohem Druck, und danach Vermischen mit Elektrolyten, die getrennt sterilisiert werden.

15 **Revendications**

1. Dialysat d'une dialyse péritonéale comprenant une solution aqueuse d'électrolytes et des agents osmotiques sucrés et un sel de sodium, de potassium ou d'un autre sel physiologiquement sain de thiosulfate ou de dithionite.

20 2. Dialysat d'une dialyse péritonéale selon la revendication 1, qui contient un agent oncotique non sucré en tant qu'agent osmotique.

25 3. Méthode de préparation d'un dialysat d'une dialyse péritonéale selon la revendication 1 ou la revendication 2, laquelle méthode comprend la stérilisation du dialysat sous une température et une pression élevées et par la suite le mélange du dialysat avec le sel physiologiquement sain qui est stérilisé séparément.

30 4. Méthode de préparation d'un dialysat d'une dialyse péritonéale selon la revendication 1 ou la revendication 2, laquelle méthode comprend la stérilisation d'un agent osmotique sucré et du sel physiologiquement sain sous une température et une pression élevées et par la suite le mélange avec des électrolytes, qui sont stérilisés séparément.

35

40

45

50

55

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- JP HEI8150930 B [0010] [0040]
- JP HEI9302388 B [0010] [0040]
- WO 0189478 A, (OLSSON LARS FRIDE ;WIESLANDER ANDERS (SE); GAMBRO LUNDIA AB (SE);) [0011]
- WO 02053094 A, (DEPPISCH REINHOLD ;WIESLANDER ANDERS (SE); GAMBRO LUNDIA AB (SE);) [0011]

Non-patent literature cited in the description

- ABSORPTION OF SODIUM BISULFITE FROM PERITONEAL DIALYSIS SOLUTIONS. **HALABY S F et al.** JOURNAL OF PHARMACEUTICAL SCIENCES. AMERICAN PHARMACEUTICAL ASSOCIATION, January 1965, vol. 54, 52-55 [0011]
- SULFIT IN INFUSIONSLOESUNGEN. **LANG K et al.** ZEITSCHRIFT FUER ERNAHRUNGSWISSENSCHAFT. STEINKOPF VERLAG, 1979, vol. 18, 37-41 [0011]
- HEAT STERILIZATION OF GLUCOSE-CONTAINING FLUIDS FOR PERITONEAL DIALYSIS: BIOLOGICAL CONSEQUENCES OF CHEMICAL ALTERATIONS. **WIESLANDER A P et al.** PERITONEAL DIALYSIS INTERNATIONAL. PERGAMON PRESS, 1995, vol. 15, S52-S60 [0011]
- INTRAPERITONEAL CIS-DIAMMINIDICHLOROPLATINIUM WITH SYSTEMIC THIOSULFATE PROTECTION. **HOWELL S B et al.** CANCER RESEARCH. AMERICAN ASSOCIATION FOR CANCER RESEARCH, March 1983, vol. 43, 1426-1431 [0011]
- **Lee K.W. et al.** A systematic approach to evaluate the modification of lens protein by glycation-induced cross linking. *Biochim.Biophys Acta*, 06 January 1999, vol. 1453 (1), 141-151 [0045]