



(86) Date de dépôt PCT/PCT Filing Date: 2003/05/26
 (87) Date publication PCT/PCT Publication Date: 2003/12/04
 (45) Date de délivrance/Issue Date: 2012/05/15
 (85) Entrée phase nationale/National Entry: 2004/11/24
 (86) N° demande PCT/PCT Application No.: JP 2003/006518
 (87) N° publication PCT/PCT Publication No.: 2003/099319
 (30) Priorité/Priority: 2002/05/24 (JP2002/151220)

(51) Cl.Int./Int.Cl. *A61K 38/17* (2006.01),
A61K 48/00 (2006.01), *A61P 43/00* (2006.01),
A61P 9/10 (2006.01)

(72) Inventeurs/Inventors:
 KADOWAKI, TAKASHI, JP;
 YAMAUCHI, TOSHIMASA, JP;
 KUBOTA, NAOTO, JP;
 TERAUCHI, YASUO, JP;
 KUBOTA, TETSUYA, JP;
 NODA, TETSUO, JP;
 NAGAI, RYOZO, JP;
 IMAI, YASUSHI, JP

(73) Propriétaire/Owner:

(54) Titre : **MEDICAMENT PREVENTIF/CURATIF POUR L'ARTERIOSCLEROSE COMPRENANT DE L'ADIPONECTINE**
 (54) Title: **PREVENTIVE/REMEDY FOR ARTERIOSCLEROSIS COMPRISING ADIPONECTIN**

(57) **Abrégé/Abstract:**

A scavenger receptor A expression down-regulator and a drug for preventing or treating arteriosclerosis which contain, as the active ingredient, a C-terminal globular domain of adiponectin, adiponectin, or a gene encoding the domain or adiponectin. According to the present invention, there is provided a preventive or therapeutic agent capable of directly preventing intimal thickening, which constitutes an essential feature of arteriosclerosis. This effect can be attained through arresting the onset and development of arteriosclerosis by reducing the expression level of scavenger receptor A in arterial walls and preventing lipid buildup in macrophages.



(73) Propriétaires(suite)/Owners(continued): JAPAN SCIENCE AND TECHNOLOGY AGENCY, JP

(74) Agent: GOWLING LAFLEUR HENDERSON LLP

Abstract

A scavenger receptor A expression down-regulator and a drug for preventing or treating arteriosclerosis which contain, as the active ingredient, a C-terminal globular domain of adiponectin, adiponectin, or a gene encoding the domain or adiponectin.

According to the present invention, there is provided a preventive or therapeutic agent capable of directly preventing intimal thickening, which constitutes an essential feature of arteriosclerosis. This effect can be attained through arresting the onset and development of arteriosclerosis by reducing the expression level of scavenger receptor A in arterial walls and preventing lipid buildup in macrophages.

DEMANDES OU BREVETS VOLUMINEUX

**LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVETS
COMPREND PLUS D'UN TOME.**

CECI EST LE TOME __1__ DE __2__

NOTE: Pour les tomes additionels, veuillez contacter le Bureau Canadien des Brevets.

JUMBO APPLICATIONS / PATENTS

**THIS SECTION OF THE APPLICATION / PATENT CONTAINS MORE
THAN ONE VOLUME.**

THIS IS VOLUME __1__ OF __2__

NOTE: For additional volumes please contact the Canadian Patent Office.

Description

Preventive/Remedy for Arteriosclerosis

Comprising Adiponectin

Technical Field

The present invention relates to a drug for preventing or treating arteriosclerosis.

Background Art

The term "arteriosclerosis" refers to a pathological condition where the walls of an artery lose elasticity and become brittle. Arteriosclerosis is one of key factors causing adult diseases, including cerebral hemorrhage, cerebral infarction, myocardial infarction, and nephrosclerosis. Known causes of arteriosclerosis include hyperlipidemia, and bacteria, viruses, or lipid peroxide in blood. However, the pathogenesis of arteriosclerosis has not yet been fully elucidated. In any case, since arteriosclerosis has been observed to begin with thickening of arterial walls caused by damage to the arterial intima or endothelium, there is keen demand for development of a drug capable of inhibiting thickening of the arterial intima.

Accordingly, an object of the present invention is to provide a drug which is effective for preventing or treating arteriosclerosis.

Disclosure of the Invention

Under the above circumstance, the present inventors have devoted their research efforts to pharmacological actions of adiponectin, which is known to have an insulin resistance reducing effect, and have found that adiponectin-gene-deficient mice show significantly thickened arterial intima; and that adiponectin, a C-terminal globular domain thereof, or a gene thereof is useful as a preventive or therapeutic drug for arteriosclerosis, on the basis of their experimental results that when apoE-deficient mice, which are employed as atherosclerosis onset model mice, are manipulated to over-express adiponectin, in particular, the C-terminal globular domain of adiponectin, the onset of arteriosclerosis is suppressed. In addition, since over-expression of adiponectin lowers the expression level of scavenger receptor A without significantly affecting the levels of free fatty acid, neutral fat, or total cholesterol in blood, the inventors have concluded that the arteriosclerosis preventive action of adiponectin lowers the expression level of scavenger receptor A, whereby accumulation of lipids in macrophages is prevented. The present invention has been accomplished on the basis of these findings.

Accordingly, the present invention provides a drug for preventing or treating arteriosclerosis, the drug containing, as an active ingredient thereof, a C-terminal globular domain of adiponectin, adiponectin, or a gene encoding the C-terminal globular domain of adiponectin or the adiponectin.

The present invention also provides a scavenger

receptor A expression down-regulator which contains, as an active ingredient thereof, a C-terminal globular domain of adiponectin, adiponectin, or a gene encoding the C-terminal globular domain of adiponectin or the adiponectin.

The present invention also provides use, in the manufacture of a drug for preventing or treating arteriosclerosis, of a C-terminal globular domain of adiponectin, adiponectin, or a gene encoding the C-terminal globular domain of adiponectin or the adiponectin.

The present invention also provides use, in the manufacture of a scavenger receptor A expression down-regulator, of a C-terminal globular domain of adiponectin, adiponectin, or a gene encoding the C-terminal globular domain of adiponectin or the adiponectin.

The present invention also provides a method for treating arteriosclerosis, comprising administering, to a subject in need thereof, an effective amount of a C-terminal globular domain of adiponectin, adiponectin, or a gene encoding the C-terminal globular domain of adiponectin or the adiponectin.

The present invention also provides a method for down-regulating the expression level of scavenger receptor A in a patient, comprising administering to the patient an effective amount of a C-terminal globular domain of adiponectin, adiponectin, or a gene encoding the C-terminal globular domain of adiponectin or the adiponectin.

Brief Description of the Drawings

Fig. 1 schematically shows a gene targeting performed on adiponectin gene deficiency, in which, a restriction map of a mouse adiponectin gene (top), an adiponectin gene targeting vector (middle), and a deduced targeting allele (bottom).

Fig. 2 shows the results of Southern blotting of ES-cell-derived DNA samples which have been digested with SpeI and EoRV. The bands of 17 kb are obtained from wild-type alleles, and those of 10.5 kb are from mutated alleles.

Fig. 3 shows the results of Southern blotting of SpeI- and EoRV-digested DNA samples from a wild-type mouse, a hetero-deficient (adipo +/-) mouse, and a homo-deficient (adipo -/-) mouse. The bands of 17 kb are obtained from wild-type alleles, and those of 10.5 kb are from mutated alleles.

Fig. 4 shows the results of Northern blotting of white adipose tissue samples from a wild-type mouse, a hetero-deficient (adipo +/-) mouse, and a homo-deficient (adipo -/-) mouse.

Fig. 5 shows blood adiponectin level of a wild-type mouse, a hetero-deficient (adipo +/-) mouse, and a homo-deficient (adipo -/-) mouse. **P<0.01.

Fig. 6 shows blood leptin level of a wild-type mouse, a hetero-deficient (adipo +/-) mouse, and a homo-deficient (adipo -/-) mouse.

Fig. 7 shows the body weight, at 6 weeks of age, of a wild-type mouse, a hetero-deficient (adipo +/-) mouse, and a

homo-deficient (adipo -/-) mouse.

Fig. 8 shows the results of an insulin tolerance test performed on a wild-type mouse and a hetero-deficient (adipo +/-) mouse at 6 weeks of age. *P<0.05.

Fig. 9 shows the results of a glucose tolerance test performed on a wild-type mouse and a hetero-deficient (adipo +/-) mouse at 6 weeks of age. *P<0.05.

Fig. 10 shows the results of a glucose tolerance test performed on a wild-type mouse and a hetero-deficient (adipo +/-) mouse after having loaded with a high-fat diet for 10 weeks. *P<0.05, **P<0.01.

Fig. 11 shows the results of an insulin tolerance test performed on a wild-type mouse and a homo-deficient (adipo -/-) mouse, at 6 weeks of age. *P<0.05, **P<0.01.

Fig. 12 shows the results of a glucose tolerance test performed on a wild-type mouse and a hetero-deficient (adipo +/-) mouse, at 6 weeks of age. *P<0.05, **P<0.01.

Fig. 13 shows levels, in blood, of free fatty acid (FFA), neutral fat (TG), total cholesterol (TC) of a wild-type mouse and a hetero-deficient (adipo +/-) mouse.

Fig. 14 shows levels, in blood, of free fatty acid (FFA), neutral fat (TG), total cholesterol (TC) of a wild-type mouse and a homo-deficient (adipo -/-) mouse.

Fig. 15 shows the inner diameter of a blood vessel of a wild-type mouse and a hetero-deficient (adipo +/-) mouse, as measured two weeks after the mice underwent cuff placement.

Fig. 16 shows the degree of intimal thickening of a

wild-type mouse and a hetero-deficient (adipo +/-) mouse, as measured two weeks after the mice underwent cuff placement.

Fig. 17 shows the degree of medial thickening of a wild-type mouse and a hetero-deficient (adipo +/-) mouse, as measured two weeks after the mice underwent cuff placement.

Fig. 18 shows the intima/media ratio of a wild-type mouse and a hetero-deficient (adipo +/-) mouse, as measured two weeks after the mice underwent cuff placement.

Fig. 19 shows the foci of arteriosclerosis in an apoE-deficient (apoE-/-:a) mouse and a gAd-overexpressed apoE-deficient (gAd Tg apoE-/-:b) mouse.

Fig. 20 shows the areas of the foci of arteriosclerosis in an apoE-deficient (apoE-/-) mouse and a gAd-overexpressed apoE-deficient (gAd Tg apoE-/-) mouse (aortic arch (b), descending aorta (c), and their sum (a)).

Fig. 21 shows the results of Oil Red O staining, reaction with anti-Mac3 antibody, and reaction with anti-scavenger receptor A antibody as observed in an apoE-deficient (apoE-/-) mouse and a gAd-overexpressed apoE-deficient (gAd Tg apoE-/-) mouse.

Best Mode for Carrying Out the Invention

Adiponectin, which is employed in the present invention, has already been cloned (Maeda, K. et al., *Biochem. Biophys. Res. Commun.* 221, 286-296 (1996), Nakano, Y. et al., *J. Biochem. (Tokyo)* 120, 802-812 (1996)), and is available through known means. SEQ ID NOs: 1 and 2 show the nucleotide

sequence and the amino acid sequence of human adiponectin. Adiponectin is composed of an N-terminal collagen-like sequence (cAd) a C-terminal globular domain (gAd; in SEQ ID NO: 1, amino acid Nos. 114 to 239 or 111 to 242). The C-terminal globular domain (gAd) is preferred as it provides stronger arteriosclerosis preventive and therapeutic effects than full length adiponectin. SEQ ID NOs: 3 and 4 show the nucleotide sequence and the amino acid sequence of mouse adiponectin. The N-terminal collagen-like sequence (cAd) of the mouse adiponectin stretches from 45 to 109 (amino acid No.), and the C-terminal globular domain (gAd) stretches from 110 to 247 (amino acid No.). According to the present invention, not only proteins comprising the amino acid sequence of any of SEQ ID NOs: 1 to 4 or an amino acid sequence having the gAd domain, but also a protein comprising a modified amino acid sequence derived from substitution, deletion, or addition of one or more amino acid residues of any of these amino acid sequences may be employed, so long as it provides an effect as exhibited by adiponectin. Examples of such mutated proteins include those having 80% or higher homology, preferably 90% or higher homology, to any of the amino acid sequences of SEQ ID NOs: 1 to 4 or an amino acid sequence including the gAd domain.

Examples of the gene which is employed in the present invention include the genes coding for adiponectin (i.e., SEQ ID NOs: 1 and 3) and a gene coding for gAd. Also, there may be employed a gene having a nucleotide sequence capable of

hybridizing with any of these genes under stringent conditions.

Adiponectin or a polypeptide which forms a portion of adiponectin (including gAd) may be isolated from cells containing the same. However, since a gene coding for adiponectin has already been cloned, the adiponectin or the polypeptide may be prepared through a DNA recombinant technique; i.e., making use of transformant cells created by use of expression vectors produced through use of the gene.

As will be described hereinbelow, adiponectin-deficient mice exhibit high levels of neutral fat in blood, but their cholesterol levels are comparable to those of wild-type mice. Moreover, adiponectin-deficient mice, representing an arteriosclerosis model, exhibited intima thickening which was twice the thickness as observed in wild-type mice. In contrast, when apoE-deficient mice, which represent a spontaneous atherosclerosis model, are caused to over-express gAd, they exhibit a significant reduction in arteriosclerotic area, preventing development of arteriosclerosis. However, overexpression of gAd induced in apoE-deficient mice only insignificantly affect general risk factors for arteriosclerosis, such as body weight and blood sugar, and free fatty acid, neutral fat, and total cholesterol in blood. On the other hand, over-expression of gAd induced in apoE-deficient mice was found to exhibit a lowered expression of scavenger receptor A in arterial walls. Scavenger receptor A is a receptor which, when macrophages engulf modified LDL,

binds to the modified LDL on the surface of a cell, and is known to play a key role as a receptor which triggers the onset of arteriosclerosis.

Accordingly, adiponectin, gAd, or a gene coding for adiponectin or gAd is useful as a down-regulator of scavenger receptor A expression, or as a drug for preventing or treating arteriosclerosis. In particular, gAd or a gene encoding gAd is very useful in that it exhibits a more potent down-regulating effect on expression of a scavenger receptor as compared with adiponectin, and stronger preventive or therapeutic effect.

For administering the drug of the present invention to a mammal including a human, pharmaceutical compositions of a variety of dosage forms may be produced through incorporation of a pharmacologically acceptable carrier to any of the aforementioned active ingredients. Among such dosage forms, preparations for injection are preferred. Examples of the pharmacologically acceptable carrier include distilled water, a solubilizer, a stabilizer, an emulsifier, and a buffer. The dose of any of the drugs may differ depending on the condition of the disease, sex, body weight, etc., and may range from 0.1 μ g to 10 mg/day or thereabouts, as reduced to the amount of adiponectin or gAd.

Examples

The present invention will next be described in more detail by way of examples, which should not construed as

limiting the invention thereto.

A. Methods

(1) Preparation of knockout mice

Screening of a 129/Sv mouse genomic library was performed using adiponectin cDNA as a probe, whereby a plurality of clones harboring a gene encoding adiponectin were obtained. A targeting vector was constructed, in which the region stretching from the translation initiation site to the translation termination site had been replaced by a neomycin-resistant gene. ES cells were transfected with the resultant targeting vector. Screening was performed through Southern blotting, whereby homologous recombinants of 5 clones were confirmed. Chimeric mice were created by means of microinjection, and the mice were crossbred with BI/6 to thereby produce F1, and then F2.

Briefly, an adiponectin-gene-deficient mouse was produced through homologous recombination as shown in Fig. 1. With an aim to knock out the mouse adiponectin gene, a targeting vector in which exons 2 and 3 that encode adiponectin were replaced with a neo resistant gene was prepared. Separate 5 homologous recombinant clones were confirmed through Southern blotting (Fig. 2). From ES cells having 129/Sv as a background, chimeric mice were produced, and in order to create a hetero-deficient mouse, they were cross-bred with BI/6. The genotype was confirmed through Southern blotting (Fig. 3).

(2) Insulin tolerance test

Human insulin was intraperitoneally administered to test mice in an amount of 0.7 mU per gram (body weight), and the mice were fasted during the tolerance test. The blood was collected from the tail vein, and blood sugar level was measured by means of a Glutest Ace (registered trademark, product of Sanwa Kagaku Kenkyusho Co., Ltd.).

(3) Glucose tolerance test

Glucose was perorally administered to test mice in an amount of 1.5 mg per gram (body weight). Prior to the administration, the mice had been fasted for at least 16 hours. The blood was collected from the fundus vein, and blood sugar level and insulin level were measured by means of a Glutest Ace (registered trademark, product of Sanwa Kagaku Kenkyusho Co., Ltd.) and a rat insulin RIA kit (product of Amersham Pharmacia Biotech), respectively.

(4) Measurement of blood lipid level

After the test mice were fasted for 16 hours, levels of free fatty acid, neutral fat, and total cholesterol, all in blood, were measured by means of a NEFAC-test, a TGL-type, and a Tchol E-type (product of Wako), respectively.

(5) Measurement of blood leptin level and blood adiponectin level

After the mice were fasted for 16 hours, levels of leptin and adiponectin, both in blood, were measured by means of a Quintikine M kit (product of R&D) and an adiponectin RIA kit (product of Linco), respectively.

(6) Creation of a thick vascular intima model through cuff

placement

A 2.0-mm polyethylene tube (PE-50) was placed in the femoral artery. When two weeks had passed, the artery was press-fixed with formalin, and removed together with the opposite-side, uncuffed artery, which served as a control artery. Each of the thus-removed blood vessels was sliced to obtain continuous ring-shaped specimens, each having a length of 10 mm. Ten specimens were taken and HE staining was performed. The inner diameter of the blood vessel, the thickness of the intima, and the thickness of the media were measured, and intima/media ratio was calculated.

(7) Preparation of gAd-overexpressed mice

According to the method described in Diabetes 48, 1822-1829 (1999), a fused gene containing a human SAP promoter and mouse gAd cDNA was prepared. Purified Hind III-XhoI fragments were microinjected to pronuclei of fertilized ova of C57BL6 mice (product of Clea Japan, Inc.). Tail DNA samples obtained from the resultant transgenic mice were subjected to Southern blotting through use of a gAd cDNA probe for the Bgl II/Hinc II site of gAd, whereby gAd overexpression of the transgenic mice was confirmed.

(8) Production of gAd-overexpressing apoE-deficient mice

gAd-overexpressing apoE-deficient mice were crossbred, to thereby produce gAd-overexpressing apoE-hetero-deficient mice. The resultant mice were crossed further with apoE-deficient mice, to thereby create apoE-deficient mice exhibiting over-expression of gAd.

(9) Measurement of blood sugar level and lipid level

Mice were fed until they were full, and their blood sugar level and levels, in blood, of free fatty acid, neutral fat, and total cholesterol were measured by means of a Glutest Ace (registered trademark, product of Sanwa Kagaku Kenkyusyo Co., Ltd.), an NEFA C-test, a TGL-type, and a Tchol E-type (Products of Wako), respectively.

(10) Evaluation of the size of arteriosclerotic foci

From each of gAd-overexpressed apoE-deficient mice (4 months old) and control apoE-deficient mice, the aortic arch and the descending aorta were removed, fixed with formalin, and then subjected to staining with Sudan IV. The arteriosclerotic foci were evaluated in terms of their size.

(11) Evaluation in terms of buildup of cholesterol ester, expression level of scavenger receptor A, and macrophage accumulation

Frozen samples of continuous ring-shaped specimens of the annulus portion of the aorta were prepared. Ten such samples were subjected to immunostaining by use of Oil Red O, anti-scavenger receptor A antibody, or anti-Mac3 antibody (a macrophage-specific marker), whereby buildup of cholesterol ester, expression level of scavenger receptor A, and macrophage accumulation were evaluated, respectively.

B. Results

(1) Mouse-adiponectin-gene-deficient mice

Through Northern blotting of white adipose tissue, the expression level of adiponectin in the hetero-deficient mice

was found to be reduced by about 60%, and the homo-deficient mice were found to exhibit completely no adiponectin expression (Fig. 4). Indeed, when blood adiponectin level was measured in the hetero-deficient mice, the magnitude of reduction was found to be about 60%, and the level in the hetero-deficient mice was found to be lower than the undetectable level (Fig. 5). With respect to the blood leptin level, no difference was observed (Fig. 6).

(2) Insulin resistance of mouse-adiponectin-gene-deficient mice

In three groups of 6-week-old mice; i.e., wild-type group, hetero-deficient (adipo +/-) group, and homo-deficient (adipo -/-) group, there was no difference in terms of body weight (Fig. 7). The 6-week-old wild-type mice and hetero-defective mice of the same age were subjected to an insulin tolerance test, to thereby check their insulin sensitivity. The degree of reduction in blood sugar level in response to exogenous insulin was statistically significantly low in the hetero-deficient mice, proving that the hetero-deficient mice had insulin resistance (Fig. 8).

Next, a glucose tolerance test was performed. No difference was observed between the two groups of wild-type mice and hetero-deficient mice in terms of blood sugar or insulin level (Fig. 9). However, as compared with the wild-type mice, the hetero-deficient mice, after having been loaded with 10-week high fat diet, exhibited a significantly high blood sugar level before and after loading with glucose,

though the body weight remained in a similar level (Fig. 10).

Afterwards, analysis on the homo-deficient mice was performed.

An insulin tolerance test performed on 6-week-old wild-type mice and homo-deficient mice of the same age. As compared with the wild-type mice or the hetero-deficient mice, the degree of reduction in blood sugar level in response to exogenous insulin was statistically significantly low in the homo-deficient mice, proving that the homo-deficient mice had insulin resistance higher than the corresponding levels of the wild-type mice and homo-deficient mice (Fig. 11).

Next, a glucose tolerance test was performed. In both stages of during fasting and after glucose loading, the homo-deficient mice exhibited blood sugar levels higher than the case of wild-type mice. This substantiates that homo-deficient mice had slightly impaired glucose tolerance in addition to insulin resistance (Fig. 12). Before administration and 30 minutes after administration, no difference was observed between the wild-type group and the homo-deficient group in terms of the insulin levels before and after glucose loading. However, the homo-deficient mice showed a somewhat low insulin level at 15 min (Fig. 12).

(3) Blood neutral fat level in adiponectin homo-deficient mice

In order to check the effect of adiponectin on lipid metabolism, levels, in blood, of free fatty acid (FFA), neutral fat (TG), and total cholesterol (TC) were determined

in wild-type, hetero-deficient, and homo-deficient mice (Figs. 13 and 14). The hetero-deficient mice did not show any difference in level of any of the three test items as compared with the wild-type mice (Fig. 13). However, the homo-deficient mice showed significantly higher blood neutral fat levels than the wild-type mice (Fig. 14).

(4) Thickening of intima in cuff-injured models of mouse adiponectin hetero-deficient mice

In order to investigate the effect of adiponectin on arteriosclerosis, the degree of intimal thickening induced by cuff placement was measured in the wild-type mice and the hetero-deficient mice for comparison therebetween. No difference was observed between the two groups in terms of the vascular inner diameter after cuff-induced injury was created (Fig. 15). When 2 weeks had elapsed after creation of cuff injury, the hetero-deficient mice showed about 1.8 times the thickness of the intima of the wild-type mice (Fig. 16). However, no difference was observed between the two groups in terms of the thickness of the media (Fig. 17). The intima/media ratio of the hetero-deficient group exhibited a ratio about two-fold that of the wild-type mice (Fig. 18).

(5) Prevention of the onset of arteriosclerosis in gAd-overexpressing apoE-deficient mice

ApoE-deficient mice, which represent a spontaneous arteriosclerosis model, were caused to overexpress gAd and studied whether or not onset of arteriosclerosis was prevented. The results are shown in Figs. 19 and 20. In Fig.

19, "a" shows the results of Sudan IV staining of aorta samples from apoE-deficient mice, and "b" shows the results of Sudan IV staining of aorta samples from gAd-overexpressed apoE-deficient mice. As is evident from the comparison between "a" and "b," gAd-overexpressed apoE-deficient mice clearly show a reduction in the incidence of arteriosclerotic foci. Fig. 20 shows comparison with respect to the area of arteriosclerotic foci. Fig. 20 shows that over-expression of gAd caused significant reduction in the area of arteriosclerotic foci, which are stained with Sudan IV, in any case of aortic arch (b), descending aorta (c), the sum of the mentioned two cases (a), indicating arresting of the onset of arteriosclerosis.

(6) Effect of gAd overexpression on arteriosclerosis risk factors in apoE-deficient mice on a normal diet

The body weight, blood sugar level, and levels, in blood, of free fatty acid, neutral fat, and total cholesterol of gAd-overexpressed apoE-deficient mice on a normal diet are shown in Table 1.

Table 1

	Mouse		Statistical significance
	apoE ^{-/-}	gAd Tg apoE ^{-/-}	
Body weight(g)	29.8±1.2	29.9±1.5	none
Plasma glucose level (mg/dl)	145±4	152±8	none
Serum total cholesterol level (mg/dl)	541±49	509±32	none
Serum triglyceride level (mg/dl)	127±52	104±24	none
Serum free fatty acid level (mEq/L)	0.53±0.08	0.57±0.04	none

Mean ± s.e. (n=5)

As shown in Table 1, when apoE-deficient mice on a normal diet were caused to over-express gAd, no significant effect was exerted on arteriosclerosis risk factors such as body weight and blood sugar, and free fatty acid, neutral fat, and total cholesterol in blood. This suggests that gAd possibly acts on vascular walls or macrophages directly, to thereby exhibit anti-arteriosclerotic activity.

(7) Mechanism of arresting onset of arteriosclerosis by gAd

With an aim to elucidate the mechanism of the interaction between gAd and vascular walls or macrophages, frozen samples of continuous ring-shaped slices of annulus portion of the aorta were subjected to immunostaining by use of Oil Red O, anti-scavenger receptor A antibody, and a macrophage-specific marker; i.e., anti-Mac3 antibody. As a result, as shown in Fig. 21, over-expression of gAd, though having no significant impact on accumulation of macrophages, were found to reduce the expression level of scavenger receptor A, suppress buildup of lipids in macrophages, and arrest the onset of arteriosclerosis.

Industrial Applicability

The present invention provides a preventive or therapeutic agent capable of directly preventing intimal thickening, which constitutes an essential feature of arteriosclerosis, wherein this effect can be attained through arresting the onset and development of arteriosclerosis by reducing the expression level of scavenger receptor A in

arterial walls and preventing lipid buildup in macrophages.

DEMANDES OU BREVETS VOLUMINEUX

**LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVETS
COMPREND PLUS D'UN TOME.**

CECI EST LE TOME 1 DE 2

NOTE: Pour les tomes additionels, veuillez contacter le Bureau Canadien des Brevets.

JUMBO APPLICATIONS / PATENTS

**THIS SECTION OF THE APPLICATION / PATENT CONTAINS MORE
THAN ONE VOLUME.**

THIS IS VOLUME 1 OF 2

NOTE: For additional volumes please contact the Canadian Patent Office.

CLAIMS

1. A drug for preventing or treating arteriosclerosis, comprising a C-terminal globular domain of adiponectin or a gene
5 encoding the domain, wherein the C-terminal globular domain consists of amino acids 114 to 239 of SEQ ID NO: 2 or amino acids 111 to 242 of SEQ ID NO: 2.

2. The drug for preventing or treating arteriosclerosis
10 as claimed in claim 1, further comprising a pharmacologically acceptable carrier.

3. A drug for downregulating scavenger receptor A expression, comprising a C-terminal globular domain of
15 adiponectin or a gene encoding the domain, wherein the C-terminal globular domain consists of amino acids 114 to 239 of SEQ ID NO: 2 or amino acids 111 to 242 of SEQ ID NO: 2.

4. Use of a C-terminal globular domain of adiponectin
20 or a gene encoding the domain in manufacture of a drug for preventing or treating arteriosclerosis, wherein the C-terminal globular domain consists of amino acids 114 to 239 of SEQ ID NO: 2 or amino acids 111 to 242 of SEQ ID NO: 2.

25 5. Use of a C-terminal globular domain of adiponectin or a gene encoding the domain in manufacture of a drug for downregulating scavenger receptor A expression, wherein the C-terminal globular domain consists of amino acids 114 to 239 of SEQ ID NO: 2 or amino acids 111 to 242 of SEQ ID NO: 2.

30

6. Use of a C-terminal globular domain of adiponectin

or a gene encoding the domain, wherein the C-terminal globular domain consists of amino acids 114 to 239 of SEQ ID NO: 2 or amino acids 111 to 242 of SEQ ID NO: 2 for preventing or treating arteriosclerosis, in a subject in need thereof.

5

7. Use of a C-terminal globular domain of adiponectin or a gene encoding the domain, or a gene encoding the domain, wherein the C-terminal globular domain consists of amino acids 114 to 239 of SEQ ID NO: 2 or amino acids 111 to 242 of SEQ ID NO: 2, for
10 down-regulating expression of scavenger receptor A in a subject in need thereof.

Application number / numéro de demande: JP03/06518

Figures: 2/3/4/19/21

Unscannable items
received with this application
(Request original documents in File Prep. Section on the 10th floor)

Documents reçu avec cette demande ne pouvant être balayés
(Commander les documents originaux dans la section de préparation des dossiers au
10^{ème} étage)

Fig. 1




Fig. 5




Fig. 6




Fig. 7




Fig. 8




Fig. 9





Fig. 10





Fig. 11




Fig. 12




Fig. 13




Fig. 14




Fig. 15




Fig. 16




Fig. 17




Fig. 18




Fig. 20

