



(11) **EP 2 311 477 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention of the grant of the patent:
09.07.2014 Bulletin 2014/28

(51) Int Cl.:
A61K 38/00 (2006.01) C07K 14/47 (2006.01)

(21) Application number: **10191552.8**

(22) Date of filing: **20.03.2007**

(54) **Use of ip3 receptor-binding protein for controlling phosphatidylinositol metabolism**

Verwendung von ip3-Rezeptorbindungsprotein zur Regelung des Phosphatidylinositol-Stoffwechsels

Utilisation de ip3-récepteur-binding- protein pour le contrôle du métabolisme phosphatidylinositol

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IS IT LI LT LU LV MC MT NL PL PT RO SE SI SK TR

(30) Priority: **20.03.2006 JP 2006077607**

(43) Date of publication of application:
20.04.2011 Bulletin 2011/16

(62) Document number(s) of the earlier application(s) in accordance with Art. 76 EPC:
07739967.3 / 2 014 297

(73) Proprietor: **Japan Science and Technology Agency Kawaguchi-shi, Saitama 332-0012 (JP)**

(72) Inventors:
• **Mikoshihira, Katsuhiko Mitaka-shi Tokyo 181-0001 (JP)**

• **Ando, Hideaki Wako-shi Saitama 351-0114 (JP)**
• **Mizutani, Akihiro Tokyo 142-0063 (JP)**

(74) Representative: **O'Brien, Simon Warwick D Young & Co LLP 120 Holborn London EC1N 2DY (GB)**

(56) References cited:
EP-A- 1 408 049

• **K.A. LAMIA ET AL.: "Increased insulin sensitivity and reduced adiposity in phosphatidylinositol 5-phosphate 4-kinase beta -/- mice", MOLECULAR AND CELLULAR BIOLOGY, vol. 24, no. 11, 1 June 2004 (2004-06-01), pages 5080-5087, XP002626464, Chicago IL USA**

EP 2 311 477 B1

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

TECHNICAL FIELD

5 **[0001]** The present invention relates to the provision of *in vitro* methods to control phosphatidylinositol metabolism in which PIPKII is involved. This is achieved by the using inositol 1,4,5-triphosphate receptor binding protein (IRBIT), DNA encoding IRBIT, siRNA for IRBIT or antibodies against IRBIT or IRBIT specific Fv, Fab, F(ab'), and scFv fragments thereof.

BACKGROUND OF THE INVENTION

10 **[0002]** When phosphatidylinositol 4,5-bisphosphate (PIP₂) is hydrolyzed through the activation of a receptor on a cell membrane, inositol 1,4,5-trisphosphate (IP₃), which is an intracellular second messenger, is generated. IP₃ binds to an IP₃ receptor (IP₃R), thereby inducing Ca²⁺ release from organelles for intracellular calcium storage (mainly, the endoplasmic reticulum). In this IP₃/Ca²⁺ signaling pathway, the IP₃ receptor plays a role in converting the IP₃ signal into Ca²⁺ signal (M. J. Berridge, Nature (1993) 361: 315-325; M. J. Berridge et al., Nat. Rev. Mol. Cell Biol. (2000) 1: 11-21; T. Furuichi and K. Mikoshiba, J. Neurochem. (1995) 64: 953-960).

15 **[0003]** The IP₃ receptor is a tetrameric intracellular IP₃-gated Ca²⁺ release channel. In mammals, there exist 3 different types of IP₃ receptors (i.e., type 1, type 2, and type 3) (T. Furuichi et al., Nature (1989) 342: 32-38; T. Sudhof et al., EMBO J. (1991) 10: 3199-3206; O. Blondel et al., J. Biol. Chem. (1993) 268: 11356-11363). Of them, the type 1 IP₃ receptor (IP₃R1) is expressed at high levels in the central nervous system and particularly in the cerebellum (P. F. Worley et al., Nature (1987) 325: 159-161; T. Furuichi et al., Recept. Channels (1993) 1: 11-24). Mouse IP₃R1 comprises 2749 amino acids and has 3 functionally different regions. Specifically, an IP₃-binding domain is present in the vicinity of the N-terminus, a channel-forming domain having a six-transmembrane region is present in the vicinity of the C terminus, and a control region is present between the two regions. The deletion mutant analysis of the IP₃-binding domain revealed that the amino acids 226-578 of the IP₃ receptor was a minimum region required for specific and high-affinity binding of a ligand. This region is referred to as the IP₃ binding core.

25 **[0004]** With an increase of cytoplasmic Ca²⁺ concentration by activation of the IP₃ receptor, the activities of a wide variety of downstream target molecules are controlled. These downstream target molecules play important roles in wide-ranging cellular responses including fertilization, development, proliferation, secretion, synaptic plasticity, and the like.

30 **[0005]** The present inventors have previously discovered a novel IP₃ receptor-binding protein and named it "IRBIT" (IP₃R-binding protein released with inositol 1,4,5-trisphosphate) (JP2004-129612A). The IP₃ receptor is widely distributed in various tissues and cells of mammals such as humans and mice (e.g., in the brain, heart, liver, kidney, pancreas, and thymus gland). Accordingly, IRBIT is inferred to be present also in such tissues or cells. The amino acid and nucleotide sequences of mouse IRBIT¹ have been determined by the present inventors (JP2004-129612A, H. Ando et al., J. Biol. Chem. (2003) 278: 10602-10612). Such IRBIT comprises 530 amino acids. Human IRBIT and mouse IRBIT share 100% identity. The region for binding to the IP₃ receptor is present in the N-terminal region of IRBIT, corresponding to amino acids 1-104 in a human or a mouse.

35 **[0006]** IRBIT is characterized in that: (1) IRBIT is a neutral protein (presumed pI: 6.48) in which the N-terminal region is relatively acidic (presumed pI: 4.98); (2) a plurality of phosphorylation sites are localized in a concentrated manner in the N-terminal region, so that phosphorylation is predicted to be necessary for interaction with IP₃R1; (3) the lysine residue at position 508, which is essential for the binding of IP₃R1 to IP₃, is also essential for interaction with IRBIT; (4) IRBIT is dissociated by IP₃ from interaction with IP₃R1; and (5) because IRBIT is dissociated from IP₃R1 and is extracted from crude microsome fractions by high salt, it is inferred that its interaction with IP₃R1 takes place due to electrostatic binding, for example (JP2004-129612A).

40 **[0007]** IRBIT has the property that it binds to the IP₃ binding region of the IP₃ receptor and is dissociated *in vitro* from the IP₃ receptor by IP₃. Therefore, it has also been revealed that IRBIT has a function of suppressing the activity of the IP₃ receptor by suppressing the binding of IP₃ to the IP₃ receptor (JP2004-129612A).

45 **[0008]** The present inventors have now found target molecules of IRBIT and important biological *in vivo* functions of IRBIT as a tertiary messenger, as described below.

Summary of the Invention

50 **[0009]** The present invention discloses the following.

[0010] In a first aspect, the present invention provides use of inositol 1, 4, 5-triphosphate (IP₃) receptor-binding protein (IRBIT) for controlling in-vitro In a cell phosphatidylinositol metabolism in which PIP₂ synthetase α , β or γ (PIPKII) is involved, wherein the IRBIT is a protein comprising an amino acid sequence shown in SEQ ID NO: 1 or SEQ ID NO:3 or a protein comprising an amino acid sequence having 90% or more identity with the amino acid sequence of SEQ ID NO: 1 or SEQ ID NO: 3 and having the same biological activity as IRBIT.

[0011] In another aspect, the present invention provides use of a DNA encoding IRBIT or vector comprising the DNA for controlling in-vitro in a cell phosphatidylinositol metabolism in which PIP₂ synthetase α , β or γ (PIPKII) is involved, wherein the IRBIT is a protein comprising an amino acid sequence shown in SEQ ID NO: 1 or SEQ ID NO: 3 or a protein comprising an amino acid sequence having 90% or more identity with the amino acid sequence of SEQ ID NO: 1 or SEQ ID NO: 3 and having the same biological activity as IRBIT, and wherein the nucleic acid is selected from the group consisting of DNAs encoding IRBIT.

[0012] In another aspect, the present invention provides use of siRNA for IRBIT for controlling in-vitro in a cell phosphatidylinositol metabolism in which PIP₂ synthetase α , β or γ (PIPKII) is involved, wherein the IRBIT is a protein comprising an amino acid sequence shown in SEQ ID NO: 1 or SEQ ID NO: 3 or a protein comprising an amino acid sequence having 90% or more identity with the amino acid sequence of SEQ ID NO: 1 or SEQ ID NO: 3 and having the same biological activity as IRBIT.

[0013] In another aspect, the present invention provides use of an antibody against IRBIT or a fragment thereof, capable of binding specifically to IRBIT, for controlling in-vitro in a cell phosphatidylinositol metabolism in which PIP₂ synthetase α , β or γ (PIPKII) is involved, wherein the IRBIT is a protein comprising an amino acid sequence shown in SEQ ID NO: 1 or SEQ ID NO: 3 or a protein comprising an amino acid sequence having 90% or more identity with the amino acid sequence of SEQ ID NO: 1 or SEQ ID NO: 3 and having the same biological activity as IRBIT, and wherein the fragment is selected from the group consisting of Fab, F(ab')₂, Fv and scFv.

[0014] In one embodiment, the control is suppression or elevation.

[0015] In another embodiment, the IRBIT is derived from a human or a mouse.

[0016] In another embodiment, the antibody is a monoclonal antibody, a polyclonal antibody, a recombinant antibody, or a single chain antibody.

[0017] In another embodiment, the IRBIT is phosphorylated.

Definitions

[0018] The term "IRBIT" as used herein refers to an IP₃ receptor-binding protein derived from a mammal, which binds to an IP₃ binding site of the IP₃ receptor and is released into cytoplasm when IP₃ binds to the receptor. As described herein, IRBIT binds to CPSF (cleavage/polyadenylation specificity factor), PIPKII (Type II phosphatidylinositol-5-phosphate 4-kinase), or pNBC1 (pancreas-type Na/HCO₃ cotransporter 1). These IRBIT targeting proteins exert important biological functions, which are involved in the protein synthesis, phosphatidylinositol metabolism, and intracellular pH maintenance respectively, through their bindings to IRBIT within cells. Thus, the IRBIT is responsible for controlling each of the biological functions.

[0019] The term "protein synthesis" as used herein refers to a series of intracellular gene transcription and translation processes. In the present invention, the control of mRNA polyadenylation in cytoplasm is involved in the protein synthesis. Moreover, the term "phosphatidylinositol metabolism" refers to the metabolism of phospholipids including IP₃.

[0020] The term "suppression" as used herein refers to decrease, reduction, or inhibition of the above-mentioned biological functions.

[0021] The term "elevation" as used herein refers to increase, rise, or enhancement of the above-mentioned biological functions.

[0022] The term "*ex vivo*" as used herein refers to a case in which cells or tissue removed from a living body are treated with the composition of the present invention and then returned into the living body.

[0023] The term "patient" as used herein refers to a mammal such as human, mouse, rat, dog, cat, or domestic animal (e.g., cattle, horse, pig, sheep, or goat), preferably human.

[0024] This specification includes all or part of the contents as disclosed in the description and/or drawings of Japanese Patent Application No. 2006-77607, to which the present application claims priority.

BRIEF DESCRIPTION OF THE DRAWINGS

[0025]

Fig. 1 shows the results of Western blotting showing the binding of IRBIT to CPSF160. In Fig. 1, "I.B." denotes Western blotting, "I.P." denotes immunoprecipitation, "HA" denotes hemagglutinin as a tag, and "load" denotes a cell extract.

Fig. 2 is a schematic diagram showing that IRBIT binds to an mRNA binding site of a CPSF160 subunit of CPSF.

Fig. 3 is a schematic diagram showing the phosphatidylinositol metabolism in which an IP₃ receptor, IRBIT, PIPKII, and IP₃ are involved. In Fig. 3, "PLC" denotes phospholipase C.

Fig. 4 shows the binding of mouse IRBIT to Myc-PIPKII α , β , and γ . Fig. 4A shows immunoprecipitation with an anti-Myc antibody. Fig. 4B shows immunoprecipitation with an anti-IRBIT antibody. In Fig. 4, "Input" denotes a cell extract,

"IP" denotes immunoprecipitation, "Control IgG (negative control)" denotes a sample subjected to immunoprecipitation using the control IgG.

Fig. 5 shows coprecipitation of mouse cerebellum PIPKII α with an anti-mouse IRBIT antibody by immunoprecipitation. "Input" is as described in Fig. 4.

Fig. 6 shows that both IRBIT and Myc-PIPKII γ are localized in the cytoplasm. "Merge" denotes merged images of a stained PIPKII image and a stained IRBIT image. Merged portions are observed yellow.

Fig. 7 shows identification of the binding site of IRBIT and PIPKII using IRBIT deletion mutants. Fig. 7A is a schematic diagram of IRBIT deletion mutants. Fig. 7B shows the binding of deletion mutants obtained by deletion from an N-terminal side of IRBIT, to Myc-PIPKII α . Fig. 7C shows the binding of deletion mutants obtained by deletion from the C-terminal side of IRBIT, to Myc-PIPKII α .

Fig. 8 shows that Ser68 and Ser71 of IRBIT bind to PIPKII. In Fig.8, "Input" denotes a cell extract, "IP" denotes immunoprecipitation, and "IB" denotes Western blotting.

Fig. 9A shows the results of Western blotting (top panel) showing the binding of IRBIT and NBC1. IRBIT (black triangle (◄)) binds to only recombinant proteins 2 and 5 containing portions specific to pNBC1 (pancreas-type). In the SDS polyacrylamide gel electrophoretic image (bottom panel of Fig. 9A), each recombinant protein (1, 2, 3, 4, 5, 6, or 7) is denoted with a black circle (●). Furthermore, Fig. 9B schematically shows the structures of these recombinant proteins in addition to the structures of pNBC1 and kNBC1 (kidney-type). In Fig. 9, "MBP" denotes a maltose-binding protein tag used for purification and "CBB" denotes coomassie brilliant blue.

Fig. 10 shows the ability of binding between NBC1 deletion mutants and IRBIT. Fig. 10A shows the results of causing expression of pNBC1 deletion mutants separately in *Escherichia coli*, purifying the resultants, and then examining the binding (or interaction) with HA-IRBIT (where "HA" denotes hemagglutinin) forcedly expressed in COS7 cells via pull down assay. Furthermore, the top panel in Fig. 10B shows the results of subjecting pulled down samples to SDS-PAGE and then staining the samples with CBB. The bottom panel in Fig. 10B shows the results of subjecting pulled down samples to Western blotting using an anti-HA antibody. Black triangles (◄) indicate electrophoretic mobilities of HA-IRBIT and black dots (•) indicate the electrophoretic mobilities of deletion mutant proteins.

Fig. 11 shows the ability of binding between IRBIT deletion mutants and pNBC1. Each type of IRBIT deletion mutant was forcedly expressed in COS7 cells in the form of fusion protein fused with GFP (green fluorescent protein) and then the binding to pNBC1 was examined via pull down assay. Both Fig. 11A and Fig. 11B show the results of subjecting cell extracts (Lysate) or pulled down samples (Pull down) each expressing a deletion mutant to SDS-PAGE and then to Western blotting using an anti-GFP antibody. Black triangle (◄) indicates an electrophoretic mobility of each deletion mutant. It was understood that all IRBITs other than the full-length (FL) had become unable to bind to pNBC1.

Fig. 12 shows the binding of endogenous IRBIT and NBC1 in cerebellum membrane fractions. A cerebellum membrane fraction extract was subjected to immunoprecipitation (IP) using a preimmunization antibody as a control and an anti-NBC1 antibody, or an anti-IRBIT antibody. Precipitates were subjected to SDS-PAGE and then Western blotting (IB) using an anti-IRBIT antibody, an anti-NBC1 antibody, or an anti-IP₃ receptor antibody was performed. Black triangle (◄) indicates an electrophoretic mobility of each protein.

Fig. 13 shows the binding of endogenous IRBIT and NBC1 in COS7 cell extracts. The COS7 cell extract was subjected to immunoprecipitation (IP) using a preimmunization antibody (control), an anti-IRBIT antibody, or an anti-NBC1 antibody. Precipitates were subjected to SDS-PAGE and then Western blotting (IB) using an anti-NBC1 antibody or an anti-IRBIT antibody was performed. Black triangle (◄) indicates an electrophoretic mobility of each protein. Three rows on the left show the results obtained using precipitates from immunoprecipitation in the absence of CaCl₂, and three rows on the right show the results obtained using precipitates from immunoprecipitation in the presence of CaCl₂.

Fig. 14 shows the activation of pNBC1 by IRBIT, as measured by a voltage-clamp method. Fig. 14A shows current values (μ A) obtained when the membrane potential was fixed at -25 mV. Furthermore, Fig. 14B shows I-V curves that represent current changes when the membrane potential was varied between -160 mV and +60 mV.

Fig. 15 shows that phosphorylation of IRBIT is required for activation of pNBC1.

PREFERRED EMBODIMENTS OF THE INVENTION

[0026] Hereinafter, the present invention will be described in more detail.

1. Composition

[0027] Although not in accordance with the present invention as claimed, there is described herein a composition comprising IRBIT, a nucleic acid controlling the expression and translation of IRBIT, and an antibody against IRBIT. The composition may be used for controlling at least one intracellular biological function selected from the group consisting

of (1) protein synthesis, (2) phosphatidylinositol metabolism, and (3) intracellular pH.

[0028] Control of the above three biological functions will be described as follows.

Control of protein synthesis

[0029] Although not in accordance with the present invention as claimed, the above control of protein synthesis is the control of mRNA polyadenylation mediated by intracellular binding of IRBIT with CPSF.

[0030] CPSF is a conjugated protein consisting of four subunits, CPSF160, CPSF100, CPSF73, and CPSF30. The present inventors have discovered this time that IRBIT binds to CPSF and particularly binds to CPSF160 as a result of coprecipitation tests involving coexpression in COS cells and immunoprecipitation (Fig. 1).

[0031] CPSF is a molecule essential for intranuclear mRNA polyadenylation reactions and is also known to have a function of regulating protein synthesis via extension of poly(A) length in the cytoplasm. Furthermore, the mRNA binding site of CPSF160 is known to be essential for CPSF to recognize mRNA to which poly(A) is added, for example (C. Barnard Daron et al., *Cell* 2004, 119: 641-651, and E. Klann et al., "Synaptic Plasticity and Translation Initiation," *Learning & Memory* 2004, 11: 365-372, Cold Spring Harbor Laboratory Press). Specifically, a section in "Cytoplasmic polyadenylation and CPEB" (E. Klann et al., (mentioned above), pp. 367 to 368) discloses that: polyadenylation is regulated by two sequences in the 3' non-translated region of mRNA; that is, cytoplasmic polyadenylation element (CPE) and AAUAAA; CPE binding protein (CPEB), which is an important regulatory protein of polyadenylation, is phosphorylated by specific protein kinase (Aurora); such kinase phosphorylates CPEB, so that CPSF interacts with CPEB on the AAUAAA sequence, poly(A) polymerase (PAP) is recruited, and a poly(A) tail portion of mRNA is extended.

[0032] In view of the above findings, it is considered that IRBIT binds to the mRNA binding site of CPSF160, so that CPSF functions are controlled (Fig. 2).

[0033] Furthermore, the present inventors have obtained this time a finding that IRBIT has a function of further enhancing polyadenylation activity (I. Kaufmann et al., *EMBO J.* (2004) 23: 616-626) in the presence of PAP and Fip1 (CPSF subunit). Based on the finding, the suppression of protein synthesis is made possible by suppression of IRBIT.

[0034] As described above, IRBIT is involved in control of mRNA polyadenylation and thus involved in control of protein synthesis through its binding to CPSF.

[0035] Therefore, IRBIT, or a substance that suppresses or enhances the generation and functions of IRBIT, makes it possible to control CPSF-associated protein synthesis

Control of phosphatidylinositol metabolism

[0036] IRBIT further binds to PIPKII, so as to suppress the activity of the enzyme (Example 2).

[0037] PIPKII is an enzyme for synthesis of phosphatidylinositol 4,5-diphosphate (PIP₂) from phosphatidylinositol pentaphosphate (PIP). Furthermore, IP₃ is produced via hydrolysis of PIP₂. IP₃ is a ligand of an IP₃ receptor. IP₃ binds to the receptor so that calcium ion (Ca²⁺) and IRBIT are together released within the cytoplasm. When the fact is taken into consideration, it can be said that IRBIT is involved in control of phosphatidylinositol metabolism. Katia A. Lamia et al., *Mol. Cell Biol.* 2004, 24: 5080-5087 disclose that inhibitors of the beta-isoform of PIPKII may be useful to treat type 2 diabetes. EP 1 408 049 A discloses that IRBIT interacts with PIPKII.

[0038] Therefore, IRBIT, or a substance that suppresses or enhances the generation and functions of IRBIT, may make it possible to control PIPKII-associated phosphatidylinositol metabolism (Fig. 3).

[0039] For example, in mammals, PIPKII encompasses 3 types of isoforms notably PIPKII α , β , and γ , and IRBIT binds to all of these isoforms. Particularly with regard to PIPKII β , it has been demonstrated that inhibition of the enzyme is useful for treatment of type 2 diabetes (Katja A. Lamia et al., *ibid*) based on the fact that a transgenic mouse with a knocked-out gene corresponding to the enzyme has high insulin sensitivity. The finding of the present inventors that IRBIT suppresses PIPKII activity suggests that IRBIT can perhaps be used for treatment of type 2 diabetes.

Control of intracellular pH

[0040] IRBIT further binds to pNBC1, so as to activate pNBC1.

[0041] Specifically, IRBIT cRNA and NBC1 cRNA were injected into *Xenopus* oocytes. The degree of pNBC1 response detected in connection with intracellular pH change was approximately 6 to 7 times higher than the degree detected before pH change (Fig. 14). This result demonstrates that IRBIT significantly enhances pNBC1 activity.

[0042] NBC1 is a 10-transmembrane protein existing on the cell membrane and functions to transport sodium ions and bicarbonates at a constant rate in the same direction across the cell membrane. *In vivo* pH is cleverly regulated by balancing between bicarbonate concentration and carbon dioxide gas concentration. Thus, it is considered that NBC1 is involved in regulation of *in vivo* pH (E. Gross and I. Kurtz, *Am. J. Physiol. Renal Physiol.* 2002, 283: F876-F887). In particular, identification of NBC1 as a causative gene of proximal renal tubular acidosis, which is a type of acidaemia

due to which blood pH approaches acidic levels suggests that the transport of bicarbonates by NBC1 plays a role essential for *in vivo* pH maintenance.

[0043] Concerning NBC1, two splicing mutants have been reported: a kidney-type (kidney-type: kNBC1) and a pancreas-type (pancreas-type: pNBC1). kNBC1 is mainly expressed in the kidney and pNBC1 is mainly expressed in the pancreas and in relatively many tissues including the cerebral nervous system. To reveal to which one of NBC1s and to which part of NBC1 the IRBIT binds, the present inventors have caused the expression of the cytoplasmic region of NBC1 as a recombinant protein in *Escherichia coli*, purified the recombinant protein, and then examined the binding of the protein with IRBIT forcedly expressed in cultured cells by pull down assay. As a result, it was revealed that IRBIT specifically and strongly binds to pNBC1-specific N-terminal 85 amino acids (Fig. 10). Furthermore, it was also revealed that the binding between IRBIT and pNBC1 is controlled by changes in specific salt concentration.

[0044] These results strongly suggest the possibility that IRBIT controls pH regulation mediated by pNBC1 in various organs in a manner depending on intracellular conditions. Moreover, patients with proximal renal tubular acidosis develop eye diseases such as glaucoma and cataract or present various symptoms including dwarfism, mental retardation, pancreatitis, and the like. pH regulation conducted by NBC1 is considered to also play an important role in organs other than the kidney. Therefore, IRBIT is also useful through mediation of pNBC1 for methods for treating eye diseases such as glaucoma and cataract and diseases such as dwarfism, mental retardation, and pancreatitis (Seth L. Alper, *Annu. Rev. Physiol.* 2002, 64: 899-923).

[0045] Therefore, although not in accordance with the present invention as claimed, IRBIT, or a substance that suppresses or enhances the generation or functions of IRBIT, may make it possible to control pNBC1-associated intracellular pH.

IP₃ receptor-binding protein (IRBIT)

[0046] IRBIT to be used in the present invention is derived from a mammal. IRBIT is known to be present in the intracellular endoplasmic reticulum of tissues such as the brain, heart, liver, kidney, pancreas, and thymus gland of a mammal (JP2004-129612 A). Preferable examples of IRBIT include human IRBIT and mouse IRBIT (JP2004-129612 A, H. Ando et al., *J. Biol. Chem.* 2003, 278: 10602-10612). Particularly, human IRBIT is preferable. The amino acid and nucleotide sequences of human IRBIT and mouse IRBIT are deposited with the GenBank under NM_006621 (see SEQ ID NOS: 1 and 2) and NM_145542 (see SEQ ID NOS: 3 and 4), respectively.

[0047] Furthermore, another preferable example of IRBIT is a protein comprising an amino acid sequence having 90% or more, preferably 95% or more, more preferably 98% or more, and most preferably 99% or more identity with the amino acid sequence shown in SEQ ID NO: 1 or SEQ ID NO: 3 and having biological activity equivalent to that of IRBIT. As used herein, the term "biological activity" refers to, in addition to S-adenosylhomocysteine hydrolase-like activity that catalyzes reversible hydrolysis of S-adenosylhomocysteine into adenosine and homocysteine, activity involved in control of biological functions including the control of mRNA polyadenylation that is mediated via binding with CPSF in protein synthesis, the control of phosphatidylinositol metabolism that is mediated via binding with PIPKII, and the control of intracellular pH that is mediated via binding with pNBC1.

[0048] Similarly, a preferable example of DNA encoding IRBIT is DNA having 90% or more, preferably 95% or more, more preferably 98% or more, and most preferably 99% or more identity with the nucleotide sequence shown in SEQ ID NO: 2 or SEQ ID NO: 4, or, DNA capable of hybridizing under stringent conditions to the nucleotide sequence of SEQ ID NO: 2 or SEQ ID NO: 4. Here, the stringent conditions consist of, but are not limited to, hybridization at approximately 45-50°C in 2-6xSSC (sodium chloride/sodium citrate), followed by washing at approximately 50-65°C with 0.2-2xSSC/0.1-1% SDS, or hybridization at 60-65°C in 6xSSC, Denhard't solution, and 0.2% SDS, followed by washing at 60-65°C with 0.2xSSC and 0.1% SDS (e.g., F. M. Ausbel et al., *Short Protocols in Molecular Biology* (3rd edition) A Compendium of Methods from Current Protocols in Molecular Biology, 1995, John Wiley & Sons, Inc.).

[0049] IRBITs derived from other mammals can also be used in the present invention. Examples of such IRBITs include IRBIT derived from an experimental animal such as a rat, a hamster, and a rabbit, IRBIT derived from a pet animal such as a dog and a cat, and IRBIT derived from a domestic animal such as cattle, a horse, a pig, sheep, and a goat. These IRBITs are prepared as follows. Probes and/or primers are prepared based on an amino acid or nucleotide sequence described in a document, databank, or the like or based on a known sequence of human or mouse IRBIT. IRBIT cDNA is cloned and/or amplified by a commonly employed technique such as a DNA cloning method or polymerase chain reaction (PCR) using a commercial library or a library constructed from the prepared animal tissues, for example. Furthermore, the thus obtained DNA encoding IRBIT is incorporated into a commercial expression vector (e.g., plasmid) having an appropriate regulatory sequence, for example. The expression vector is transformed into appropriate host cells or host cells are transfected with the expression vector. The thus obtained cells are cultured in appropriate medium, so as to cause expression of IRBIT DNA. The thus generated IRBIT protein can be collected. A series of these techniques are described in, for example, J. Sambrook et al., *Molecular Cloning A Laboratory Manual*, 1989, Cold Spring Harbor Laboratory Press, F. M. Ausbel et al., *Short Protocols in Molecular Biology* (3rd edition) A Compendium of Methods from

Current Protocols in Molecular Biology, 1995, John Wiley & Sons, Inc., Experimental Medicine, Separate Volume, 4th edition, Edited by Masami Matsumura et al., "New Genetic Engineering Handbook" (2003) YODOSHA, Tokyo, Japan, and the like. IRBIT homologs derived from various mammals can be obtained according to such techniques described in these documents.

5 [0050] A mammalian tissue containing an IRBIT gene is homogenized using a homogenizer and then centrifuged at approximately 10,000 rpm, so that a supernatant is obtained. Subsequently, total RNA is collected by a guanidine acidic phenol method, for example, cDNA is synthesized according to a standard method, and then DNA encoding IRBIT can be obtained from the cDNA. For example, ISOGEN (trademark) of NIPPON GENE is commercially available as a kit for RNA extraction and can also be used herein.

10 [0051] The size of a probe for detection of DNA encoding IRBIT is generally 30 or more nucleotides and preferably 50 to 100 or more nucleotides. In general, a label such as a fluorescent label (e.g., fluorescamine, rhodamine, or their derivatives thereof), a radioactive isotope label (e.g., ³²P), or the like is bound to a probe. Thus, the binding of DNA encoding target IRBIT with the probe can be detected.

15 [0052] The size of a primer for amplification of DNA encoding IRBIT generally ranges from 15 to 30 nucleotides and preferably ranges from 20 to 25 nucleotides. Primers to be used herein should have sequences complementary to the 3' terminal sequences of the sense strand and the antisense strand of DNA encoding IRBIT. However, when the sequence of IRBIT-encoding DNA to be amplified is unknown, a plurality of primers are prepared based on known IRBIT sequences, a template DNA is amplified by PCR, and then formal primers are prepared based on the thus amplified template DNA, so that target template DNA can be amplified by PCR.

20 [0053] PCR is generally performed by performing approximately 20 to 40 cycles each consisting of DNA denaturation, primer annealing, and elongation reaction. DNA denaturation is a step for separating double-stranded DNA into single-chain DNAs, which is performed by approximately 15 seconds to 1 minute of treatment generally at 94°C. Primer annealing is a step for annealing a primer to complementary single-chain template DNA. The optimum temperature or time for annealing depends on the nucleotide sequence of a primer or the length thereof. In general, annealing treatment is performed at approximately 55-60°C for approximately 30 seconds to 1 minute. Elongation reaction is a step for elongation of template DNA in the presence of 4 types of dNTP and heat-resistant DNA polymerase. In general, elongation reaction is performed at 72°C for approximately 30 seconds to 10 minutes. Before the initiation of the cycles, DNA can be completely denatured by heating at 94°C for approximately 1 to 5 minutes. Furthermore, after completion of the entire cycles, approximately 1 to 5 minutes of heat treatment can also be performed at 72°C. Heat-resistant DNA polymerase is commercially available and *Thermus aquaticus* (Taq) polymerase (marketed by TaKaRa, PerkinElmer, Pharmacia, or the like) can be used, for example. Concerning PCR techniques, Protein, Nucleic Acid, and Enzyme "Frontiers of PCR from Basic Technology to Application," vol. 41, No. 5, April 1996, Extra Number, KYORITSU SHUPPAN, Tokyo, Japan can be referred, for example.

30 [0054] An expression vector to be used herein may be any vector that can be used in prokaryote- or eukaryote-derived cells. A vector can contain regulatory sequences such as a promoter, a replication origin, a ribosome-binding site, a multicloning site, and a terminator. As expression vectors, plasmid or viral vectors, or the like, and particularly commercially-available vectors such as pGEX-4T-1 (Amersham Pharmacia Biotech), pBluescript II SK, pHS 19, pHS 15, pG-1, and pXT1 (Stratagene Corporation), pMAL and pTYB series (Daiichi Pure Chemicals), pQE series (Qiagen), pET series (Novagen), pSVK3 and pSVL SV40 (Pharmacia), pcDNA1 and pcDM8 (Funakoshi), and pHB6, pVB6, pHM6, pVM6, and pXM (Roche Diagnostics) can be adequately selected and used.

35 [0055] Examples of host cells include bacteria of the genus *Escherichia*, such as *Escherichia coli*, bacteria of the genus *Bacillus*, such as *Bacillus subtilis*, and bacteria of the genus *Pseudomonas*, bacteria of the genus *Corynebacterium*, yeasts such as yeast of the genus *Saccharomyces*, yeast of the genus *Pichia*, and yeast of the genus *Schizosaccharomyces*, insect cells, plant cells, and mammalian cells (e.g., CHO, COS, and HEK293 cells).

40 [0056] Examples of a method for introducing DNA encoding IRBIT into host cells include a calcium phosphate method, a lipofection method, an electroporation method, and a method using infection with viruses such as adenovirus or retrovirus (Experimental Medicine, Separate Volume, 4th Edition, Edited by Masami Matsumura et al., "New Genetic Engineering Handbook (2003) YODOSHA, Tokyo, Japan).

45 [0057] More specifically, concerning mouse IRBIT, IRBIT purification from mature mouse cerebellum and cDNA cloning and expression are disclosed in JP2004-129612 A. The disclosure is available for reference.

50 [0058] The biological activity of IRBIT can be determined based on an analogous assay method since IRBIT has homology with S-adenosylhomocysteine hydrolase that catalyzes reversible hydrolysis of S-adenosylhomocysteine into adenosine and homocysteine. This assay method can be performed according to the method of C. -S. Yuan et al., (J. Biol. Chem. 1996, 271: 28009-28016), for example. This is briefly explained as follows. This assay method comprises performing the above hydrolysis reaction using IRBIT (approximately 3 μg) and rabbit S-adenosylhomocysteine hydrolase (Sigma) (approximately 2.5 μg), causing the thus generated product (homocysteine) to react with 5,5'-dithiobis (2-nitrobenzoate) (Sigma), measuring the thus obtained color at 412 nm using a spectrophotometer, and then finding the absorbance.

Nucleic acid controlling IRBIT expression and translation

[0059] In the present invention, examples of nucleic acids that control IRBIT expression and translation include DNA encoding IRBIT, antisense RNA of mRNA encoding IRBIT or a fragment thereof, ribozyme that makes it possible to cleave mRNA encoding IRBIT, functional RNA such as siRNA (small interfering RNA), and vector DNA containing such DNA or RNA (actually, DNA encoding RNA).

[0060] DNA encoding IRBIT is derived from a mammal such as a human or a mouse and can be prepared by a technique as explained in the above section, "IP₃ receptor-binding protein (IRBIT)." Preferable DNA encoding IRBIT is DNA comprising the nucleotide sequence that encodes the amino acid sequence shown in SEQ ID NO: 1 or SEQ ID NO: 3. More preferable DNA is DNA comprising the nucleotide sequence shown in SEQ ID NO: 2 or SEQ ID NO: 4 and encoding human or mouse IRBIT.

[0061] DNA encoding IRBIT is inserted into an expression vector (e.g., a plasmid or viral vector) and thus can be used for the intracellular expression of IRBIT.

[0062] A plasmid vector for expression of DNA encoding IRBIT can contain, in addition to a DNA sequence encoding IRBIT and a promoter, regulatory sequences such as a drug resistance gene (e.g., a neomycin resistance gene, an ampicillin resistance gene, a puromycin resistance gene, and a hygromycin resistance gene), a terminator, a multiple cloning site, a replication origin, and a ribosome-binding site.

[0063] As a viral vector for expression of DNA encoding IRBIT, an adenoviral vector, an adeno-associated viral vector, a lentiviral vector, a retroviral vector (e.g., leukemia viral vector), a herpes viral vector, or the like can be used, for example. A preferable type of a viral vector is deficient in replication competence, for example, so as not to cause disease when the vector is applied to a human. For example, in the case of an adenoviral vector, a replication competence-deficient adenoviral vector (e.g., pAdeno-X (Invitrogen)) prepared by deletion of E1 and E3 genes can be used. A method described in the document can be used for construction of such a viral vector (e.g., U.S. Patent No. 5252479 and International Publication WO94/13788).

[0064] As a promoter, a promoter that enables expression of foreign DNA within mammalian cells can be used. Examples of such promoter include a cytomegalovirus (CMV) promoter, a SV40 promoter, and an EF promoter.

[0065] In the case of a plasmid vector, a complex of such a plasmid vector with positively charged liposome such as Lipofectamine, Lipofectin, CellFECTIN, or positively charged cholesterol is formed, the complex is capsulated, and then the resultant can be introduced into a living patient's body (e.g., Mamoru Nakanishi et al., Protein, Nucleic Acid, and Enzyme, Vol. 44, No. 11, pp. 48-54, 1999, KYORITSU SHUPPAN, Tokyo, Japan; Clinical Cancer research 59: 4325-4333, 1999; Wu et al., J. Biol. Chem. 1987, 262: 4429). Furthermore, in the case of a viral vector, such a viral vector is introduced into an affected part for infecting cells, so that gene transfer into cells can be performed (L. Zender et al., Proc. Natl. Acad. Sci. U.S.A. (2003), 100: 77797-7802; H. Xia et al., Nature Biotech. (2002), 20: 1006-1010; X. F. Qin et al., Proc. Natl. Acad. Sci. U.S.A. (2003), 100: 183-188; G. M. Barton et al., Proc. Natl. Acad. Sci. U.S.A. (2002), 99: 14943-14945; J. D. Hommel et al., Nature Med. (2003), 9: 1539-1544). In particular, it has been confirmed that gene transfer can be performed with very high efficiencies into various cell species with the use of an adenoviral vector or an adeno-associated viral vector. Such vectors are not incorporated into the genome, so that their effects are exerted temporarily and the safety of such vectors is thought to be much higher than other viral vectors.

[0066] Moreover, antisense RNA of mRNA encoding IRBIT or a fragment thereof can inhibit the translation of an IRBIT gene to an IRBIT protein.

[0067] The above fragment can comprise a sequence consisting of the number of nucleotides, which is approximately continuous 30 or more nucleotides, 50 or more nucleotides, 70 or more nucleotides, 100 or more nucleotides, 150 or more nucleotides, 200 or more nucleotides, or 250 or more nucleotides, but is the same or less than the full-length sequence of an IRBIT gene or mRNA.

[0068] Antisense RNA or a fragment thereof may comprise halogen (fluorine, chlorine, bromine or iodine) and a modification group such as a methyl, carboxymethyl, or thio group.

[0069] The above antisense nucleic acid can be synthesized using a known DNA/RNA synthesis technique or a DNA recombination technique. When such antisense nucleic acid is synthesized by a DNA recombination technique, polymerase chain reaction (PCR) is performed using vector DNA containing the nucleotide sequence of IRBIT as a template and primers that sandwich a sequence to be amplified, so as to amplify the target sequence. If necessary, the product is cloned into a vector, and then antisense DNA can be generated. Alternatively, DNA having the thus obtained amplified target sequence is inserted into a vector, and then the vector is introduced into eukaryotic or prokaryotic cells, so that antisense RNA can be obtained with the use of the transcription system.

[0070] Similar inhibition can also be performed using ribozyme that enables cleavage of mRNA encoding IRBIT or functional RNA such as siRNA.

[0071] siRNA can contain a sense strand sequence (derived from mRNA sequence that is encoded by the nucleotide sequence of SEQ ID NO: 2 (human IRBIT) or SEQ ID NO: 4 (mouse IRBIT), for example) with a length of approximately continuous 18 to 30, preferably approximately 19 to 25, and further preferably approximately 20 to 23 nucleotides and

a complementary sequence thereof that is an antisense strand sequence. Here, the term "sense strand sequence" refers to the same nucleotide sequence as that of a target site of the above mRNA. Furthermore, the term "antisense strand sequence" refers to a nucleotide sequence complementary to the sense strand sequence. A sense strand and antisense strand can anneal together to form double-stranded siRNA. siRNA can further comprise an overhang (e.g., UU) consisting of 1 to 5 nucleotides at each 3' end of sense and antisense strands.

[0072] For selection of a sense strand sequence of siRNA, known understandings for selection of a target site of target IRBIT mRNA can be employed. For example, criteria that can be employed are: that (a) GC content ranges from approximately 30% to 70% and is preferably approximately 50%, (b) all nucleotides are equal and G is discontinuous, (c) the 5' terminal nucleotides of an antisense strand are A and U (D. M. Dykxhoorn et al., *Nature Rev. Mol. Cell Biol.* (2003), 77: 7174-7181; A. Khvorova et al., *Cell* (2003), 115: 209-216). Furthermore, a candidate gene site on IRBIT mRNA that is a target site of the above siRNA can be inferred using the mfold RNA secondary structure prediction program (J. A. Jaeger et al., *Methods in Enzymology* 1989, 183: 281-306; D. H. Mathews et al., *J. Mol. Biol.* 1999, 288: 911-940). For example, a sequence that siRNA can target can be determined by inferring based on the above findings and then actually confirming the effects. Examples of siRNA that can be used in the present invention include, but are not limited to, the following sequences.

IRBIT siRNA-1: AAAUCCAGUUUGCUGAUGACA (SEQ ID NO: 5)

IRBIT siRNA-2: AACUCAGAAUGAAGUAGCUGC (SEQ ID NO: 6)

[0073] siRNA can be synthesized using a known chemical synthesis technique. For example, siRNA can be obtained via chemical synthesis using a conventionally used DNA/RNA autosynthesizer or by commissioning the synthesis thereof to a company handling the synthesis of siRNA and the like (e.g., Funakoshi (Tokyo, Japan), Dharmacon, or Ambion).

[0074] When siRNA is introduced into cells or tissues, siRNA is directly injected into cells or tissues or a vector that enables siRNA expression is preferably used. Alternatively, a complex is formed using siRNA or a vector and liposome such as Lipofectamine, Lipofectin, CellFECTIN, or another positively-charged liposome (e.g., positively charged cholesterol) or a microcapsule and then the complex can also be used (e.g., Mamoru Nakanishi et al., *Protein, Nucleic Acid, and Enzyme*, Vol. 44, No. 11, pp. 48-54, 1999, KYORITSU SHUPPAN, Tokyo, Japan; *Clinical Cancer research* 59: 4325-4333, 1999; Wu et al., *J. Biol. Chem.* 1987, 262: 4429).

[0075] A vector for expression of siRNA contains a DNA sequence encoding siRNA or a precursor thereof under regulation of a promoter.

[0076] An example of an expression vector is a hairpin vector. This type of vector contains DNA encoding hairpin RNA in which the above sense strand RNA sequence and the above antisense strand RNA sequence are covalently bound via a single-chain loop sequence, wherein the DNA is a vector such that the hairpin RNA is formed via intracellular transcription and then the hairpin RNA is processed by a dicer to form the above siRNA. A poly T sequence comprising 1 to 6 and preferably 1 to 5 Ts is ligated to the 3' end of hairpin DNA encoding siRNA as a transcription termination signal sequence or for overhanging. It is desired that short hairpin RNA (shRNA) as an siRNA precursor transcribed from vector DNA have an overhang comprising 2 to 4 Us at the 3' end of the antisense strand. Because of the presence of such overhang, sense strand RNA and antisense strand RNA can increase their stability against degradation by nuclease. One endogenous dicer is present in a human and is responsible for converting long-chain dsRNA or precursor micro RNA (miRNA) into siRNA and mature miRNA, respectively. Examples of a promoter include a pol III promoter such as human- or mouse-derived U6 promoter, and an H1 promoter, a pol II promoter, and a cytomegalovirus promoter.

[0077] Another example of an expression vector is a tandem vector. This vector contains a DNA sequence encoding a sense strand RNA sequence that composes the above siRNA and a DNA sequence encoding an antisense strand RNA sequence consecutively. The vector further contains DNA in which a promoter is ligated to the 5' end of each strand and a poly T sequence is ligated to the 3' end of each strand, wherein the DNA is a vector such that the sense strand RNA and the antisense strand RNA hybridize to each other after intracellular transcription, so as to form the above siRNA.

[0078] An example of a promoter for such tandem vector is a pol III promoter such as a human- or a mouse-derived U6 promoter or an H1 promoter, or a cytomegalovirus promoter. Furthermore, a poly T sequence is a poly T sequence comprising 1 to 6 and preferably 1 to 5 Ts. Such a tandem vector is introduced into cells and then transcribed into RNAs corresponding to a sense strand and an antisense strand. The strands hybridize to each other, so that a target siRNA can be generated.

[0079] The above hairpin and tandem vectors are plasmid vectors or viral vectors. A plasmid vector can be prepared using techniques described in the following Examples or methods described in documents. Alternatively, a commercially available vector system, such as piGENE™ U6 vector and piGENE™ H1 vector (TAKARA BIO INC., Kyoto, Japan) can also be used (T. R. Brummelkamp et al., *Science* (2002), 296: 550-553; N. S. Lee et al., *Nature Biotech.* (2002), 20: 500-505; M. Miyagishi et al., *Nat. Biotechnol.* (2002), 20: 497-500; P. J. Paddison et al., *Genes & Dev.* (2002), 16: 948-958; T. Tusch, *Nature Biotech* (2002), 20: 446-448; C. P. Paul et al., *Nature Biotech.* (2002), 20: 505-508; Edited by Kazumasa Tahira et al., *RNAi Experimental Protocols*, YODOSHA (Tokyo, Japan), 2003).

[0080] Another nucleic acid that is an active ingredient of the composition of the present invention is ribozyme. Ribozyme is RNA having catalytic activity and has activity of cleaving mRNA that corresponds to a target IRBIT gene of the present invention. The expression of the gene is inhibited or suppressed by this cleavage.

[0081] It is known that a cleavable target sequence of ribozyme is generally NUX (N = A, G, C, U; X = A, C, U), such as a sequence containing GUC triplet. Such ribozyme contains hammerhead-type ribozyme. The hammerhead-type ribozyme can contain a nucleotide sequence that composes a sensor site, a nucleotide sequence that contains a region capable of forming cavities that stably capture Mg²⁺ ion only when RNA binds to the sensor site, and a nucleotide sequence containing a region complementary to a sequence in the vicinity of the cleavage site of the target RNA.

[0082] For delivery of the ribozyme of the present invention into cells or into a patient's living body, ribozyme is enclosed within liposome (preferably, positively charged liposome) (JP Patent Publication (Kokai) No. 9-216825 A (1997)) and then the liposome is incorporated into a viral vector such as an Adeno-associated virus (JP Patent Publication (Kohyo) No. 2002-542805 A). With the use of these methods, a drug delivery system can be constructed.

[0083] Ribozyme can be incorporated into a vector so that it can be expressed. Examples of a promoter for expression of ribozyme include pol II and pol III promoters. A promoter is preferably a pol III promoter such as a mammal-derived tRNA promoter and is more preferably a tRNA^{Val} promoter (S. Koseki et al., J. Virol., 73: 1868-1877, 1999).

Antibody

[0084] An antibody against IRBIT or a fragment thereof can also be used for controlling the above biological functions.

[0085] Examples of such antibody include polyclonal antibodies, monoclonal antibodies, recombinant antibodies, human antibodies, humanized antibodies, chimeric antibodies, single-chain antibodies, Fab fragments, F(ab')₂ fragments, Fv, scFv, bispecific antibodies, and synthetic antibodies.

[0086] The class or subclass of an antibody may be any type. Examples of such class or subclass include IgG, IgM, IgE, IgD, IgA, IgG₁, IgG₂, IgG₃, IgG₄, IgA₁, and IgA₂.

[0087] An antibody to be used herein may also be derivatized via pegylation, acetylation, glycosylation, amidation, or the like.

[0088] For preparation of a polyclonal antibody, a water-in-oil emulsion containing IRBIT as an immunogen (approximately 1 μg to 100 μg) and if necessary an adjuvant such as Freund's complete or incomplete adjuvant, aluminium hydroxide (alum), muramyl dipeptide, or lipid A is injected intradermally or intravenously for immunization into non-human animals such as rabbits, guinea pigs, mice, rats, sheep, or goats. Approximately 2 to 4 weeks later, boost was performed via injection once or twice with adjuvant-free IRBIT. Blood was collected on a trial basis, so as to confirm that the antibody titer has increased sufficiently. Subsequently, blood was collected from the animals and then antiserum was collected by centrifugation. If necessary, purification is performed by ammonium sulfate fractionation, DEAE ion exchange chromatography, or the like, so that IgG can be obtained.

[0089] A hybridoma that secretes a monoclonal antibody can be prepared according to Kohler and Milstein's technique (Nature 1975, 256: 495-497). Specifically, a hybridoma can be obtained by extracting the spleen or the lymph node from an immunized animal, fusing antibody-producing cells contained therein to myeloma cells derived from a mammal such as a mouse, a rat, or a guinea pig, and then performing HAT selection. Cell fusion can be performed using polyethylene glycol (e.g., molecular weight ranging from 1500 to 6000), for example. For production of a target antibody, the reactivity of the hybridoma to the immunizing antigen in a culture supernatant can be measured using a conventional method such as enzyme immunoassay, radioimmunoassay, or a fluorescent antibody technique.

[0090] Furthermore, for preparation of a monoclonal antibody from a hybridoma, a hybridoma is cultured *in vitro* and then the monoclonal antibody may be isolated from the culture supernatant. Alternatively, a hybridoma is cultured *in vivo* in the ascites or the like of a mouse, a rat, a guinea pig, or the like and then the monoclonal antibody may be isolated from the ascites.

[0091] Alternatively, a gene that encodes a monoclonal antibody is cloned from antibody-producing cells such as hybridomas, and then the resultant is incorporated into a vector, and then the vector is introduced into mammalian cells (e.g., CHO), so that a recombinant antibody can also be prepared (P. J. Delves et al., ANTIBODY PRODUCTION ESSENTIAL TECHNIQUES., 1997, John Wiley&Sons).

[0092] A human antibody can be produced by a phage display library method (T. C. Thomas et al., Mol. Immunol. 33: 1389-1401, 1996) or a method using a human antibody-producing animal (e.g., a mouse or cattle) (I. Ishida et al., Cloning Stem Cell 4: 91-102, 2002), for example.

[0093] For example, a human antibody-producing mouse can be produced by a method that involves introducing a human chromosome fragment containing a human-antibody-producing gene into a human artificial chromosome, incorporating the artificial chromosome into a mouse embryonic stem cell genome or the like with the use of a microcell method, injecting the recombinant embryonic stem cell into a blastocyst, implanting the blastocyst in the uterus of a foster mother mouse, causing the mouse to deliver chimeric mice, and thus producing homozygous progeny mice that contains the human antibody gene and is capable of producing human antibodies through mating of male and female chimeric

mice or mating of chimeric mice with wild-type mice (e.g., Japanese Republication (Saikohyo) No. 02/092812 A (1990). International Publication WO 98/24893, and WO 96/34096). The human-antibody-producing transgenic mouse is immunized with the IRBIT protein of the present invention as an antigen, the spleen is excised, and then a hybridoma is formed by fusing the spleen cells with mouse myeloma cells, so that a target monoclonal antibody can be selected.

[0094] The phage display library method involves screening for DNA that encodes a target antibody from the immunoglobulin gene library directly obtained from untreated human lymphocytes and then establishing physical association between the DNA and the antibody chain with the use of phage particles, thereby enriching the phages presenting the antibody having affinity for the target via affinity screening. With the use of this method, an antibody having binding affinity for a target can be synthesized in a large amount by general techniques (e.g., JP2003-527832 A).

[0095] The composition described herein can be used for treatment of diseases or disorders.

Therapeutic composition

[0096] The present invention is associated with such diseases or disorders caused by abnormalities in phosphatidylinositol metabolism. Although not in accordance with the present invention as claimed, there is also described herein diseases or disorders caused by abnormalities in protein synthesis and intracellular pH.

[0097] In the case of protein synthesis, the protein synthesis associated with the binding of IRBIT to CPSF is controlled. In this case, IRBIT enhances protein synthesis and a substance that suppresses the functions of IRBIT (the above nucleic acid or antibody) can suppress protein synthesis. An example of an abnormality in protein synthesis, in which CPSF is involved, is tumor.

[0098] In the case of phosphatidylinositol metabolism, one example of diseases associated with abnormal activation of PIPKII is type 2 diabetes. IRBIT has an effect of suppressing PIPKII activity, so that IRBIT can be used for treatment of type 2 diabetes.

[0099] In the case of intracellular pH, pNBC 1 plays an important role in maintenance of intracellular pH. In particular, examples of diseases caused by such pH approaching or changing to acidic levels include eye diseases such as glaucoma and cataract and diseases such as dwarfism, mental retardation, and pancreatitis. IRBIT binds to pNBC1, so as to make it possible to maintain intracellular pH at a normal level.

[0100] The content of an active ingredient of the composition described herein ranges from, but is not limited to, approximately 1 μ g to 100 mg. The content can be varied depending on the types of active ingredient.

[0101] The dose of IRBIT in the composition described herein ranges from approximately 1 μ g to 1 mg and preferably approximately ranges from 50 μ g to 500 μ g per dosage unit, but is not limited to such ranges.

[0102] The dose of nucleic acid in the composition described herein ranges from, in terms of siRNA, antisense nucleic acid, or ribozyme, approximately 1 nM to 100 μ M and preferably approximately ranges from 10 nM to 50 μ M per dosage unit, but is not limited thereto.

[0103] The dose of an antibody or a fragment thereof in the composition described herein ranges from, but is not limited to, approximately 1 to 100 mg/ml and preferably approximately 5 to 70 mg/ml per dosage unit.

[0104] However, the above dose or dosage can be varied depending on conditions, age, sex, severity, or the like of a patient and should be determined based on the judgment made by a medical specialist.

[0105] Although not in accordance with the present invention as claimed, the composition described herein can generally contain a pharmaceutically acceptable carrier (specifically, an excipient or a diluent), such as sterilized water, physiological saline, buffer, or nonaqueous liquid (e.g., oil of almonds, plant oil, or ethanol). The composition can further contain a pharmaceutically acceptable stabilizer (e.g., an amino acid such as methionine), a preservative (methyl p-hydroxybenzoate or sorbic acid), an isotonic agent (e.g., sodium chloride), an emulsifying agent (e.g., lecithin or gum arabic), a suspending agent (e.g., a cellulose derivative), or the like.

[0106] Although not in accordance with the present invention as claimed, examples of preferable pharmaceutical preparations could include solutions, suspending agents, and emulsifying agents.

[0107] Although not in accordance with the present invention as claimed, examples of the route of administration of the composition described herein could include oral administration and parental administration (e.g. intravenous administration and local administration). Although not in accordance with the present invention as claimed, examples of local administration include surgical operation or a method that involves direct endoscopic injection into an affected part. Although not in accordance with the present invention as claimed, furthermore, the composition described herein could be administered in a single or in divided doses to a patient based on the treatment plan determined by a medical specialist at constant time intervals of 1 week, 2 weeks, 3 weeks, 1 month, 2 months, 6 months, 1 year, or the like.

[0108] Although not in accordance with the present invention as claimed, furthermore, an antibody or a fragment thereof could be delivered to a patient as described below. Such an antibody or a fragment thereof alone or the same encapsulated within a liposome (preferably, positively charged liposome), a microcapsule, nanoparticle, or the like is delivered in combination with generally an appropriate carrier (an excipient or a diluent) via oral or parenteral route (e.g., intravenous administration or local administration).

[0109] Although not in accordance with the present invention as claimed, the composition described herein could be used *in vitro*, *in vivo*, or *ex vivo*.

[0110] Although not in accordance with the present invention as claimed, *ex vivo* cells or tissues that have been once removed from the patient's body can be returned to the body after treatment with the active ingredients of the present invention. Accordingly, cells or tissues in which abnormalities occur in protein synthesis, phosphatidylinositol metabolism, or intracellular pH can be caused to return to a **normal state**.

2. Use examples of IRBIT

[0111] The present invention provides a use of IRBIT in control of intracellular phosphatidylinositol metabolism *in vitro* or *ex vivo*, although the latter is not in accordance with the present invention as claimed.

[0112] This method is based on the fact that IRBIT has an effect of suppressing PIPKII **activity**.

[0113] Although not in accordance with the present invention as claimed, there is also described herein a use of IRBIT in control of intracellular protein synthesis *in vitro* or *ex vivo*.

[0114] This method is based on the fact that IRBIT has an effect of binding to CPSF, **thereby controlling CPSF functions**.

[0115] Although not in accordance with the present invention as claimed, there is also described herein a use of IRBIT in control of intracellular pH *in vitro* or *ex vivo*.

[0116] This method is based on the fact that IRBIT has an effect of activating pNBC1. Moreover, the activation of pNBC1 requires phosphorylation of IRBIT

[0117] Although not in accordance with the present invention as claimed, as described above, IRBIT can be used *in vitro* for screening for a substance that makes it possible to control intracellular protein synthesis, phosphatidylinositol metabolism, or intracellular pH. Although not in accordance with the present invention as claimed, furthermore, IRBIT can be used *ex vivo* for causing abnormal conditions of cells or tissues (including abnormal protein synthesis, abnormal phosphatidylinositol metabolism, or abnormal intracellular pH) to return to normal conditions.

3. Screening

[0118] Although not in accordance with the present invention as claimed, there is also described herein a method for screening for a substance, comprising measuring the binding of IRBIT to CPSF, PIPKII, or pNBC1 in the presence of candidate substances and then identifying a substance that suppresses or enhances the binding.

[0119] Regarding the above binding, the binding of IRBIT to CPSF, PIPKII, or pNBC1 can be measured *in vitro* or within cells (in particular, mammalian cells) in the presence of candidate substances. Examples of mammalian cells include CHO, COS, HEK293, HeLa, and NIH3T3.

[0120] The thus identified substance can be used for treatment or diagnosis, for example. In particular, such substance controls at least one intracellular biological function selected from the group consisting of intracellular protein synthesis, phosphatidylinositol metabolism, and intracellular pH.

[0121] When the above binding is performed *in vitro*, for example, IRBIT and CPSF, PIPKII, or pNBC1 are caused to present in an appropriate buffer, a candidate substance is added to the buffer, and then the level of binding of IRBIT to CPSF, PIPKII, or pNBC1 can be detected by SDS-PAGE and the immunoblot method. This system is effective for detection of a substance that suppresses or inhibits the above binding. The recombinant protein of IRBIT, CPSF, PIPKII, or pNBC1 can be prepared by techniques similar to those described in the section of the above IRBIT.

[0122] When the binding is performed within cells, DNAs encoding the following proteins are incorporated into the same vector or different vectors so that IRBIT and CPSF, PIPKII, or pNBC1 can be expressed simultaneously or separately and then mammalian cells are transformed or transfected with the vectors. The translated proteins are caused to present within cells and particularly in the cytoplasm. Preferably, vector DNA contains no secretion signal sequence.

[0123] The amino acid and nucleotide sequences of CPSF, PIPKII, or pNBC1 are available from the GenBank, from documents, or the like. The amino acid and nucleotide sequences are deposited under accession Nos: AB092504; AF030558 and AF033355; or NM_003759 and NM_018760; respectively. The amino acid and nucleotide sequences of IRBIT are as described above.

[0124] Expression vectors are any vectors that can be used in preferably mammalian cells. A vector can contain regulatory sequences such as a promoter, an enhancer, a replication origin, a ribosome-binding site, a multicloning site, a terminator, and poly A signal. As an expression vector, a commercially available vector such as pSG5, pXT1 (Stratagene), pSVK3, pBPV, pMSG and pSVL SV40 (Pharmacia), pHM6, pVM6, and pXM (Roche Diagnostics) can be adequately selected and used.

[0125] Examples of a promoter include a CMV promoter, an SV40 promoter, and an EF promoter.

[0126] Examples of a method for introducing DNA encoding IRBIT, CPSF, PIPKII, or pNBC1 into host cells include a calcium phosphate method, a lipofection method, an electroporation method, and methods using viral infection with

adenovirus, retrovirus, or the like (Experimental Medicine, Separate Volume, 4th edition, edited by Masami Matsumura et al., "New genetic engineering handbook" (2003) YODOSHA, Tokyo, Japan) .

[0127] Alternatively, non-human transgenic animals (e.g., mice) are produced by incorporating an IRBIT gene exogeneously into the genome by a known technique using oocytes or embryonic stem cells of a non-human animal so that the gene can be forcedly expressed. Furthermore, non-human transgenic animals (e.g., mice) are produced by incorporating a CPSF, PIPKII, or pNBC1 gene exogeneously into the genome, so that the gene can be forcedly expressed. Non-human chimeric animals and their progenies can thus be produced by mating both transgenic animals and then producing non-human chimeric animals and their progenies capable of expressing the IRBIT gene and the CPSF, PIPKII, or pNBC1 gene.

[0128] The ways in which the binding of IRBIT (forcedly expressed within cells or within a non-human transgenic animal) to CPSF, PIPKII, or pNBC1 is affected in the presence of candidate substances incorporated within cells or animals are examined by measuring the binding according to a pull down method, an immunoprecipitation method, or the like. At the same time, the effects on intracellular synthesis of a specific protein, the effects of the aforementioned binding on phosphatidylinositol metabolism, and the effects on intracellular pH are examined.

[0129] The effects on intracellular protein synthesis can be measured by the Western blot method or the like using an antibody against a specific protein, for example.

[0130] The effects on phosphatidylinositol metabolism can be measured via quantification of PIP2 level using a [³H] label or a PIP2-binding protein, for example.

[0131] The effects on intracellular pH can be measured by intracellular pH measurement or the like using a fluorescent pH indicator or the like.

[0132] Examples of candidate substances include, but are not limited to, organic small molecules, peptides, polypeptides, proteins, nucleosides, oligonucleotides, polynucleotides, and nucleic acids (DNA or RNA).

[0133] As described above, IRBIT has extremely important significance for all cells in regulation of intracellular metabolism, pH change, ion balance, and phospholipid metabolism, control of protein synthesis, control of Ca²⁺ release, and the like. It has been revealed herein that through control of IRBIT concentration or expression pattern, various biological functions described above can be controlled. Substances that are identified by the screening method of the present invention are useful for such control.

EXAMPLES

[0134] Hereinafter, the present invention will be described in detail by examples as follows, but the scope of the present invention is not limited by these examples.

<Example 1>

Involvement of IRBIT in cytoplasmic poly (A) addition reaction

[0135] CPSF is a group of conjugated proteins consisting of 4 subunits: CPSF160, CPSF100, CPSF73, and CPSF30. First, to reveal to which one of the 4 subunits IRBIT binds, myc-tagged cDNAs prepared by tagging each subunit with myc were expressed together with IRBIT in various combinations by COS cells or the like and then the way in which coprecipitation occurs upon immunoprecipitation of IRBIT or CPSF was examined.

[0136] Mouse-derived CPSF was used herein. CPSF was cloned by RT-PCR based on the full-length sequence under GenBank accession No. AB092504 and the nucleotide sequence of each of four subunits (H. Ando et al., 2003, *ibid*). Mouse-derived IRBIT was used herein. IRBIT was cloned by RT-PCR based on the nucleotide sequence under GenBank accession No. NM_145542 (H. Ando et al., 2003, *ibid*).

[0137] DNA was prepared by fusing DNA encoding an myc tag to cDNA encoding the full-length of or each subunit of CPSF. The thus prepared DNA was inserted into a vector for mammals. Meanwhile, cDNA encoding IRBIT was also inserted into the same vector. COS cells were transformed with the thus prepared vector and then the above DNA was co-expressed.

[0138] Cells were separated by centrifugation and then lysed. Cytoplasmic fractions were collected and then the binding of IRBIT to CPSF was detected using an anti-IRBIT antibody (JP2004-129612 A) and an anti-myc antibody.

[0139] As a result, IRBIT was revealed to bind to CPSF via the CPSF160 subunit (Fig. 1).

[0140] Furthermore, a similar experiment conducted using CPSF160 deletion mutants revealed that IRBIT binds to the mRNA binding site of CPSF 160 (Fig. 2).

[0141] The mRNA binding site of CPSF 160 is a region essential for CPSF to recognize mRNA to which poly (A) is added. Hence, the fact that IRBIT binds to this region suggests that IRBIT may inhibit CPSF functions. To confirm this, the binding of IRBIT with CPSF and the effects of the binding on polyadenylation reaction were examined with the use of an *in vitro* rearrangement system using purified proteins.

[0142] CPSF is a molecule essential for mRNA polyadenylation reaction, but the location at which CPSF functions is mainly within the nucleus. CPSF functions for RNA just after its transcription from DNA; that is, CPSF functions as a member involved in the process of the maturation reaction of such RNA until the RNA is released from the nucleus in the form of mRNA. However, exceptionally, in other situations such as in the process of oocyte maturation or local synthesis of new proteins in neurons, CPSF is involved in cytoplasmic mRNA polyadenylation and regulates the length of poly (A), so as to regulate the protein synthesis of a target molecule (Daron C. Barnard et al., Cell 2004, 119: 641-651). This cytoplasmic polyadenylation reaction is an exceptional phenomenon, but is an essential phenomenon for oocyte maturation and the establishment of neuroplasticity. When the intracellular distribution of IRBIT is examined, almost no IRBIT is present within the nucleus. Accordingly, the present inventors conducted examinations with a focus on the involvement of IRBIT in cytoplasmic polyadenylation reaction.

[0143] A *Xenopus* oocyte system is better for examination of cytoplasmic polyadenylation reaction. Hence, the present inventors first performed cloning and isolation of IRBIT cDNA from *Xenopus* oocytes according to a standard method. Unlike mammalian cells, 3 types of IRBIT mRNAs were expressed in *Xenopus* oocytes, and the 3 types of IRBIT mRNAs were revealed to bind to CPSF160.

[0144] The 3' terminus of IP₃ receptor mRNA has a sequence that may be subjected to a cytoplasmic polyadenylation reaction. Actually, in cerebellum Purkinje cells, the mRNA is present within dendrites. This suggests the possibility that IRBIT released from the IP₃ receptor controls IP₃ receptor protein synthesis via binding with CPSF.

[0145] Moreover, the present inventors discovered that IRBIT has an effect of further enhancing polyadenylation activity in the presence of PAP and Fip1 (CPSF subunits) (I. Kaufmann et al., EMBO J. (2004) 23: 616-626).

[0146] It was demonstrated based on the above results that IRBIT controls protein synthesis via the binding to CPSF within cells.

<Example 2>

Interaction between IRBIT and PIP kinase type II (PIPKII)

[0147] The present inventors further searched for molecules that interact with IRBIT.

[0148] With procedures similar to those used in Example 1, FLAG-IRBIT was overexpressed in HEK293 cells, immunoprecipitation was performed using an anti-FLAG antibody, and then proteins that had been coprecipitated with IRBIT were analyzed using a mass spectroscopic analyzer. As a result, phosphatidylinositol-5-phosphate 4-kinase γ (PIPKII γ), which is an enzyme for phosphorylating phospholipids, was identified. The PIPKII family (PIPKII α , PIPKII β , and PIPKII γ) is the family of enzymes involved in the synthesis of PIP₂. In view of the fact that IP₃ is produced by hydrolysis of PIP₂, a signaling mechanism in which the IP₃ receptor, IRBIT, and PIPKII γ are involved may be present (Fig. 3).

[0149] Furthermore, the following experiment was conducted and the results are as described below.

(1) Mouse IRBIT and Myc-PIPKII α , β , or γ were over-expressed in COS-7, followed by immunoprecipitation. Specifically, cells were solubilized with a lysis buffer (10 mM HEPES, 100 mM NaCl, 2 mM EDTA, 1% P-40, pH 7.4) and then centrifuged (20000 \times g, 30 minutes) to collect supernatants. Subsequently, an anti-Myc antibody or an anti-IRBIT antibody was added and then reaction was performed for 1 hour. Furthermore, protein G Sepharose[®] was added before one hour reaction, and the immune complex was washed with a lysis buffer and then eluted with a SDS-PAGE sample buffer. Subsequently, Western blotting was performed using an anti-Myc antibody or an anti-IRBIT antibody. As a result, IRBIT was coprecipitated with Myc-PIPKII α , β , and γ (Fig. 4).

(2) Immunoprecipitation was performed using an anti-IRBIT antibody and the mouse cerebellum, so that PIPKII α was coprecipitated (Fig. 5). The *in vivo* binding of IRBIT with PIPKII was confirmed based on the results.

(3) Mouse IRBIT and Myc-PIPKII γ were overexpressed in COS-7 and then immunostaining was performed. Specifically, after fixation with 4% paraformaldehyde, transmembrane treatment with 0.1% Triton X-100 and blocking treatment with 2% goat serum were performed, a mouse anti-Myc antibody and a rabbit anti-IRBIT antibody were added, and then reaction was performed at room temperature for 1 hour. Alexa488-conjugated anti-mouse IgG antibody and Alexa594-conjugated anti-rabbit IgG antibody were added as secondary antibodies, followed by 45 minutes of reaction at 37°C. As a result, both IRBIT and Myc-PIPKII γ were revealed to be localized within the cytoplasm (Fig. 6).

(4) Mouse IRBIT deletion mutants (60-530, 78-530, 105-530, 1-277, 1-104, 1-90, and 1-77 in the amino acid sequence of SEQ ID NO: 3) were prepared by amplifying the corresponding sequences by PCR and then cloning them into GFP fusion protein expression vectors pEGFP-C1 (Clontech). The binding of each of these mutants with PIPKII α was examined, revealing that a serine-rich region existing in the N-terminal region of IRBIT is important for the above binding (Fig. 7).

(5) Point mutants (T52A, T58A, S62A, S64A, S66A, S68A, S70A, S71A, T72A, S74A, S76G, S77A, S80A, D83A, S90A, and T97A in the amino acid sequence of SEQ ID NO: 3) of the mouse IRBIT serine-rich region were prepared

using a site-directed mutagenesis kit (Stratagene). The binding of each of these mutants with PIPKII α was examined. As a result, IRBIT Ser68 and Ser71 were found to be important for the above binding (Fig. 8).

[0150] It was confirmed based on the above results that IRBIT binds to the PIPKII family. Furthermore, IRBIT suppresses PIPKII activity, as revealed from preliminary data. Thus, it was concluded that IRBIT controls the activity of phosphatidylinositol metabolism.

<Example 3>

10 Activation of NBC1 (Na/HCO₃ cotransporter 1) by IRBIT

[0151] The present inventors further identified an NBC1 (sodium bicarbonate co-transporter 1) protein as a protein that binds to IRBIT, which transports sodium ions and bicarbonate on the cell membrane.

[0152] It is known that NBC1 includes two splicing mutants, p type and k type (Seth L. Alper, Annu. Rev. Physiol. 2002, 64: 899-923). It was revealed that IRBIT binds to p-type NBC1 (referred to as pNBC1) of these mutants and that the binding of IRBIT to pNBC1 requires phosphorylation of some serine residues in IRBIT. Experiments and results conducted are as specifically described below.

20 (1) Identification of IRBIT-binding region of NBC1

[0153] The p-type-specific N-terminal sequence (amino acids 1-85) of pNBC1 was revealed to be essential for binding with IRBIT. Other deletion mutants were prepared within the 85 amino acids, and then the binding of these mutants with IRBIT was studied using pull down assay. Specifically, the following experiment was conducted.

[0154] HA-IRBIT was overexpressed in COS-7 cells and then cell extracts were prepared. A recombinant protein of a MBP-pNBC1 deletion mutant was added. After reaction, the bound proteins were pulled down with an amylose resin. HA-IRBIT was detected by Western blotting using an anti-HA antibody.

[0155] As a result, it was demonstrated that binding with IRBIT is possible if the 62 N-terminal amino acids of the p-type-specific sequence are present (Fig. 9 and Fig. 10). However, if an N-terminal portion or a C-terminal portion was deleted from the 62 amino acids, the binding with IRBIT became impossible to observe. Thus, it was concluded that an IRBIT-binding region with a length shorter than the 62 amino acids makes it difficult for the binding to take place. It was suggested based on the results that a sequence with a length of several specific amino acids of pNBC1 is not involved in the binding with IRBIT, but rather that a three-dimensional structure comprising dozens of amino acids may be required for binding with IRBIT.

35 (2) Identification of pNBC1-binding region of IRBIT

[0156] Next, various mouse IRBIT (full-length 530 amino acids; SEQ ID NO: 3) deletion mutants were expressed in COS7 cells and then the ability of each mutant to bind to pNBC1 was examined by pull down assay. Specifically, the following experiment was conducted.

[0157] GFP-IRBIT deletion mutants were overexpressed in COS-7 cells and then cell extracts were prepared. A MBP-pNBC1 (1-85) recombinant protein was added. After reaction, the bound proteins were pulled down with an amylose resin. GFP-IRBIT deletion mutants were detected by Western blotting using an anti-GFP antibody.

[0158] The results are shown in Fig. 11. As shown in Fig. 11, it was demonstrated that even deletion mutants (1-104 and 1-277) expressing the IRBIT N-terminal portions (confirmed to bind to the IP₃ receptor) were unable to bind to pNBC1. Moreover, the deletion mutant (105-530) expressing an IRBIT C-terminal portion (previously confirmed to be unable to bind to the IP₃ receptor) was also unable to bind to pNBC1. A serine phosphorylation site demonstrated to be necessary for binding with pNBC1 is completely contained within 1-104. Thus, it was considered that although a short pNBC1-binding sequence is present in 1-104, the binding with pNBC1 may be inhibited by some kind of three-dimensional structure. Hence, deletion mutants were prepared via further fine deletion of 1-104, and then the binding of each of these mutants with pNBC1 was analyzed. However, none of these mutants were observed to bind to pNBC1.

[0159] Based on the above results, it was revealed that the entire structure, including the IRBIT N-terminus and the C-terminus, is required for binding with pNBC1. It was also revealed that the binding manner of IRBIT with pNBC1 is different from that of IRBIT with IP₃ receptor, and such binding can be achieved only in the presence of the IRBIT N-terminus.

55 (3) Binding of endogenous IRBIT with NBC1

[0160] Next, an experiment was conducted to confirm the binding of endogenous IRBIT to NBC 1 using immunopre-

cipitation.

[0161] NBC1 had been previously identified as an IRBIT-binding protein in a cerebellum membrane fraction. First, immunoprecipitation was performed using a cerebellum membrane fraction extract. Furthermore, since IRBIT is known to also bind to an IP₃ receptor in a cerebellum membrane fraction, whether or not a triple complex of NBC1, IRBIT, and an IP₃ receptor could be formed was examined. Specifically, the following experiment was conducted.

[0162] After solubilization of the cerebellum membrane fraction with a surfactant, an anti-IRBIT antibody, an anti-NBC antibody, or a control antibody was added to perform a reaction. Moreover, protein G Sepharose® was added to extract immune complexes. Subsequently, Western blotting was performed using the anti-IRBIT antibody, the anti-NBC antibody, or the anti-IP₃R antibody.

[0163] As a result, it was revealed that the anti-NBC1 antibody immunoprecipitate contained endogenous IRBIT and the anti-IRBIT antibody immunoprecipitate contained endogenous NBC1. Thus it was confirmed that endogenous IRBIT and NBC1 form a complex in a cerebellum membrane fraction (Fig. 12). Meanwhile, as already reported by Ando et al., (2003, *ibid*), the IP₃ receptor was detected in the anti-IRBIT immunoprecipitate, but no IP₃ receptor could be detected in the anti-NBC1 precipitate. These results demonstrate that most of the endogenous NBC1 does not form any triple complex containing IRBIT and the IP₃ receptor.

[0164] Furthermore, to confirm that the binding of IRBIT to NBC1 is universal, the binding of endogenous IRBIT to NBC1 was examined by the immunoprecipitation similar to the above method using COS7 cell extracts. Specifically, the following experiment was conducted.

[0165] An anti-IRBIT antibody, an anti-NBC antibody, or a control antibody was added to the COS-7 cell extract to perform a reaction. Moreover, protein G Sepharose® was added to extract immune complexes. Subsequently, Western blotting was performed using an anti-IRBIT antibody or an anti-NBC antibody.

[0166] Fig. 13 shows the results. As shown in Fig. 13, when COS7 cells were extracted with the same buffer as that used for preparation of a cerebellum membrane fraction extract, no binding of endogenous IRBIT to NBC1 was detected. However, when a similar experiment was conducted with addition of 2 mM CaCl₂ to the buffer, the binding of endogenous IRBIT to NBC1 was detected. This buffer had contained 2 mM EDTA in advance, so it was considered that the calcium ion concentration was several μM when 2 mM CaCl₂ was added. These results suggest that the binding of endogenous NBC 1 to IRBIT might be controlled differently (i.e., in different manners) in a cerebellum membrane fraction and COS7 cells.

(4) Activation of NBC1 by IRBIT

[0167] Mouse IRBIT cRNA and the cRNA of human pNBC1 or kNBC1 were simultaneously injected into cultured cells of *Xenopus* oocytes. The membrane potential was set at -25 mV, and then current changes were measured when an ND96 solution (96 mM NaCl, 2 mM KCl, 1 mM MgCl₂, 5 mM HEPES, and pH 7.4) was replaced by an ND96+HCO₃⁻ solution. As controls, IRBIT, pNBC1, and kNBC1 were used.

[0168] As a result, only the pNBC1 activity was enhanced approximately 6 to 7 times by simultaneous injection of IRBIT (Fig. 14A). Similarly, when the membrane potential was varied within a range from -160 mV to +60 mV in the ND96 + HCO₃⁻ solution, only the pNBC1 activity was significantly enhanced by simultaneous injection of IRBIT (Fig. 14B).

[0169] Furthermore, an IRBIT mutant (S68A, S71A, S74A, or S77A) was prepared in which an IRBIT phosphorylation site had been substituted with alanine (A). The cRNA of the IRBIT mutant and the cRNA of human pNBC1 were simultaneously injected into *Xenopus* oocytes in a manner similar to the above. The membrane potential was set at -25 mV, and then current changes were measured when the ND96 solution was replaced by the ND96+HCO₃⁻ solution.

[0170] As a result, the effects of IRBIT to enhance the pNBC1 activity were observed to disappear in all IRBIT mutants in which the phosphorylation sites had been substituted with alanine (Fig. 15). This demonstrates that phosphorylation of IRBIT is required for activation of pNBC1.

INDUSTRIAL APPLICABILITY

[0171] IRBIT is involved in protein synthesis, phosphatidylinositol metabolism, or intracellular pH maintenance within mammalian cells via its binding to CPSF, PIPKII, or pNBC1. Hence, IRBIT controlling such biological functions, a nucleic acid controlling IRBIT expression and translation, or an antibody against IRBIT may be useful for treating diseases arising from abnormalities in such functions. Hence, the composition and methods according to the present invention are useful for treating diseases arising from abnormalities in such functions.

[0172] According to the present invention, it is revealed that IRBIT is extremely significant in all cells for regulating intracellular phospholipid metabolism. Although not in accordance with the present invention as claimed, through control of the concentration or expression pattern of IRBIT, various biological functions can be controlled. IRBIT is strongly expressed in the cerebral nervous system such as in the choroid plexus, neurons, and glial cells, and it also contributes to the control of the functions of the cerebral nervous system. Furthermore, IRBIT may be useful for treating diseases

EP 2 311 477 B1

that are developed when the functions of pNBCI, which are important in intracellular pH maintenance, are activated, and in particular, when pH approaches or changes to acidic levels. Examples of such diseases include eye diseases such as glaucoma and cataract and diseases such as dwarfism, mental retardation, and pancreatitis. Furthermore, IRBIT has an effect of suppressing or inhibiting PIPKII activity, so that IRBIT could perhaps be used for treating type 2 diabetes. As described above, three proteins that are targets of IRBIT were specified by the present invention. IRBIT, an inhibitor of IRBIT, and an agent for enhancing IRBIT are useful for controlling phosphatidylinositol metabolism in a cell in vitro in which the PIP₂ synthetase α , β , or γ PIPKII is involved.

SEQUENCE LISTING

[0173]

<110> Japan science and Technology Agency

<120> Control of intracellular target molecule by IP3 receptor-binding protein

<130> P035168EPA

<150> JP 2006-77607

<151> 2006-03-20

<160> 6

<170> PatentIn Ver. 2.1

<210> 1

<211> 530

<212> PRT

<213> Homo sapiens

<400> 1

EP 2 311 477 B1

Met Ser Met Pro Asp Ala Met Pro Leu Pro Gly Val Gly Glu Glu Leu
 1 5 10 15
 Lys Gln Ala Lys Glu Ile Glu Asp Ala Glu Lys Tyr Ser Phe Met Ala
 20 25 30
 Thr Val Thr Lys Ala Pro Lys Lys Gln Ile Gln Phe Ala Asp Asp Met
 35 40 45
 Gln Glu Phe Thr Lys Phe Pro Thr Lys Thr Gly Arg Arg Ser Leu Ser
 50 55 60
 Arg Ser Ile Ser Gln Ser Ser Thr Asp Ser Tyr Ser Ser Ala Ala Ser
 65 70 75 80
 Tyr Thr Asp Ser Ser Asp Asp Glu Val Ser Pro Arg Glu Lys Gln Gln
 85 90 95
 Thr Asn Ser Lys Gly Ser Ser Asn Phe Cys Val Lys Asn Ile Lys Gln
 100 105 110
 Ala Glu Phe Gly Arg Arg Glu Ile Glu Ile Ala Glu Gln Asp Met Ser
 115 120 125
 Ala Leu Ile Ser Leu Arg Lys Arg Ala Gln Gly Glu Lys Pro Leu Ala
 130 135 140
 Gly Ala Lys Ile Val Gly Cys Thr His Ile Thr Ala Gln Thr Ala Val
 145 150 155 160
 Leu Ile Glu Thr Leu Cys Ala Leu Gly Ala Gln Cys Arg Trp Ser Ala
 165 170 175
 Cys Asn Ile Tyr Ser Thr Gln Asn Glu Val Ala Ala Ala Leu Ala Glu
 180 185 190
 Ala Gly Val Ala Val Phe Ala Trp Lys Gly Glu Ser Glu Asp Asp Phe
 195 200 205
 Trp Trp Cys Ile Asp Arg Cys Val Asn Met Asp Gly Trp Gln Ala Asn
 210 215 220
 Met Ile Leu Asp Asp Gly Gly Asp Leu Thr His Trp Val Tyr Lys Lys
 225 230 235 240
 Tyr Pro Asn Val Phe Lys Lys Ile Arg Gly Ile Val Glu Glu Ser Val
 245 250 255

45

50

55

EP 2 311 477 B1

Thr Gly Val His Arg Leu Tyr Gln Leu Ser Lys Ala Gly Lys Leu Cys
 260 265 270
 Val Pro Ala Met Asn Val Asn Asp Ser Val Thr Lys Gln Lys Phe Asp
 5 275 280
 Asn Leu Tyr Cys Cys Arg Glu Ser Ile Leu Asp Gly Leu Lys Arg Thr
 290 295 300
 Thr Asp Val Met Phe Gly Gly Lys Gln Val Val Val Cys Gly Tyr Gly
 10 305 310 315 320
 Glu Val Gly Lys Gly Cys Cys Ala Ala Leu Lys Ala Leu Gly Ala Ile
 325 330 335
 Val Tyr Ile Thr Glu Ile Asp Pro Ile Cys Ala Leu Gln Ala Cys Met
 15 340 345 350
 Asp Gly Phe Arg Val Val Lys Leu Asn Glu Val Ile Arg Gln Val Asp
 355 360 365
 Val Val Ile Thr Cys Thr Gly Asn Lys Asn Val Val Thr Arg Glu His
 20 370 375 380
 Leu Asp Arg Met Lys Asn Ser Cys Ile Val Cys Asn Met Gly His Ser
 385 390 395 400
 Asn Thr Glu Ile Asp Val Thr Ser Leu Arg Thr Pro Glu Leu Thr Trp
 25 405 410 415
 Glu Arg Val Arg Ser Gln Val Asp His Val Ile Trp Pro Asp Gly Lys
 420 425 430
 Arg Val Val Leu Leu Ala Glu Gly Arg Leu Leu Asn Leu Ser Cys Ser
 30 435 440 445
 Thr Val Pro Thr Phe Val Leu Ser Ile Thr Ala Thr Thr Gln Ala Leu
 450 455 460
 Ala Leu Ile Glu Leu Tyr Asn Ala Pro Glu Gly Arg Tyr Lys Gln Asp
 35 465 470 475 480
 Val Tyr Leu Leu Pro Lys Lys Met Asp Glu Tyr Val Ala Ser Leu His
 485 490 495
 Leu Pro Ser Phe Asp Ala His Leu Thr Glu Leu Thr Asp Asp Gln Ala
 40 500 505 510
 Lys Tyr Leu Gly Leu Asn Lys Asn Gly Pro Phe Lys Pro Asn Tyr Tyr
 515 520 525
 Arg Tyr
 45 530

<210> 2

<211> 2677

<212> DNA

50 <213> Homo sapiens

<400> 2

55

5
 cagagtgcc tttctcccc cctcttcccc ctcccgggag ctgccagtac ttgacgtggc 60
 gtcaccgccc tctaccctcg ctttgcgtgc gtgtttgcgt acagcggagg tggcggcgcg 120
 ggcaggtcgg agctcggagc tgctgcttct ggttctcttg tggccaccgt cgctgtccgg 180
 ctgccttggg ctgccgaaca gacaaggcgt gggccacagc acctcagaag ccgacgcagc 240
 tcgacgcagg ggccggcagg aggggtgggcg atcgcgtgtc ggagggcgcc gcgcgggcag 300
 gcgggagggg gccagagggg gaaagagggc ggggcggcgg gtcagccgct ggccgggccc 360

10
 gcgggggaat gtcgatgcct gacgcgatgc cgctgcccgg ggtcggggag gagctgaagc 420
 aggccaagga gatcgaggac gccgagaagt actccttcat ggccaccgtc accaaggcgc 480
 ccaagaagca aatccagttt gctgatgaca tgcaggagtt caccaaattc cccacaaaaa 540
 ctggccgaag atctttgtct cgctcgatct cacagtctct cactgacagc tacagttcag 600
 ctgcatccta cacagatagc tctgatgatg aggtttctcc ccgagagaag cagcaaacca 660
 actccaaggg cagcagcaat tctgtgtgta agaacatcaa tttggacgcc tttggacgcc 720
 gggagattga gattgcagag caagacatgt ctgctctgat ttactcagg aaacgtgctc 780
 agggggagaa gcccttggct ggtgctaaaa tagtgggctg tacacacatc acagcccaga 840
 cagcgggtgt gattgagaca ctctgtgcc tgggggctca gtgccgctgg tctgcttgta 900
 acatctactc aactcagaat gaagtagctg cagcactggc tgaggctgga gttgcagtgt 960
 tcgcttgaa gggcgagtca gaagatgact tctggtggtg tattgaccgc tgtgtgaaca 1020
 tggatgggtg gcaggccaac atgatcctgg atgatggggg agacttaacc cactgggttt 1080
 ataagaagta tccaacgtg ttaagaaga tccgaggcat tgtggaagag agcgtgactg 1140
 gtgttcacag gctgtatcag ctctccaaag ctgggaagct ctgtgttccg gccatgaacg 1200
 tcaatgattc tgttaccaaa cagaagtttg ataacttgta ctgctgccga gaatccattt 1260
 tggatggcct gaagaggacc acagatgtga tgtttggtgg gaaacaagtg gtggtgtgtg 1320
 gctatgggtg ggtaggcaag ggctgctgtg ctgctctcaa agctcttggg gcaattgtct 1380
 acattaccga aatcgacccc atctgtgctc tgcaggcctg catggatggg ttcagggtgg 1440
 taaagctaaa tgaagtcac cggcaagtcg atgtcgtaat aacttgcaca ggaaataaga 1500
 atgtagtac acgggagcac ttggatcgca tgaaaaacag ttgtatcgta tgcaatatgg 1560
 gccactccaa cacagaaatc gatgtgacca gcctccgcac tccggagctg acgtgggagc 1620
 gagtacgttc tcaggtggac catgtcatct ggccagatgg caaacgagtt gtctctctgg 1680
 cagccacaac tctactcaat ttgagctgct ccacagttec cacctttgtt tgtccatca 1740
 agcaggatgt gtacttgctt gcaactgtag aactctataa tgcacccgag gggcgataca 1800
 catcatttga tgcccacctt cctaagaaaa tggatgaata cgttgccagc ttgcatctgc 1860
 acaaaaatgg gccattcaaa cctaattatt acagagctga cagatgacca agcaaaatat ctgggactca 1920
 ccagtccacc tgaaccacac actctaaaga aatatttttt aagataactt ttattttctt 2040
 ctactcctt tctcttggat ttttttctta taatttcatt ctgttttttt catctcatta 2100
 tccaagttct gcagaccaca caggaacttg ctatcatggct ctttagatga aatagaagtt 2160
 caggggtccc cactctagtc actaaagaag gattttactc ccccagcca gaaaggtgat 2220
 tcttctcttt accatttctg gggactttag tcttaattag gtaccttatt aacaggaaat 2280
 gctaaggtag cttctctgtg gaacaatctg caatgtctaa atcgccttaa aagagcccat 2340
 ttcttagctg ctgaaatcag tgcctttca cttcttcaga gaagcaggga tggtagctac 2400
 ccggcaggta ggttagatgt ggggtgtgca tgtaatttc ccttagaagt tccaagccct 2460
 gtttcctgcg taaaggtggt atgtccagtt cagagatgtg tataatgagc atggcttggt 2520
 aagatcagga ggcccacttg gatttatagt atagcccttc ctccactccc accagacttg 2580
 ctcatttttc gagtttttaa ctgactaca ctctatttga gtttaatttt gtctcttagg 2640
 atttatttct gttgtccaaa aaaaaaaaaa aaaaaaa 2677

<210> 3

<211> 530

45 <212> PRT

<213> Mus musculus

<400> 3

50

55

EP 2 311 477 B1

Met Ser Met Pro Asp Ala Met Pro Leu Pro Gly Val Gly Glu Glu Leu
 1 5 10 15
 Lys Gln Ala Lys Glu Ile Glu Asp Ala Glu Lys Tyr Ser Phe Met Ala
 5 20 25 30
 Thr Val Thr Lys Ala Pro Lys Lys Gln Ile Gln Phe Ala Asp Asp Met
 35 40 45
 Gln Glu Phe Thr Lys Phe Pro Thr Lys Thr Gly Arg Arg Ser Leu Ser
 10 50 55 60
 Arg Ser Ile Ser Gln Ser Ser Thr Asp Ser Tyr Ser Ser Ala Ala Ser
 65 70 75 80
 Tyr Thr Asp Ser Ser Asp Asp Glu Val Ser Pro Arg Glu Lys Gln Gln
 15 85 90 95
 Thr Asn Ser Lys Gly Ser Ser Asn Phe Cys Val Lys Asn Ile Lys Gln
 100 105 110

20

25

30

35

40

45

50

55

EP 2 311 477 B1

Ala Glu Phe Gly Arg Arg Glu Ile Glu Ile Ala Glu Gln Asp Met Ser
115 120 125

Ala Leu Ile Ser Leu Arg Lys Arg Ala Gln Gly Glu Lys Pro Leu Ala
5 130 135 140

Gly Ala Lys Ile Val Gly Cys Thr His Ile Thr Ala Gln Thr Ala Val
145 150 155 160

Leu Ile Glu Thr Leu Cys Ala Leu Gly Ala Gln Cys Arg Trp Ser Ala
10 165 170 175

Cys Asn Ile Tyr Ser Thr Gln Asn Glu Val Ala Ala Ala Leu Ala Glu
180 185 190

Ala Gly Val Ala Val Phe Ala Trp Lys Gly Glu Ser Glu Asp Asp Phe
15 195 200 205

Trp Trp Cys Ile Asp Arg Cys Val Asn Met Asp Gly Trp Gln Ala Asn
210 215 220

Met Ile Leu Asp Asp Gly Gly Asp Leu Thr His Trp Val Tyr Lys Lys
20 225 230 235 240

Tyr Pro Asn Val Phe Lys Lys Ile Arg Gly Ile Val Glu Glu Ser Val
245 250 255

Thr Gly Val His Arg Leu Tyr Gln Leu Ser Lys Ala Gly Lys Leu Cys
25 260 265 270

Val Pro Ala Met Asn Val Asn Asp Ser Val Thr Lys Gln Lys Phe Asp
275 280 285

Asn Leu Tyr Cys Cys Arg Glu Ser Ile Leu Asp Gly Leu Lys Arg Thr
30 290 295 300

Thr Asp Val Met Phe Gly Gly Lys Gln Val Val Val Cys Gly Tyr Gly
305 310 315 320

Glu Val Gly Lys Gly Cys Cys Ala Ala Leu Lys Ala Leu Gly Ala Ile
35 325 330 335

Val Tyr Ile Thr Glu Ile Asp Pro Ile Cys Ala Leu Gln Ala Cys Met
340 345 350

Asp Gly Phe Arg Val Val Lys Leu Asn Glu Val Ile Arg Gln Val Asp
40 355 360 365

Val Val Ile Thr Cys Thr Gly Asn Lys Asn Val Val Thr Arg Glu His
370 375 380

Leu Asp Arg Met Lys Asn Ser Cys Ile Val Cys Asn Met Gly His Ser
45 385 390 395 400

Asn Thr Glu Ile Asp Val Thr Ser Leu Arg Thr Pro Glu Leu Thr Trp
405 410 415

Glu Arg Val Arg Ser Gln Val Asp His Val Ile Trp Pro Asp Gly Lys
50 420 425 430

Arg Val Val Leu Leu Ala Glu Gly Arg Leu Leu Asn Leu Ser Cys Ser
435 440 445

Thr Val Pro Thr Phe Val Leu Ser Ile Thr Ala Thr Thr Gln Ala Leu
450 455 460

Ala Leu Ile Glu Leu Tyr Asn Ala Pro Glu Gly Arg Tyr Lys Gln Asp
55 465 470 475 480

EP 2 311 477 B1

Val Tyr Leu Leu Pro Lys Lys Met Asp Glu Tyr Val Ala Ser Leu His
 485 490 495

Leu Pro Ser Phe Asp Ala His Leu Thr Glu Leu Thr Asp Asp Gln Ala
 500 505 510

Lys Tyr Leu Gly Leu Asn Lys Asn Gly Pro Phe Lys Pro Asn Tyr Tyr
 515 520 525

Arg Tyr
 530

<210> 4
 <211> 2427
 <212> DNA
 <213> Mus musculus
 <400> 4

20 tgctgttgct tctggttctg tggccgccgt cgctgtccgg caggetccgg tctcggagcc 60
 gacgaggcgc gcgagcgagg gtcccagccc ttcggaagcc caagcagctc ggcgcggggc 120
 ctggcgggaa ggcgggagcg cgcgtggccg agggcgccgc gcggacgggc gggcgcccgt 180
 gagggamaga ggcgggggag cggggtagc cgcgggcccgg gccggccggg ggatgtcgat 240
 gcctgacgag atgccgctgc ccggggctgg ggaggagctg aacaggcca aggagatcga 300
 ggacgccgag aagtactcct tcatggccac ggtcaccaag gctcccaga agcaaatcca 360
 gtttgctgat gacatgcaag agttcaccaa attccctact aagactggcc ggagatcttt 420
 gtctcgttcc atctcacaat cctccacaga cagctacagt tcagctgcat cctatacaga 480
 tagctctgat gatgaggttt cccctcgaga gaagcagcaa accaactcga agggcagcag 540
 caatttctgt gtgaagaaca tcaagcaggc agagtttggg cgccgggaga ttgagattgc 600
 agagcaagac atgtctgctc tgatttact caggaaacgt gctcaggagc agaagccttt 660
 ggctggtgct aaaatagtag gctgtacgca catcacggcc cagacagcgg tattaattga 720
 gaccctttgt gccctgggag ctgagtgccc ttggctgccc tgcaacatct attcaactca 780
 30 gaatgaagta gctgcagcac tggctgaggc tggagtcgag gtgtttgctt ggaagggcga 840
 gtcagaagat gatttctggt ggtgcattga ccgctgtgtc aacatggatg ggtggcaggc 900
 taacatgatc ctggatgatg ggggagactt aaccactgg gtttataaga agtatccaaa 960
 cggtgttaag aagatccgag gcattgtgga agagagcgtg actggtgttc acaggctgta 1020
 tcagctctcc aaagctggga agctctgtgt tccagccatg aatgtcaatg attctgttac 1080
 caaacagaag tttgataacc tgtactgctg ccgagaatcc attttggatg gcctgaagag 1140
 35 gaccacggat gtgatgtttg gtgggaaaca ggtgggtggg tgtggctatg gtgaggtagg 1200
 aaagggctgc tgtgctgctc tcaagggcct tggagcaatt gtctacataa cagaaattga 1260
 ccccatctgt gctctgcagg cctgcatgga tgggttcagg gtgggtgaagc tgaatgaagt 1320
 catccggcag gtggacgctg taattacttg cacaggaat aagaatgtag tgactcggga 1380
 gcacttggac cgaatgaaaa atagttgtat tgtgtgcaat atgggccatt ccaacacgga 1440
 40 gatcgagctg accagcctcc gactccaga actaacatgg gagcgtgtac gttctcagg 1500
 ggaccatgtc atctggcctg atggcaaacg ggtcgtcctt cttagcagagg gccgtttact 1560
 taatctgagc tgctccacag tccctacctt tgttctttcc atcacggcta caacacaggc 1620
 tttggcactg atagagcttt acaacgccc ggagggagcg tacaacagg atgtgtactt 1680
 gcttcctaag aagatggatg aatatgttgc cagcttgcac ytaccatcat ttgatgccca 1740
 cctgacagaa ctgacagatg accaagcaaa gtatctggga ctcaacaaaa atgggccatt 1800
 45 caaacctaata tattacagat actaatggac atagtacagt gaccagtcca catgaaccac 1860
 gcaactctaa tagagtattt ttaagataa cttttatttt cttcttatta ctttctttt 1920
 gatttttttt ttctatcatt tcattgttgt tttctcatct catcatttga gttttgcaga 1980
 ccacacagga acttgctcca tagctcttta ggtgaaactg aggtcaaagg tttctcacc 2040
 aagtcactaa aaggggttaa ctctgctgcc cagaaaagtt attctttaac catttctggg 2100
 aactttgatc gtatttagtt accttattaa cagaaaatgc taaggcatct tctatgtgga 2160
 50 acaatctaca gtgtctaaat tgccttaaaa gagcctgttc cttagctgctg gaactagtgc 2220
 tctttcactt cttcagagga gccaggatgg tacttcccag ccaggtaggt tagatgtagg 2280
 tgggtcatgt cagcttccca tagacactct aagccctgtt tcctgtgtaa ggtgggyatg 2340
 tctggcagag atgcgctgct tgttcaactc agtaggttca ctggggttg tagtccagcc 2400
 ttccaccagt ctctctcatt gttctag 2427

<210> 5
 <211> 21
 <212> RNA
 <213> Artificial sequence

<220>

<223> siRNA

<400> 5

5 aaauccaguu ugcugaugac a 21

<210> 6

<211> 21

<212> RNA

10 <213> Artificial Sequence

<220>

<223> siRNA

15 <400> 6

aacucagaau gaaguagcug c 21

Claims

- 20
1. Use of inositol 1,4,5-triphosphate (IP₃) receptor-binding protein (IRBIT) for controlling phosphatidylinositol metabolism in a cell in-vitro in which the PIP₂ synthetase isoform α , β or γ PIPKII (Type II phosphatidylinositol-5-phosphate 4-kinase) is involved, wherein the IRBIT is a protein comprising an amino acid sequence shown in SEQ ID NO: 1 or SEQ ID NO: 3 or a protein comprising an amino acid sequence having 90% or more identity with the amino acid sequence of SEQ ID NO: 1 or SEQ ID NO: 3 and having the same biological activity as IRBIT.
 - 25
 2. Use of a DNA encoding IRBIT or vector comprising the DNA for controlling phosphatidylinositol metabolism in a cell in-vitro in which the PIP₂ synthetase isoform α , β or γ PIPKII is involved, wherein the IRBIT is a protein comprising an amino acid sequence shown in SEQ ID NO: 1 or SEQ ID NO: 3 or a protein comprising an amino acid sequence having 90% or more identity with the amino acid sequence of SEQ ID NO: 1 or SEQ ID NO: 3 and having the same biological activity as IRBIT, and wherein the nucleic acid is selected from the group consisting of DNAs encoding IRBIT.
 - 30
 3. Use of siRNA for IRBIT for controlling phosphatidylinositol metabolism in a cell in vitro in which the PIP₂ synthetase isoform α , β or γ PIPKII is involved, wherein the IRBIT is a protein comprising an amino acid sequence shown in SEQ ID NO: 1 or SEQ ID NO: 3 or a protein comprising an amino acid sequence having 90% or more identity with the amino acid sequence of SEQ ID NO: 1 or SEQ ID NO: 3 and having the same biological activity as IRBIT, and wherein the siRNA consists of the nucleotide sequence of SEQ ID NO: 5 or 6.
 - 35
 4. Use of an antibody against IRBIT or a fragment thereof, capable of binding specifically to IRBIT, for controlling phosphatidylinositol metabolism in a cell in vitro in which the PIP₂synthetase isoform α , β or γ PIPKII is involved, wherein the IRBIT is a protein comprising an amino acid sequence shown in SEQ ID NO: 1 or SEQ ID NO: 3 or a protein comprising an amino acid sequence having 90% or more identity with the amino acid sequence of SEQ ID NO: 1 or SEQ ID NO: 3 and having the same biological activity as IRBIT, and wherein the fragment is selected from the group consisting of Fab, F(ab')₂, Fv and scFv.
 - 40
 - 45
 5. The use according to any one of claims 1-4, wherein the control is suppression or elevation.
 6. The use according to claim 4, wherein the antibody is a monoclonal antibody, a polyclonal antibody, a recombinant antibody, or a single chain antibody.
 - 50
 7. The use according to any one of claims 1 and 4, wherein the IRBIT is phosphorylated.

55 Patentansprüche

1. Verwendung von Inositol-1,4,5-triphosphat(IP₃)-rezeptorbindendem Protein (IRBIT) für die Steuerung des Phosphatidylinositolmetabolismus in einer Zelle in vitro, an dem die PIP₂-Synthetaseisoform α -, β - oder γ -PIPKII (Typ-

II-Phosphatidylinositol-5-phosphat-4-Kinase) beteiligt ist, wobei es sich bei dem IRBIT um ein Protein, umfassend eine Aminosäuresequenz gemäß SEQ ID NO: 1 oder SEQ ID NO: 3 oder ein Protein, umfassend eine Aminosäuresequenz mit 90% oder mehr Identität zu der Aminosäuresequenz von SEQ ID NO: 1 oder SEQ ID NO: 3 aufweist und das über dieselbe biologische Aktivität wie IRBIT verfügt, handelt.

- 5 2. Verwendung einer DNA, die für IRBIT codiert, oder eines Vektors, der die DNA umfasst, für die Steuerung des Phosphatidylinositolmetabolismus in einer Zelle *in vitro*, an dem die PIP₂-Synthetaseisoform α -, β - oder γ -PIPKII beteiligt ist, wobei es sich bei dem IRBIT um ein Protein, umfassend eine Aminosäuresequenz gemäß SEQ ID NO: 1 oder SEQ ID NO: 3 oder ein Protein, umfassend eine Aminosäuresequenz mit 90% oder mehr Identität zu der Aminosäuresequenz von SEQ ID NO: 1 oder SEQ ID NO: 3 aufweist und das über dieselbe biologische Aktivität wie IRBIT verfügt, handelt, und wobei die Nukleinsäure aus der Gruppe bestehend aus DNAs, die für IRBIT codieren, ausgewählt ist.
- 10 3. Verwendung von siRNA für IRBIT für die Steuerung des Phosphatidylinositolmetabolismus in einer Zelle *in vitro*, an dem die PIP₂-Synthetaseisoform α -, β - oder γ -PIPKII beteiligt ist, wobei es sich bei dem IRBIT um ein Protein, umfassend eine Aminosäuresequenz gemäß SEQ ID NO: 1 oder SEQ ID NO: 3 oder ein Protein, umfassend eine Aminosäuresequenz mit 90% oder mehr Identität zu der Aminosäuresequenz von SEQ ID NO: 1 oder SEQ ID NO: 3 aufweist und das über dieselbe biologische Aktivität wie IRBIT verfügt, handelt, und wobei die siRNA aus der Nukleotidsequenz von SEQ ID NO: 5 oder 6 besteht.
- 15 4. Verwendung eines Antikörpers gegen IRBIT oder eines Fragments davon, das fähig ist, spezifisch an IRBIT zu binden, für die Steuerung des Phosphatidylinositolmetabolismus in einer Zelle *in vitro*, an dem die PIP₂-Synthetaseisoform α -, β - oder γ -PIPKII beteiligt ist, wobei es sich bei dem IRBIT um ein Protein, umfassend eine Aminosäuresequenz gemäß SEQ ID NO: 1 oder SEQ ID NO: 3 oder ein Protein, umfassend eine Aminosäuresequenz mit 90% oder mehr Identität zu der Aminosäuresequenz von SEQ ID NO: 1 oder SEQ ID NO: 3 aufweist und das über dieselbe biologische Aktivität wie IRBIT verfügt, handelt, und wobei das Fragment aus der Gruppe bestehend aus Fab, F(ab')₂, Fv und scFv ausgewählt ist.
- 20 5. Verwendung nach einem der Ansprüche 1-4, wobei es sich bei der Steuerung um Suppression oder Erhöhung handelt.
- 25 6. Verwendung nach Anspruch 4, wobei es sich bei dem Antikörper um einen monoklonalen Antikörper, einen polyclonalen Antikörper, einen rekombinanten Antikörper oder einen Einzelkettenantikörper handelt.
- 30 7. Verwendung nach einem der Ansprüche 1 und 4, wobei das IRBIT phosphoryliert ist.

Revendications

- 40 1. Utilisation de protéine de liaison de récepteur d'inositol 1,4,5-triphosphate (IP₃) (IRBIT) pour contrôler le métabolisme du phosphatidylinositol dans une cellule *in vitro* dans laquelle l'isoforme de PIP₂ synthétase α , β ou γ PIPKII (phosphatidylinositol-5-phosphate 4-kinase de type II) est impliquée, l'IRBIT étant une protéine comprenant une séquence d'acides aminés décrite dans SEQ ID NO: 1 ou SEQ ID NO: 3 ou une protéine comprenant une séquence d'acides aminés ayant 90 % ou plus d'identité avec la séquence d'acides aminés de SEQ ID NO: 1 ou SEQ ID NO: 3 et ayant la même activité biologique qu'IRBIT.
- 45 2. Utilisation d'un ADN codant pour IRBIT ou vecteur comprenant l'ADN pour contrôler le métabolisme du phosphatidylinositol dans une cellule *in vitro* dans laquelle l'isoforme de PIP₂ synthétase α , β ou γ PIPKII est impliquée, l'IRBIT étant une protéine comprenant une séquence d'acides aminés décrite dans SEQ ID NO: 1 ou SEQ ID NO: 3 ou une protéine comprenant une séquence d'acides aminés ayant 90 % ou plus d'identité avec la séquence d'acides aminés de SEQ ID NO: 1 ou SEQ ID NO: 3 et ayant la même activité biologique qu'IRBIT, et l'acide nucléique étant choisi dans le groupe constitué d'ADN codant pour IRBIT.
- 50 3. Utilisation d'ARNsi pour IRBIT pour contrôler le métabolisme du phosphatidylinositol dans une cellule *in vitro* dans laquelle l'isoforme de PIP₂ synthétase α , β ou γ PIPKII est impliquée, l'IRBIT étant une protéine comprenant une séquence d'acides aminés décrite dans SEQ ID NO: 1 ou SEQ ID NO: 3 ou une protéine comprenant une séquence d'acides aminés ayant 90 % ou plus d'identité avec la séquence d'acides aminés de SEQ ID NO: 1 ou SEQ ID NO: 3 et ayant la même activité biologique qu'IRBIT, et l'ARNsi étant constitué de la séquence nucléotidique de SEQ
- 55

EP 2 311 477 B1

ID NO: 5 ou 6.

- 5
4. Utilisation d'un anticorps contre IRBIT ou un fragment de celui-ci, capable de se lier spécifiquement à IRBIT, pour contrôler le métabolisme du phosphatidylinositol dans une cellule *in vitro* dans laquelle l'isoforme de PIP₂ synthétase α , β ou γ PIPKII est impliquée, l'IRBIT étant une protéine comprenant une séquence d'acides aminés décrite dans SEQ ID NO: 1 ou SEQ ID NO: 3 ou une protéine comprenant une séquence d'acides aminés ayant 90 % ou plus d'identité avec la séquence d'acides aminés de SEQ ID NO: 1 ou SEQ ID NO: 3 et ayant la même activité biologique qu'IRBIT, et le fragment étant choisi dans le groupe constitué de Fab, F(ab')₂, Fv et scFv.
- 10
5. Utilisation selon l'une quelconque des revendications 1 à 4, dans laquelle le contrôle est la suppression ou l'élévation.
6. Utilisation selon la revendication 4, dans laquelle l'anticorps est un anticorps monoclonal, un anticorps polyclonal, un anticorps recombinant, ou un anticorps monocaténaire.
- 15
7. Utilisation selon l'une quelconque des revendications 1 et 4, dans laquelle l'IRBIT est phosphorylé.

20

25

30

35


40

45

50

55

Fig. 1



- 1: IRBIT alone
- 2: CPSF complex alone
- 3: IRBIT+CPSF
- 4: IRBIT+CPSF-160 kDa subunit
- 5: IRBIT+CPSF-100 kDa subunit
- 6: IRBIT+CPSF-73 kDa subunit
- 7: IRBIT+CPSF-30 kDa subunit

Fig. 2




Fig. 3




Fig. 4




Fig. 5





Fig. 6






Fig. 8





Fig. 9

A



B





Fig. 10

A



B





Fig. 11

A



B




Fig. 12




Fig. 13





Fig. 14

A



B





Fig. 15



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- JP 2004129612 A [0005] [0006] [0007] [0046] [0057] [0138]
- JP 2006077607 A [0024] [0173]
- EP 1408049 A [0037]
- US 5252479 A [0063]
- WO 9413788 A [0063]
- JP 9216825 A [0082]
- JP 2002542805 A [0082]
- JP 2092812 A [0093]
- WO 9824893 A [0093]
- WO 9634096 A [0093]
- JP 2003527832 A [0094]

Non-patent literature cited in the description

- **M. J. BERRIDGE.** *Nature*, 1993, vol. 361, 315-325 [0002]
- **M. J. BERRIDGE et al.** *Nat. Rev. Mol. Cell Biol.*, 2000, vol. 1, 11-21 [0002]
- **T. FURUICHI ; K. MIKOSHIBA.** *J. Neurochem.*, 1995, vol. 64, 953-960 [0002]
- **T. FURUICHI et al.** *Nature*, 1989, vol. 342, 32-38 [0003]
- **T. SUDHOF et al.** *EMBO J.*, 1991, vol. 10, 3199-3206 [0003]
- **O. BLONDEL et al.** *J. Biol. Chem.*, 1993, vol. 268, 11356-11363 [0003]
- **P. F. WORLEY et al.** *Nature*, 1987, vol. 325, 159-161 [0003]
- **T. FURUICHI et al.** *Recept. Channels*, 1993, vol. 1, 11-24 [0003]
- **H. ANDO et al.** *J. Biol. Chem.*, 2003, vol. 278, 10602-10612 [0005] [0046]
- **C. BARNARD DARON et al.** *Cell*, 2004, vol. 119, 641-651 [0031]
- Synaptic Plasticity and Translation Initiation. **E. KLANN et al.** *Learning & Memory*. Cold Spring Harbor Laboratory Press, 2004, vol. 11, 365-372 [0031]
- **E. KLANN et al.** *Cytoplasmic polyadenylation and CPEB*, 367-368 [0031]
- **I. KAUFMANN et al.** *EMBO J.*, 2004, vol. 23, 616-626 [0033] [0145]
- **KATIA A. LAMIA et al.** *Mol. Cell Biol.*, 2004, vol. 24, 5080-5087 [0037]
- **E. GROSS ; I. KURTZ.** *Am. J. Physiol. Renal Physiol.*, 2002, vol. 283, F876-F887 [0042]
- **SETH L. ALPER.** *Annu. Rev. Physiol.*, 2002, vol. 64, 899-923 [0044] [0152]
- Short Protocols in Molecular Biology. **F. M. AUSBEL et al.** *A Compendium of Methods from Current Protocols in Molecular Biology*. John Wiley & Sons, Inc, 1995 [0048] [0049]
- **J. SAMBROOK et al.** *Molecular Cloning A Laboratory Manual*. Cold Spring Harbor Laboratory Press, 1989 [0049]
- *Experimental Medicine. New Genetic Engineering Handbook*. YODOSHA, 2003 [0049] [0056]
- *Frontiers of PCR from Basic Technology to Application. PCR techniques, Protein, Nucleic Acid, and Enzyme*. KYORITSU SHUPPAN, April 1996, vol. 41 [0053]
- **C. -S. YUAN et al.** *J. Biol. Chem.*, 1996, vol. 271, 28009-28016 [0058]
- **MAMORU NAKANISHI et al.** *Protein, Nucleic Acid, and Enzyme*. KYORITSU SHUPPAN, 1999, vol. 44, 48-54 [0065] [0074]
- *Clinical Cancer research*, 1999, vol. 59, 4325-4333 [0065] [0074]
- **WU et al.** *J. Biol. Chem.*, 1987, vol. 262, 4429 [0065] [0074]
- **L. ZENDER et al.** *Proc. Natl. Acad. Sci. U.S.A.*, 2003, vol. 100, 77797-7802 [0065]
- **H. XIA et al.** *Nature Biotech.*, 2002, vol. 20, 1006-1010 [0065]
- **X. F. QIN et al.** *Proc. Natl. Acad. Sci. U.S.A.*, 2003, vol. 100, 183-188 [0065]
- **G. M. BARTON et al.** *Proc. Natl. Acad. Sci. U.S.A.*, 2002, vol. 99, 14943-14945 [0065]
- **J. D. HOMMEL et al.** *Nature Med.*, 2003, vol. 9, 1539-1544 [0065]
- **D. M. DYKXHOORN et al.** *Nature Rev. Mol. Cell Biol.*, 2003, vol. 77, 7174-7181 [0072]
- **A. KHVOROVA et al.** *Cell*, 2003, vol. 115, 209-216 [0072]
- **J. A. JAEGER et al.** *Methods in Enzymology*, 1989, vol. 183, 281-306 [0072]
- **D. H. MATHEWS et al.** *J. Mol. Biol.*, 1999, vol. 288, 911-940 [0072]
- **T. R. BRUMMELKAMP et al.** *Science*, 2002, vol. 296, 550-553 [0079]

EP 2 311 477 B1

- **N. S. LEE et al.** *Nature Biotech.*, 2002, vol. 20, 500-505 [0079]
- **M. MIYAGISHI et al.** *Nat. Biotechnol.*, 2002, vol. 20, 497-500 [0079]
- **P. J. PADDISON et al.** *Genes & Dev.*, 2002, vol. 16, 948-958 [0079]
- **T. TUSCH.** *Nature Biotech.*, 2002, vol. 20, 446-448 [0079]
- **C. P. PAUL et al.** *Nature Biotech.*, 2002, vol. 20, 505-508 [0079]
- RNAi Experimental Protocols. YODOSHA, 2003 [0079]
- **S. KOSEKI et al.** *J. Virol.*, 1999, vol. 73, 1868-1877 [0083]
- *Nature*, 1975, vol. 256, 495-497 [0089]
- **P. J. DELVES et al.** ANTIBODY PRODUCTION ESSENTIAL TECHNIQUES. John Wiley&Sons, 1997 [0091]
- **T. C. THOMAS et al.** *Mol. Immunol.*, 1996, vol. 33, 1389-1401 [0092]
- **I. ISHIDA et al.** *Cloning Stem Cell*, 2002, vol. 4, 91-102 [0092]
- Experimental Medicine. New genetic engineering handbook. YODOSHA, 2003 [0126]
- **DARON C. BARNARD et al.** *Cell*, 2004, vol. 119, 641-651 [0142]