

(19)



(11)

EP 2 012 128 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention of the grant of the patent:
22.05.2013 Bulletin 2013/21

(51) Int Cl.:
G01N 33/574 (2006.01)

(21) Application number: **07740484.6**

(86) International application number:
PCT/JP2007/057048

(22) Date of filing: **30.03.2007**

(87) International publication number:
WO 2007/119606 (25.10.2007 Gazette 2007/43)

(54) METHOD FOR DETERMINATION OF PROGNOSIS OF PROSTATE CANCER, AND DIAGNOSTIC AGENT FOR USE IN THE METHOD

VERFAHREN ZUR BESTIMMUNG DER PROGNOSE VON PROSTATAKREBS UND DIAGNOSTISCHES MITTEL ZUR VERWENDUNG IN DEM VERFAHREN

MÉTHODE POUR DÉTERMINER UN PRONOSTIC DE CANCER DE LA PROSTATE, ET AGENT DIAGNOSTIQUE À UTILISER AVEC LA MÉTHODE

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IS IT LI LT LU LV MC MT NL PL PT RO SE SI SK TR

(56) References cited:
JP-A- 2002 511 564 JP-A- 2004 532 390
US-A1- 2005 026 196

(30) Priority: **05.04.2006 JP 2006104315**

(43) Date of publication of application:
07.01.2009 Bulletin 2009/02

(73) Proprietor: **Yokohama City University**
Yokohama-shi
Kanagawa 236-0027 (JP)

(72) Inventors:
• **RYO, Akihide**
Yokohama-shi, Kanagawa 236-0004 (JP)
• **AOKI, Ichiro**
Yokohama-shi, Kanagawa 236-0004 (JP)
• **SASAKI, Takeshi**
Yokohama-shi, Kanagawa 232-0024 (JP)

(74) Representative: **Kador, Utz Ulrich**
Kador & Partner
Corneliusstrasse 15
80469 München (DE)

- **RYO A ET AL: "REGULATION OF NF-KB SIGNALING BY PIN1-DEPENDENT PROLYL ISOMERIZATION AND UBIQUITIN-MEDIATED PROTEOLYSIS OF P65/REIA" MOLECULAR CELL, CELL PRESS, CAMBRIDGE, MA, US, vol. 12, 1 December 2003 (2003-12-01), pages 1413-1426, XP008056951 ISSN: 1097-2765**
- **ANDELA VALENTINE B ET AL: "NFKappaB: a pivotal transcription factor in prostate cancer metastasis to bone." CLINICAL ORTHOPAEDICS AND RELATED RESEARCH OCT 2003, no. 415 Suppl, October 2003 (2003-10), pages S75-S85, XP009116083 ISSN: 0009-921X**
- **LESSARD L ET AL: "Nuclear localisation of nuclear factor-kappaB transcription factors in prostate cancer: an immunohistochemical study" BRITISH JOURNAL OF CANCER, vol. 93, no. 9, October 2005 (2005-10), pages 1019-1023, XP002526077 ISSN: 0007-0920**
- **HUANG SUYUN ET AL: "Blockade of NF-kappaB activity in human prostate cancer cells is associated with suppression of angiogenesis, invasion, and metastasis" ONCOGENE, vol. 20, no. 31, 12 July 2001 (2001-07-12), pages 4188-4197, XP002526078 ISSN: 0950-9232**

EP 2 012 128 B1

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

Technical Field

5 **[0001]** The present invention relates to a method for determining probability that a prostate cancer will metastasize.

Background Art

10 **[0002]** In the U.S., prostate cancer has the highest prevalence among cancers in males. In Japan too, the prevalence of the prostate cancer is consistently increasing due to the change of lifestyle such as westernization of dietary life (Non-patent Literature 1). Since no specific symptoms are manifested in the early stage of prostate cancer, and since prostate cancer is often complicated with benign prostatic hyperplasia, the symptoms similar to those of benign prostatic hyperplasia, and dysuria, nocturia, urinary urgency and the like are found in some cases. Therefore, recently, even though no symptom is observed, most of prostate cancer is detected by measurement of PSA prostate-specific antigen

15 in the blood in medical examination or health screening. Prostate cancer is characterized by high frequency of metastasis, and in some cases, advanced cancer is found because of low back pain due to bone metastasis (Non-patent Literature 2). **[0003]** As the factor deciding the prognosis of prostate cancer, stage (in the following) thereof before surgery is important.

20 Stage A: cases where prostate cancer was accidentally found in surgery of benign prostatic hyperplasia or bladder cancer

Stage B: cases where the cancer is confined in the prostate

Stage C: cases where the cancer is not metastasized to other organs (such as bone, lymph node, lung and liver), but the cancer is exposed to the outside of the coating membrane (the membrane coating the periphery of prostate)

25 Stage D: cases where the cancer is metastasized to other organs such as bone and lymph node

[0004] For the cases of Stage A to Stage C, basically, total extirpation of prostate is carried out. However, in a considerable number of cases, recurrence of the cancer within several years after the surgery is observed. Especially, refractoriness to hormone therapy and metastasis are important factors which aggravate the survival prognosis, so that to estimate these factors at an early stage has a great clinical significance in the selection of therapy or treatment of the patient. However, up to now, no molecular marker is known which enables to estimate the prognosis of prostate cancer, especially the metastasis thereof, at an early stage.

30 **[0005]** Andela et al investigated the potential role of the transcription factor NF- κ B as a central regulator of prostate cancer metastasis using the prostate adenocarcinoma cell line, PC-3, in a series of in vitro studies. These in vitro experiments suggest NF- κ B as an important target to prevent prostate cancer bone metastasis ("NF- κ B: a pivotal transcription factor in prostate cancer metastasis to bone." Clinical orthopaedics and related research, number 415S, October 2003 (2003-10), pages S75-S85).

35 **[0006]** On the other hand, it is known that blockade of NF- κ B (nuclear factor, NF)- κ B activity, which is one of the transcription factors, inhibits angiogenesis, infiltration and metastasis in human prostate cancer (Non-patent Literature 3). More particularly, Non-patent Literature 3 discloses that when a mutated I κ B α gene having an activity to inhibit NF- κ B activity was introduced into metastatic human prostate cancer cells PC-3M and the cells were transplanted to nude mice, angiogenesis, infiltration and metastasis were inhibited in the group wherein NF- κ B activity was reduced by the introduction of the mutated I κ B α gene when compared with the group wherein the human prostate cancer cells PC-3M into which the mutated I κ B α gene was not introduced were transplanted. Non-patent Literature 4 discloses that in prostate cancer cells, expression of NF- κ B significantly occurs in the nuclei.

45 **[0007]** US 2005-026196 A1 discloses a method for predicting whether a patient suffering from a prostate cancer is at risk of experiencing a cancer progression or recurrence which comprises the steps of: obtaining a Gleason score of said prostate tumour sample; assessing the proportion of NF- κ B localized in the nuclei of said tumour sample with regard to all NF- κ B present in said tumour sample. Non-patent Literature 5 discloses that by the action of Pin1 which is a type of peptidyl-prolyl isomerase that specifically acts on the sequence of phosphorylated threonine or serine and subsequent proline, which sequence exists in a protein, on p65/RelA which is a subunit of NF- κ B and in which the 254th amino acid threonine is phosphorylated, the binding of NF- κ B-p65/RelA to I κ B α that is an inhibition factor of NF- κ B is inhibited, so that nuclear accumulation and stability of NF- κ B-p65/RelA are increased and, in turn, the NF- κ B activity is increased.

50 **[0008]**

55

Non-patent Literature 1: Wakai K. Descriptive epidemiology of prostate cancer in Japan and Western countries Nippon Rinsho. 2005 Feb;63(2):207-12.

Non-patent Literature 2: Loberg RD, Gayed BA, Olson KB, Pienta KJ. A paradigm for the treatment of prostate

cancer bone metastases based on an understanding of tumor cell-microenvironment interactions. J Cell Biochem. 2005 Oct 15;96(3):439-46.

Non-patent Literature 3: Huang S, Pettaway CA, Uehara H, Bucana CD, Fidler IJ. Blockade of NF-kappaB activity in human prostate cancer cells is associated with suppression of angiogenesis, invasion, and metastasis. Oncogene. 2001 Jul 12;20(31):4188-97.

Non-patent Literature 4: Lessard L, Begin LR, Gleave ME, Mes-Masson AM, Saad F. Nuclear localisation of nuclear factor-kappaB transcription factors in prostate cancer: an immunohistochemical study. Br J Cancer. 2005 Oct 31; 93(9):1019-23.

Non-patent Literature 5: Ryo A, Suizu F, Yoshida Y, Perrem K, Liou YC, Wulf G, Rottapel R, Yamaoka S, Lu KP. Regulation of NF-kappaB signaling by Pin1-dependent prolyl isomerization and ubiquitin-mediated proteolysis of p65/RelA. Mol Cell. 2003 Dec;12(6):1413-26.

Disclosure of the Invention

Problems to be Solved by the Invention

[0009] An object of the present invention is to provide a method for determining probability that a prostate cancer will metastasize and a diagnostic reagent therefor. Means for Solving the Problems

[0010] The present inventors intensively studied to discover that the percentage that NF-κB-p65/RelA has the 254th amino acid threonine which is phosphorylated is significantly higher in the prostate cancer cells in the cases where the bone metastasis was observed than in the cases where bone metastasis was not observed, and that the probability that a prostate cancer will metastasize can be determined at an early stage by examining whether the 254th amino acid threonine in NF-κB-p65/RelA is phosphorylated or not, thereby completing the present invention.

[0011] That is, the present invention provides a method for determining probability that a prostate cancer will metastasize, the method comprising measuring human NF-κB-p65/RelA in which 254th amino acid threonine is phosphorylated, in a prostate tissue separated from human. The present invention can be utilized for a diagnostic reagent for determining probability that a prostate cancer will metastasize, the reagent comprising an antibody or an antigen-binding fragment thereof, the antibody undergoing antigen-antibody reaction with human NF-κB-p65/RelA in which 254th amino acid threonine is phosphorylated but not undergoing antigen-antibody reaction with human NF-κB-p65/RelA in which 254th amino acid threonine is not phosphorylated.

Effects of the Invention

[0012] By the present invention, means by which the probability that a prostate cancer will metastasize can be determined at an early stage was first provided. By the present invention, since the probability that a prostate cancer will metastasize can be determined at an early stage, therapy or treatment of the patient can be properly selected, so that the present invention is expected to greatly contribute to the therapy of prostate cancer.

Brief Description of the Drawings

[Fig. 1]

[0013] A figure which comparatively shows the ratio of strongly positive, positive and negative results of immunohistochemical staining of prostate tissues in the cases where bone metastasis was observed and not observed, respectively, which immunohistochemical staining was carried in the Example of the present invention. Best Mode for Carrying out the Invention

[0014] NF-κB is a transcription factor which is expressed widely in most cells. Since it is expressed in nuclei of cells, it is called nuclear factor. NF-κB-p65/RelA is one of the subunits of NF-κB. The amino acid sequence of human NF-κB-p65/RelA and the cDNA sequence encoding it are known, and are shown in GenBank Accession No. M62399. The base sequence of the cDNA of human NF-κB-p65/RelA and the amino acid sequence encoded thereby are shown in SEQ ID NO: 1 in SEQUENCE LISTING, and the amino acid sequence alone is shown in SEQ ID NO: 2.

[0015] In the method of the present invention, the human NF-κB-p65/RelA in which 254th amino acid threonine is phosphorylated (hereinafter also referred to as "254 phosphorylated NF-κB-p65/RelA" for convenience). The term "threonine is phosphorylated" means that the hydroxyl group in the side chain of threonine is converted to a phosphoric acid ester. Phosphorylated threonine *per se* is well-known. In the present invention, the term "measure" includes any of detection, semi-quantification and quantification. The term "measure the 254 phosphorylated NF-κB-p65/RelA" means to measure the 254 phosphorylated NF-κB-p65/RelA distinctly from the NF-κB-p65/RelA whose 254th threonine is not phosphorylated (hereinafter also referred to as "254 unphosphorylated NF-κB-p65/RelA" for short).

[0016] An example of the preferred method for measuring the 254 phosphorylated NF- κ B-p65/RelA is an immunoassay using an antibody (hereinafter also referred to as "254 phosphorylated NF- κ B-p65/RelA-specific antibody") or an antigen-binding fragment thereof, which antibody undergoes antigen-antibody reaction with 254 phosphorylated NF- κ B-p65/RelA but does not undergo antigen-antibody reaction with 254 unphosphorylated NF- κ B-p65/RelA.

5 **[0017]** The 254 phosphorylated NF- κ B-p65/RelA-specific antibody may be a polyclonal antibody or a monoclonal antibody. The 254 phosphorylated NF- κ B-p65/RelA-specific polyclonal antibody can be obtained by preparing a polyclonal antibody by a conventional method using as an immunogen 254 phosphorylated NF- κ B-p65/RelA or a fragment thereof containing the 254th threonine; and removing, from the obtained polyclonal antibody, by adsorption the antibodies which also undergo antigen-antibody reaction with 254 unphosphorylated NF- κ B-p65/RelA (see the Example below).

10 The monoclonal antibody can be obtained by preparing hybridomas each producing an anti-254 phosphorylated NF- κ B-p65/RelA monoclonal antibody by a conventional method using as an immunogen 254 phosphorylated NF- κ B-p65/RelA or a fragment thereof containing the 254th threonine; selecting, from the obtained hybridomas, those producing the 254 phosphorylated NF- κ B-p65/RelA-specific antibody; culturing the selected hybridomas *in vitro* or in abdominal cavity of an animal, or the like; and recovering the 254 phosphorylated NF- κ B-p65/RelA-specific antibody produced by the hybridomas from the culture medium or ascites. In the immunoassay, the 254 phosphorylated NF- κ B-p65/RelA-specific antibody may be used, or a fragment thereof having a binding activity to the antigen, such as Fab fragment or F(ab')₂ fragment (in the present invention, referred to as "antigen-binding fragment") may also be used.

15 **[0018]** As the immunogen used for the preparation of the anti-254 phosphorylated NF- κ B-p65/RelA antibody, the whole molecule of 254 phosphorylated NF- κ B-p65/RelA or a fragment thereof containing the 254th threonine may be used, as mentioned above. In the latter case, the size of the peptide fragment is preferably not less than 10 amino acids, more preferably not less than 12 amino acids. Since a 254 phosphorylated NF- κ B-p65/RelA-specific antibody was obtained in the Example described below using a fragment having 12 amino acids ((VFRT(PO3H2)PPYADPSC) (SEQ ID NO: 3), the "(PO3H2)" after T means that the hydroxyl group in the side chain of this threonine (T) is phosphorylated), a size of 12 amino acids is sufficient. Needless to say, however, those having a longer size (the longest is the whole molecule of 254 phosphorylated NF- κ B-p65/RelA) can also be used as the immunogen. The peptide fragment of the 254 phosphorylated NF- κ B-p65/RelA which can be used as the immunogen can easily be chemically synthesized using a commercially available peptide synthesizer. Since the phosphorylated threonine *per se* is commercially available, a peptide fragment containing the phosphorylated threonine as the 254th threonine can be synthesized by using the commercially available threonine as a material in the step of binding the 254th threonine when the peptide is chemically synthesized. In cases where a peptide fragment is used as the immunogen, the peptide fragment may be used as the immunogen as it is, or the peptide fragment bound to a protein carrier such as bovine serum albumin, casein or keyhole limpet hemocyanin, may be used as the immunogen. Since a 254 phosphorylated NF- κ B-p65/RelA-specific antibody was obtained in the Example below by administering a peptide fragment consisting of 12 amino acids as it is, binding to the carrier protein is not indispensable.

30 **[0019]** The immunoassay can be carried out by immunohistochemical staining to a prostate tissue separated from the living body, or by a well-known immunoassay such as sandwich method, competition method or agglutination method using as a test sample a cell homogenate of a prostate tissue separated from the living body or the purified or partially purified product thereof containing NF- κ B-p65/RelA. Among these methods, immunohistochemical staining is preferred. The immunohistochemical staining can be carried out by a conventional method except that the 254 phosphorylated NF- κ B-p65/RelA-specific antibody is used as the antibody (see the Example below). Since a kit for immunohistochemical staining is commercially available, the immunohistochemical staining can be easily carried out using the commercially available kit and the above-described 254 phosphorylated NF- κ B-p65/RelA-specific antibody as the antibody.

35 **[0020]** As will be concretely described in the Example below, in the prostate cancer tissue metastasized to bone, the percentage that the reactivity with the 254 phosphorylated NF- κ B-p65/RelA-specific antibody is positive (positive rate) was statistically significantly higher than in the cases where bone metastasis was not observed. Moreover, among the cases where the 254 phosphorylated NF- κ B-p65/RelA was positive, in the cases where the abundance of 254 phosphorylated NF- κ B-p65/RelA was especially high (in the cases where the immunohistochemical staining was strongly positive), the percentage of bone metastasis was especially high. Further, in most cases where bone metastasis occurred, the immunohistochemical staining was positive irrespective of the histological malignancy (Gleason grade). These results show that the 254 phosphorylated NF- κ B-p65/RelA can be utilized as a prostate cancer metastasis marker indicating the probability of metastasis of prostate cancer, and whether metastasis will occur in the future or not can be estimated irrespective of the histological malignancy.

40 **[0021]** The present invention will now be described more concretely by way of an example thereof. However, the present invention is not restricted to the Example below.

55

Example

Materials and Methods

5 1. Preparation of 254 phosphorylated NF- κ B-p65/RelA-specific Antibody

[0022] A phosphorylated peptide (VFRT(PO3H2)PPYADPSC) (SEQ ID NO: 3) which was a fragment of p65/RelA, containing the 254th threonine (12 amino acids from the 251st valine to 262nd cysteine) wherein the 254th threonine had a phosphate group attached thereto was chemically synthesized using a peptide synthesizer. As the phosphorylated threonine to which a phosphate group was added, a commercially available product was used. The obtained peptide fragment was mixed with an equal amount of Freund complete adjuvant (first immunization) or Freund incomplete adjuvant (second and subsequent immunization), and an emulsion was prepared by sonication. The resulting viscous emulsion was subcutaneously administered to the back of a rabbit (New Zealand white rabbit). The immunization was conducted totally 4 times at 2-week intervals. At the time of third immunization, a small amount of the antigen protein was sampled, and antibody titer was measured by ELISA. Two weeks after the fourth immunization, the total blood was collected from the carotid artery under anesthesia with ketamine-xylazine. The collected blood was left to stand at room temperature for 1 hour, and then at 4°C overnight, followed by centrifugation at 3000 rpm for 10 minutes to obtain a serum.

[0023] Using a commercially available antibody-purification kit (Amersham Bio., HiTrap Protein A HP), antibody was purified from the obtained antiserum. The buffers used were as follows:

20	Binding buffer	20 mM sodium phosphate, pH7.0
	Elution buffer	0.1 M sodium citrate, pH3.0
	Neutralization buffer	1.0 M Tris-HCl, pH9.0

[0024] Concrete purification operation was as follows:

- (1) Preparation of column: A syringe was connected without introducing bubbles into the column, and 25 mL of ultrapure water was applied at a rate of 5 drops/second.
- (2) Equilibration of Column: 25 mL of binding buffer was applied at a rate of 5 drops/second.
- (3) Addition of sample: The prepared sample (dilution of antiserum) was applied at a rate of 5 drops/second, and the adsorbed components were washed, followed by application of 25 mL of binding buffer at a rate of 5 drops/second.
- (4) Elution of antibody: 25 mL of elution buffer was applied at a rate of 5 drops/second, the eluted solution was collected in 3 mL fractions (300 μ L of neutralization buffer was added to each collection tube), the concentration of each of the eluted fractions was measured, the absorbance at 280 nm of each fraction was measured, the antibody fraction was recovered and dialyzed to PBS(-) overnight to exchange the buffer.

[0025] From the obtained polyclonal antibody, the antibody which underwent antigen-antibody reaction also with 254 unphosphorylated NF- κ B-p65/RelA was removed by adsorption. More particularly, a biotinylated 254 unphosphorylated NF- κ B-p65/RelA (VFRTPPYADPSC) (SEQ ID NO: 3) chemically synthesized with a peptide synthesizer was bound to commercially available streptavidin-magnetic beads (amount of total bound antigen: 0.01 mg; amount of beads: 0.2 mg), and the beads were mixed with 10-fold diluted antibody solution (0.3 mL) at 4°C for 2 hours. Then the non-specific antibody adsorbed to the unphosphorylated peptide was removed using a magnet, and the supernatant was recovered to obtain a 254 phosphorylated NF- κ B-p65/RelA-specific antibody.

45 2. Immunohistochemical Staining

[0026] Using a commercially available immunohistochemical staining kit (VECTASTAIN ABC Kit), immunohistochemical staining with the 254 phosphorylated NF- κ B-p65/RelA-specific antibody obtained in the above-described 1 was performed on prostate tissues separated from patients. The concrete operation of the immunohistochemical staining was as follows:

[0027]

- (1) Deparaffinization and hydrophilization of each tissue section with xylene and ethanol
- (2) Washing twice (each for 5 minutes) with 100 mL each of Tris buffered physiological saline (TBS).
- (3) Inactivation of endogenous peroxidase (1 hour) with 60 mL of TBS supplemented with 2 mL of 30% H₂O₂ (1%)
- (4) Washing twice (each for 5 minutes) with 100 mL each of TBS.
- (5) Blocking (1 hour) with goat serum in TBS (3 mL) (final concentration: 10%)

- (6) Washing once (for 5 minutes) with 100 mL of TBST (0.1% Tween 20 (trade name)-containing TBS)
- (7) Primary antibody (254 phosphorylated NF- κ B-p65/RelA-specific antibody prepared in the above-described 1) diluted (50-fold to 150-fold) with the blocking solution was added to the sample and the resultant was left to stand at room temperature for 2 hours or at 4°C overnight.
- 5 (8) Washing twice (each for 5 minutes) with 100 mL each of TBS
- (9) Secondary antibody (500 μ L of TBS supplemented with 2 μ L of biotinylated secondary antibody (goat anti-rabbit IgG antibody, included in the kit)) was added to the sample and the resultant was allowed to react for 2 hours.
- (10) Washing twice (each for 5 minutes) with 100 mL each of TBS
- 10 (11) Preparation of AB solution (10 mL of TBS supplemented with 2 drops each of A: avidin solution (C solution) and B: biotinylated peroxidase (D solution)). AB solution was prepared 30 minutes before use and allowed to react.
- (12) AB solution is added to the sample, and the resultant is allowed to react for 2 hours.
- (13) Washing twice (each for 5 minutes) with 100 mL each of TBS
- 15 (14) DAB solution (TBS (5 mL) containing 50 μ g (10 μ g/mL) of diaminobenzidine (DAB) which is a substrate of peroxidase and 5 μ L (0.03%) of H₂O₂ (produced by VECTOR) is added to the sample, and the resultant is allowed to react for about 5 to 10 minutes in the dark. Coloring is checked sometimes.
- (15) Washing once (for 5 minutes) with 100 mL of TBS
- (16) Staining of nuclei of cells in blue with hematoxylin reagent
- (17) Ethanol substitution, dropping xylene and a sealant, and seal with a cover glass

20 3. Materials

[0028] Prostate cancer operation samples extirpated in Yokohama City University Hospital and related hospitals and pre-operation needle biopsies (total 49 cases, breakdown was as follows:)

Cases where bone metastasis occurred within 3 to 5 years (27 cases)

25 Cases where bone metastasis did not occur in the period mentioned above (22 cases)

Bone metastatic site samples of prostate cancer (6 cases)

4. Results

30 **[0029]** Immunostaining was performed on the prostate cancer tissue samples by immunohistochemistry with the antibody. The samples used were the needle biopsy tissue samples or the extirpated prostate samples of the cases (27 cases) where bone metastasis occurred during the follow-up period of 3 to 5 years (including the cases where bone metastasis was found at the first visit), and the cases (22 cases) where bone metastasis did not occur during the above-mentioned period. Bone metastatic site samples of prostate cancer (6 cases) were also stained, and evaluation at the metastatic site was also conducted.

35 **[0030]** As a result of immunohistochemical staining, in prostate cancer cases where bone metastasis occurred, positive image (brown) was observed in the primary focus and in the metastatic focus in the prostate cancer cells. In the staining pattern, strong staining was observed chiefly in the nuclei. Further, irrespective of histological malignancy (Gleason grade), in most of the bone metastasis cases, the results were positive and in most of the cases where bone metastasis did not occur, the results were negative.

40 **[0031]** The results of staining were classified into 3 grades, namely, negative, positive, and strongly positive by two or more pathologists. The negative grade was the cases where staining with the antibody was not observed; the strongly positive grade was the cases where strong staining with the antibody was observed; and the positive grade was the cases where staining with the antibody was observed, but the degree of staining was not so strong as the strongly positive grade.

45 **[0032]** The results are shown in Table 1 below and Fig. 1. As shown in Table 1 and Fig. 1, irrespective of the histological malignancy, in the cases where bone metastasis occurred, 19 cases were strongly positive grade, 8 cases were positive grade and 0 case was negative grade, so that the antibody reaction was positive in 100% of the cases. On the other hand, in the cases where bone metastasis did not occur, 1 case was strongly positive grade, 6 cases were positive grade and 15 cases were negative grade, so that most of them was negative grade. Since the follow-up period is short, it is thought that there is a possibility that in the strongly positive case and the positive cases in the cases where bone metastasis did not occur, bone metastasis may occur in the future. Further, in the site of bone metastasis, among the 6 cases, 3 cases were strongly positive grade and 3 cases were positive grade, so that the positive rate was 100% too. To determine whether there is a significant difference between the groups with bone metastasis and without bone metastasis in the results of staining with the antibody, a statistical test (chi square test) was carried out. As a result, in the bone metastasis-positive group, the staining with the antibody was significantly strongly positive or positive with a 99% significant difference ($P < 0.01$ ($\chi^2_{(2)} = 31.2 > \chi^2_{(2)}(0.01) = 9.21$). By these results, it was proved that the antibody can be used as a molecular marker (diagnostic reagent) for determining the probability that the prostate cancer will metas-

tasized.
[0033]

Table 1

5

Immunohistochemical Staining	Number of Cases	
	without metastasis	with metastasis
Strongly Positive	1	19
Positive	6	8
Negative	15	0
Total	22	27

10

15

SEQUENCE LISTING

[0034]

<110> Yokohama City University

20

<120> Method for predicting probability of metastasis of prostate cancer and diagnostic drug therefor

<130> 06PF342-PCT

25

<160> 3

<170> PatentIn version 3.1

30

<210> 1

<211> 1767

<212> DNA

<213> Homo sapiens

35

<220>

<221> CDS

<222> (80)..(1735)

<223>

40

<400> 1

45

50

55

EP 2 012 128 B1

gaattccggc gaatggctcg tctgtagtgc acgccgcggg cccagctgcg accccggccc 60
 cgccccggg accccggcc atg gac gaa ctg ttc ccc ctc atc ttc ccg gca 112
 5 Met Asp Glu Leu Phe Pro Leu Ile Phe Pro Ala
 1 5 10
 gag cca gcc cag gcc tct ggc ccc tat gtg gag atc att gag cag ccc 160
 Glu Pro Ala Gln Ala Ser Gly Pro Tyr Val Glu Ile Ile Glu Gln Pro
 15 20 25
 10 aag cag cgg ggc atg cgc ttc cgc tac aag tgc gag ggg cgc tcc gcg 208
 Lys Gln Arg Gly Met Arg Phe Arg Tyr Lys Cys Glu Gly Arg Ser Ala
 30 35 40
 ggc agc atc cca ggc gag agg agc aca gat acc acc aag acc cac ccc 256
 Gly Ser Ile Pro Gly Glu Arg Ser Thr Asp Thr Thr Lys Thr His Pro
 15 45 50 55
 acc atc aag atc aat ggc tac aca gga cca ggg aca gtg cgc atc tcc 304
 Thr Ile Lys Ile Asn Gly Tyr Thr Gly Pro Gly Thr Val Arg Ile Ser
 60 65 70 75
 20 ctg gtc acc aag gac cct cct cac cgg cct cac ccc cac gag ctt gta 352
 Leu Val Thr Lys Asp Pro Pro His Arg Pro His Pro His Glu Leu Val
 80 85 90
 gga aag gac tgc cgg gat ggc ttc tat gag gct gag ctc tgc ccg gac 400
 Gly Lys Asp Cys Arg Asp Gly Phe Tyr Glu Ala Glu Leu Cys Pro Asp
 25 95 100 105
 cgc tgc atc cac agt ttc cag aac ctg gga atc cag tgt gtg aag aag 448
 Arg Cys Ile His Ser Phe Gln Asn Leu Gly Ile Gln Cys Val Lys Lys
 110 115 120
 30 cgg gac ctg gag cag gct atc agt cag cgc atc cag acc aac aac aac 496
 Arg Asp Leu Glu Gln Ala Ile Ser Gln Arg Ile Gln Thr Asn Asn Asn
 125 130 135
 ccc ttc caa gtt cct ata gaa gag cag cgt ggg gac tac gac ctg aat 544
 Pro Phe Gln Val Pro Ile Glu Glu Gln Arg Gly Asp Tyr Asp Leu Asn
 35 140 145 150 155
 gct gtg cgg ctc tgc ttc cag gtg aca gtg cgg gac cca tca ggc agg 592
 Ala Val Arg Leu Cys Phe Gln Val Thr Val Arg Asp Pro Ser Gly Arg

40

45

50

55

EP 2 012 128 B1

	160					165					170						
5	ccc Pro	ctc Leu	cgc Arg	ctg Leu 175	ccg Pro	cct Pro	gtc Val	ctt Leu	cct Pro 180	cat His	ccc Pro	atc Ile	ttt Phe	gac Asp 185	aat Asn	cg Arg	640
	gcc Ala	ccc Pro	aac Asn 190	act Thr	gcc Ala	gag Glu	ctc Leu	aag Lys 195	atc Ile	tgc Cys	cga Arg	gtg Val	aac Asn 200	cga Arg	aac Asn	tct Ser	688
10	ggc Gly	agc Ser 205	tgc Cys	ctc Leu	ggt Gly	ggg Gly	gat Asp 210	gag Glu	atc Ile	ttc Phe	cta Leu	ctg Leu 215	tgt Cys	gac Asp	aag Lys	gtg Val	736
15	cag Gln 220	aaa Lys	gag Glu	gac Asp	att Ile	gag Glu 225	gtg Val	tat Tyr	ttc Phe	acg Thr	gga Gly 230	cca Pro	ggc Gly	tgg Trp	gag Glu	gcc Ala 235	784
	cga Arg	ggc Gly	tcc Ser	ttt Phe	tcg Ser 240	caa Gln	gct Ala	gat Asp	gtg Val	cac His 245	cga Arg	caa Gln	gtg Val	gcc Ala	att Ile 250	gtg Val	832
20	ttc Phe	cg Arg	acc Thr	cct Pro 255	ccc Pro	tac Tyr	gca Ala	gac Asp	ccc Pro 260	agc Ser	ctg Leu	cag Gln	gct Ala	cct Pro 265	gtg Val	cg Arg	880
25	gtc Val	tcc Ser	atg Met 270	cag Gln	ctg Leu	cg Arg	cg Arg	cct Pro 275	tcc Ser	gac Asp	cg Arg	gag Glu	ctc Leu 280	agt Ser	gag Glu	ccc Pro	928
	atg Met	gaa Glu 285	ttc Phe	cag Gln	tac Tyr	ctg Leu	cca Pro 290	gat Asp	aca Thr	gac Asp	gat Asp	cg Arg 295	cac His	cg Arg	att Ile	gag Glu	976
30	gag Glu 300	aaa Lys	cg Arg	aaa Lys	agg Arg	aca Thr 305	tat Tyr	gag Glu	acc Thr	ttc Phe	aag Lys 310	agc Ser	atc Ile	atg Met	aag Lys	aag Lys 315	1024
35	agt Ser	cct Pro	ttc Phe	agc Ser	gga Gly 320	ccc Pro	acc Thr	gac Asp	ccc Pro	cg Arg 325	cct Pro	cca Pro	cct Pro	cga Arg	cg Arg 330	att Ile	1072
	gct Ala	gtg Val	cct Pro	tcc Ser 335	cg Arg	agc Ser	tca Ser	gct Ala	tct Ser 340	gtc Val	ccc Pro	aag Lys	cca Pro	gca Ala 345	ccc Pro	cag Gln	1120
40	ccc Pro	tat Tyr	ccc Pro 350	ttt Phe	acg Thr	tca Ser	tcc Ser	ctg Leu 355	agc Ser	acc Thr	atc Ile	aac Asn	tat Tyr 360	gat Asp	gag Glu	ttt Phe	1168
45	ccc Pro	acc Thr 365	atg Met	gtg Val	ttt Phe	cct Pro	tct Ser 370	ggg Gly	cag Gln	atc Ile	agc Ser	cag Gln 375	gcc Ala	tcg Ser	gcc Ala	ttg Leu	1216
	gcc Ala 380	ccg Pro	gcc Ala	cct Pro	ccc Pro	caa Gln 385	gtc Val	ctg Leu	ccc Pro	cag Gln 390	gct Ala	cca Pro	gcc Ala	cct Pro	gcc Ala	cct Pro 395	1264
50	gct Ala	cca Pro	gcc Ala	atg Met	gta Val 400	tca Ser	gct Ala	ctg Leu	gcc Ala	cag Gln 405	gcc Ala	cca Pro	gcc Ala	cct Pro	gtc Val 410	cca Pro	1312
	gtc Val	cta Leu	gcc Ala	cca Pro 415	ggc Gly	cct Pro	cct Pro	cag Gln 420	gct Ala	gtg Val	gcc Ala	cca Pro	cct Pro	gcc Ala 425	ccc Pro	aag Lys	1360
55	ccc Pro	acc Thr	cag Gln	gct Ala	ggg Gly	gaa Glu	gga Gly	acg Thr	ctg Leu	tca Ser	gag Glu	gcc Ala	ctg Leu	ctg Leu	cag Gln	ctg Leu	1408

EP 2 012 128 B1

	430	435	440	
5	cag ttt gat gat gaa gac ctg ggg gcc ttg ctt ggc aac agc aca gac			1456
	Gln Phe Asp Asp Glu Asp Leu Gly Ala Leu Leu Gly Asn Ser Thr Asp			
	445	450	455	
10	cca gct gtg ttc aca gac ctg gca tcc gtc gac aac tcc gag ttt cag			1504
	Pro Ala Val Phe Thr Asp Leu Ala Ser Val Asp Asn Ser Glu Phe Gln			
	460	465	470	
15	cag ctg ctg aac cag ggc ata cct gtg gcc ccc cac aca act gag ccc			1552
	Gln Leu Leu Asn Gln Gly Ile Pro Val Ala Pro His Thr Thr Glu Pro			
	480	485	490	
20	atg ctg atg gag tac cct gag gct ata act cgc cta gtg aca ggg gcc			1600
	Met Leu Met Glu Tyr Pro Glu Ala Ile Thr Arg Leu Val Thr Gly Ala			
	495	500	505	
25	cag agg ccc ccc gac cca gct cct gct cca ctg ggg gcc ccg ggg ctc			1648
	Gln Arg Pro Pro Asp Pro Ala Pro Ala Pro Leu Gly Ala Pro Gly Leu			
	510	515	520	
30	ccc aat ggc ctc ctt tca gga gat gaa gac ttc tcc tcc att gcg gac			1696
	Pro Asn Gly Leu Leu Ser Gly Asp Glu Asp Phe Ser Ser Ile Ala Asp			
	525	530	535	
35	atg gac ttc tca gcc ctg ctg agt cag atc agc tcc taa gggggtgacg			1745
	Met Asp Phe Ser Ala Leu Leu Ser Gln Ile Ser Ser			
	540	545	550	
40	cctgccctcc ccagagcact gg			1767
30	<210> 2			
	<211> 551			
	<212> PRT			
	<213> Homo sapiens			
35	<400> 2			
40	Met Asp Glu Leu Phe Pro Leu Ile Phe Pro Ala Glu Pro Ala Gln Ala			
	1 5 10 15			
45	Ser Gly Pro Tyr Val Glu Ile Ile Glu Gln Pro Lys Gln Arg Gly Met			
	20 25 30			
50	Arg Phe Arg Tyr Lys Cys Glu Gly Arg Ser Ala Gly Ser Ile Pro Gly			
	35 40 45			
55	Glu Arg Ser Thr Asp Thr Thr Lys Thr His Pro Thr Ile Lys Ile Asn			
	50 55 60			
60	Gly Tyr Thr Gly Pro Gly Thr Val Arg Ile Ser Leu Val Thr Lys Asp			
	65 70 75 80			
65	Pro Pro His Arg Pro His Pro His Glu Leu Val Gly Lys Asp Cys Arg			
	85 90 95			
70	Asp Gly Phe Tyr Glu Ala Glu Leu Cys Pro Asp Arg Cys Ile His Ser			
	100 105 110			

EP 2 012 128 B1

Phe Gln Asn Leu Gly Ile Gln Cys Val Lys Lys Arg Asp Leu Glu Gln
 115 120 125
 5 Ala Ile Ser Gln Arg Ile Gln Thr Asn Asn Asn Pro Phe Gln Val Pro
 130 135 140
 10 Ile Glu Glu Gln Arg Gly Asp Tyr Asp Leu Asn Ala Val Arg Leu Cys
 145 150 155 160
 Phe Gln Val Thr Val Arg Asp Pro Ser Gly Arg Pro Leu Arg Leu Pro
 165 170 175
 15 Pro Val Leu Pro His Pro Ile Phe Asp Asn Arg Ala Pro Asn Thr Ala
 180 185 190
 20 Glu Leu Lys Ile Cys Arg Val Asn Arg Asn Ser Gly Ser Cys Leu Gly
 195 200 205
 Gly Asp Glu Ile Phe Leu Leu Cys Asp Lys Val Gln Lys Glu Asp Ile
 210 215 220
 25 Glu Val Tyr Phe Thr Gly Pro Gly Trp Glu Ala Arg Gly Ser Phe Ser
 225 230 235 240
 30 Gln Ala Asp Val His Arg Gln Val Ala Ile Val Phe Arg Thr Pro Pro
 245 250 255
 Tyr Ala Asp Pro Ser Leu Gln Ala Pro Val Arg Val Ser Met Gln Leu
 260 265 270
 35 Arg Arg Pro Ser Asp Arg Glu Leu Ser Glu Pro Met Glu Phe Gln Tyr
 275 280 285
 40 Leu Pro Asp Thr Asp Asp Arg His Arg Ile Glu Glu Lys Arg Lys Arg
 290 295 300
 Thr Tyr Glu Thr Phe Lys Ser Ile Met Lys Lys Ser Pro Phe Ser Gly
 305 310 315 320
 45 Pro Thr Asp Pro Arg Pro Pro Pro Arg Arg Ile Ala Val Pro Ser Arg
 325 330 335
 50 Ser Ser Ala Ser Val Pro Lys Pro Ala Pro Gln Pro Tyr Pro Phe Thr
 340 345 350
 Ser Ser Leu Ser Thr Ile Asn Tyr Asp Glu Phe Pro Thr Met Val Phe
 355 360 365
 55 Pro Ser Gly Gln Ile Ser Gln Ala Ser Ala Leu Ala Pro Ala Pro Pro
 370 375 380

EP 2 012 128 B1

Gln Val Leu Pro Gln Ala Pro Ala Pro Ala Pro Ala Pro Ala Met Val
 385 390 395 400
 5 Ser Ala Leu Ala Gln Ala Pro Ala Pro Val Pro Val Leu Ala Pro Gly
 405 410 415
 10 Pro Pro Gln Ala Val Ala Pro Pro Ala Pro Lys Pro Thr Gln Ala Gly
 420 425 430
 15 Glu Gly Thr Leu Ser Glu Ala Leu Leu Gln Leu Gln Phe Asp Asp Glu
 435 440 445
 20 Asp Leu Gly Ala Leu Leu Gly Asn Ser Thr Asp Pro Ala Val Phe Thr
 450 455 460
 25 Asp Leu Ala Ser Val Asp Asn Ser Glu Phe Gln Gln Leu Leu Asn Gln
 465 470 475 480
 30 Gly Ile Pro Val Ala Pro His Thr Thr Glu Pro Met Leu Met Glu Tyr
 485 490 495
 35 Pro Glu Ala Ile Thr Arg Leu Val Thr Gly Ala Gln Arg Pro Pro Asp
 500 505 510
 40 Pro Ala Pro Ala Pro Leu Gly Ala Pro Gly Leu Pro Asn Gly Leu Leu
 515 520 525
 45 Ser Gly Asp Glu Asp Phe Ser Ser Ile Ala Asp Met Asp Phe Ser Ala
 530 535 540 545
 50 Leu Leu Ser Gln Ile Ser Ser
 545 550

<210> 3
 <211> 12
 40 <212> PRT
 <213> Artificial

<220>
 <223> peptide fragment used as an immunogen
 45
 <400> 3

Val Phe Arg Thr Pro Pro Tyr Ala Asp Pro Ser Cys
 1 5 10

Claims

1. A method for determining probability that a prostate cancer will metastasize, said method comprising measuring human NF- κ B-p65/RelA in which 254th amino acid threonine is phosphorylated, in a prostate tissue separated from human.
2. The method according to claim 1, carried out by an immunoassay using an antibody or an antigen-binding fragment

thereof, said antibody undergoing antigen-antibody reaction with human NF- κ B-p65/RelA in which 254th amino acid threonine is phosphorylated but not undergoing antigen-antibody reaction with human NF- κ B-p65/RelA in which 254th amino acid threonine is not phosphorylated.

- 5 3. The method according to claim 2, wherein said immunoassay is immunohistochemical staining.
4. The method according to according to any one of claims 1 to 3, wherein the metastasis is bone metastasis.

10 **Patentansprüche**

1. Verfahren zum Bestimmen der Wahrscheinlichkeit, daß Prostatakrebs Metastasen bildet, wobei dieses Verfahren das Messen von Human-NF- κ B-p65/RelA, worin die 254. Aminosäure Threonin phosphoryliert ist, in einem vom Menschen gewonnenen Prostatagewebe aufweist.
- 15 2. Verfahren nach Anspruch 1, das durch einen Immunoassay durchgeführt wird, wobei ein Antikörper oder ein Antigen bindendes Fragment davon verwendet wird, wobei der Antikörper eine Antigen-Antikörper-Reaktion mit Human-NF- κ B-p65/RelA, worin die 254. Aminosäure Threonin phosphoryliert ist, jedoch keine Antigen-Antikörper-Reaktion mit Human-NF- κ B-p65/RelA eingeht, worin die 254. Aminosäure Threonin nicht phosphoryliert ist.
- 20 3. Verfahren nach Anspruch 2, wobei der Immunoassay das Immunohistochemische Anfärben ist.
4. Verfahren nach einem der Ansprüche 1 bis 3, wobei die Metastasen Knochenmetastasen sind.

25

Revendications

1. Procédé destiné à déterminer la probabilité qu'un cancer de la prostate se métastase, ledit procédé comprenant la mesure du NF - κ B - p65 / RelA humain dans lequel la thréonine d'acide aminé en position 254 est phosphorylée, dans un tissu de prostate séparé d'un corps humain.
- 30 2. Procédé selon la revendication 1, exécuté avec un immunoessai en utilisant un anticorps ou un fragment de liaison antigène de celui-ci, ledit anticorps subissant une réaction antigène-anticorps avec le NF - κ B - p65 / RelA humain dans lequel la thréonine d'acide aminé en position 254 est phosphorylée, mais ne subissant pas une réaction antigène-anticorps avec le NF - κ B - p65 / RelA humain dans lequel la thréonine d'acide aminé en position 254 n'est pas phosphorylée.
- 35 3. Procédé selon la revendication 2, dans lequel ledit immunoessai est une coloration immunohistochimique.
- 40 4. Procédé selon l'une quelconque des revendications 1 à 3, dans lequel la métastase est une métastase osseuse.

45

50

55




Fig.1

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 2005026196 A1 [0007]

Non-patent literature cited in the description

- NF- κ B: a pivotal transcription factor in prostate cancer metastasis to bone. *Clinical orthopaedics and related research*, October 2003, S75-S85 [0005]
- **WAKAI K.** Descriptive epidemiology of prostate cancer in Japan and Western countries. *Nippon Rinsho*, February 2005, vol. 63 (2), 207-12 [0008]
- **LOBERG RD ; GAYED BA ; OLSON KB ; PIANTA KJ.** A paradigm for the treatment of prostate cancer bone metastases based on an understanding of tumor cell-microenvironment interactions. *J Cell Biochem.*, 15 October 2005, vol. 96 (3), 439-46 [0008]
- **HUANG S ; PETTAWAY CA ; UEHARA H ; BUCANA CD ; FIDLER IJ.** Blockade of NF- κ B activity in human prostate cancer cells is associated with suppression of angiogenesis, invasion, and metastasis. *Oncogene*, 12 July 2001, vol. 20 (31), 4188-97 [0008]
- **LESSARD L ; BEGIN LR ; GLEAVE ME ; MES-MASSON AM ; SAAD F.** Nuclear localisation of nuclear factor- κ B transcription factors in prostate cancer: an immunohistochemical study. *Br J Cancer*, 31 October 2005, vol. 93 (9), 1019-23 [0008]
- **RYO A ; SUIZU F ; YOSHIDA Y ; PERREM K ; LIOU YC ; WULF G ; ROTTAPPEL R ; YAMAOKA S ; LU KP.** Regulation of NF- κ B signaling by Pin1-dependent prolyl isomerization and ubiquitin-mediated proteolysis of p65/RelA. *Mol Cell*, December 2003, vol. 12 (6), 1413-26 [0008]