



US007960530B2

(12) **United States Patent**
Miyawaki et al.

(10) **Patent No.:** **US 7,960,530 B2**

(45) **Date of Patent:** **Jun. 14, 2011**

(54) **FLUORESCENT PROTEIN**
(75) Inventors: **Atsushi Miyawaki**, Saitama (JP);
Hidekazu Tsutsui, Saitama (JP);
Satoshi Karasawa, Tokyo (JP)

WO 03/104460 12/2003
WO 03/104461 12/2003
WO 2004/018671 3/2004
WO 2004/111235 12/2004
WO 2005/054464 6/2005

(73) Assignee: **Riken**, Saitama (JP)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 1009 days.

(21) Appl. No.: **10/561,040**

(22) PCT Filed: **Jun. 16, 2004**

(86) PCT No.: **PCT/JP2004/008790**

§ 371 (c)(1),
(2), (4) Date: **Oct. 16, 2006**

(87) PCT Pub. No.: **WO2004/111236**

PCT Pub. Date: **Dec. 23, 2004**

(65) **Prior Publication Data**

US 2007/0072259 A1 Mar. 29, 2007

(30) **Foreign Application Priority Data**

Jun. 16, 2003 (JP) 2003-170330

(51) **Int. Cl.**

C07H 21/04 (2006.01)

C12P 21/06 (2006.01)

(52) **U.S. Cl. ... 536/23.5**; 530/350; 435/69.1; 435/320.1;
435/325

(58) **Field of Classification Search** 536/23.5;
530/350; 435/69.1, 320.1, 325
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

2003/0157643 A1* 8/2003 Almond et al. 435/69.1
2005/0032085 A1 2/2005 Labas et al.
2005/0106661 A1 5/2005 Miyawaki et al.
2006/0154296 A1 7/2006 Miyawaki et al.
2006/0240472 A1 10/2006 Miyawaki et al.

FOREIGN PATENT DOCUMENTS

WO 02/42323 5/2002
WO 03/033693 4/2003
WO 03/042401 5/2003
WO 03/054191 7/2003

OTHER PUBLICATIONS

Gurskaya et al., "Color transitions in coral's fluorescent proteins by site-directed mutagenesis," *BMC Biochemistry*, vol. 2, No. 6, Jul. 10, 2001, 7 pages.

Labas et al., "Diversity and evolution of the green fluorescent protein family," *Proceedings of the National Academy of Sciences U.S.A.*, vol. 99, No. 7, Apr. 2, 2002, pp. 4256-4261.

Yang et al., "The molecular structure of green fluorescent protein," *Nature Biotechnology*, vol. 14, No. 10, Oct. 1996, pp. 1246-1251.

Matz et al., "Family of the green fluorescent protein: journey to the end of the rainbow," *BioEssays*, vol. 24, 2002, pp. 953-959.

Miyawaki, "Green Fluorescent Protein-like Proteins in Reef Anthozoa Animals," *Cell Structure and Function*, vol. 27, 2002, pp. 343-347.

Campbell et al., "A monomeric red fluorescent protein," *Proceedings of the National Academy of Sciences U.S.A.*, vol. 99, No. 12, Jun. 11, 2002 pp. 7877-7882.

Matz et al., "Fluorescent proteins from nonbioluminescent Anthozoa species," *Nature Biotechnology*, vol. 17, No. 10, Oct. 1999, pp. 969-973.

Fradkov et al., "Novel fluorescent protein from *Discosoma* coral and its mutants possesses a unique far-red fluorescence," *FEBS Letters*, vol. 479, No. 3, Aug. 18, 2000, pp. 127-130.

U.S. Appl. No. 10/492,081 (Miyawaki et al.), filed Oct. 10, 2002.

U.S. Appl. No. 10/498,505 (Miyawaki et al.), filed Dec. 20, 2002.

U.S. Appl. No. 10/525,365 (Miyawaki et al.), filed Aug. 22, 2003.

U.S. Appl. No. 10/561,041 (Miyawaki et al.), filed Jun. 16, 2004.

U.S. Appl. No. 10/581,551 (Miyawaki et al.), filed Dec. 3, 2004.

U.S. Appl. No. 11/390,215 (Miyawaki et al.), filed Mar. 28, 2006.

Tsien, R.Y., "The Green Fluorescent Protein," *Annual Review of Biochemistry*, vol. 67, 1998, pp. 509-544.

* cited by examiner

Primary Examiner — Chih-Min Kam

(74) *Attorney, Agent, or Firm* — Greenblum & Bernstein, P.L.C.

(57) **ABSTRACT**

An object of the present invention is to provide a novel fluorescent protein derived from *favia favius*. The present invention provides a fluorescent protein derived from *favia favius* having the following properties:

- (1) an excitation maximum wavelength is 507 nm;
- (2) a fluorescence maximum wavelength is 517 nm;
- (3) a molar absorption coefficient at 482 nm is 80,000;
- (4) a quantum yield is 0.68; and
- (5) pH sensitivity of the fluorescence maximum is stable at pH 5 to pH 11.

3 Claims, 7 Drawing Sheets

Fig. 1




Fig. 2




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8





Fig. 9



1

FLUORESCENT PROTEIN

This application is the National Stage of PCT/JP2004/008790, filed Jun. 16, 2004.

SEQUENCE LISTING

The instant application contains a Sequence Listing which has been submitted in ASCII format via EFS-Web and is hereby incorporated by reference in its entirety. Said ASCII copy, created on Jan. 5, 2011, is named P28994.txt and is 40,504 bytes in size.

TECHNICAL FIELD

The present invention relates to a novel fluorescent protein. More specifically, the present invention relates to a novel fluorescent protein derived from *favia favius* and the use thereof.

BACKGROUND ART

Green fluorescent protein (GFP) derived from *Aequorea victoria*, a jellyfish, has many purposes in biological systems. Recently, various GFP mutants have been produced based on the random mutagenesis and semi-rational mutagenesis, wherein a color is changed, a folding property is improved, luminance is enhanced, or pH sensitivity is modified. Fluorescent proteins such as GFP are fused with other proteins by gene recombinant technique, and monitoring of the expression and transportation of the fusion proteins is carried out.

One of the most commonly used types of GFP mutant is Yellow fluorescent protein (YFP). Among *Aequorea*-derived GFP mutants, YFP exhibits the fluorescence with the longest wavelength. The values ϵ and Φ of the majority of YFPs are 60,000 to 100,000 $M^{-1}cm^{-1}$ and 0.6 to 0.8, respectively (Tsien, R. Y. (1998). *Ann. Rev. Biochem.* 67, 509-544). These values are comparable to those of the general fluorescent group (fluorescein, rhodamine, etc.). Accordingly, improvement of the absolute luminance of YFP is nearly approaching its limit.

In addition, cyan fluorescent protein (CFP) is another example of the GFP mutant. Of this type of protein, ECFP (enhanced cyan fluorescent protein) has been known. Moreover, red fluorescent protein (RFP) has been isolated from sea anemone (*Discosoma* sp.). Of this type of protein, DsRed has been known. Thus, 4 types of fluorescent proteins, that are, green fluorescent protein, yellow fluorescent protein, cyan fluorescent protein, and red fluorescent protein, have successfully been developed. The range of the spectrum has significantly been expanded.

Also, there are many cnidarians which emit fluorescence. Cloning of genes of fluorescent proteins from cnidarians has been attempted, but cloning of more genes is necessary in order to increase repertoire of fluorescent and biochemical properties.

DISCLOSURE OF THE INVENTION

An object of the present invention is to provide a novel fluorescent protein derived from *favia favius*.

In order to achieve the above object, the present inventors have conducted intensive studies. They have designed adequate primers based on the amino acid sequence information of existing fluorescent proteins, amplified the gene obtained from the cDNA library derived from *favia favius* using the aforementioned primers, and then succeeded in

2

cloning a novel gene encoding a fluorescent protein. Further, the present inventors have analyzed fluorescence properties and pH sensitivity of the obtained fluorescent protein derived from *favia favius*. The present invention has been completed based on such findings.

Thus, the present invention provides a fluorescent protein derived from *favia favius* having the following properties:

- (1) an excitation maximum wavelength is 507 nm;
- (2) a fluorescence maximum wavelength is 517 nm;
- (3) a molar absorption coefficient at 482 nm is 80,000;
- (4) a quantum yield is 0.68; and
- (5) pH sensitivity of the fluorescence maximum is stable at pH 5 to pH 11.

Another aspect of the present invention provides a fluorescent protein having any of the following amino acid sequences:

- (a) an amino acid sequence shown in SEQ ID NO: 1; or
- (b) an amino acid sequence comprising a deletion, substitution and/or addition of one or several amino acids with respect to the amino acid sequence shown in SEQ ID NO: 1, and having fluorescent properties.

A further aspect of the present invention provides a fluorescent protein having an amino acid sequence derived from any of the following amino acid sequences by substitution from asparagine to histidine at 62th amino acid residue:

- (a) an amino acid sequence shown in SEQ ID NO: 1; or
- (b) an amino acid sequence comprising a deletion, substitution and/or addition of one or several amino acids with respect to the amino acid sequence shown in SEQ ID NO: 1, and having fluorescent properties.

A further aspect of the present invention provides a fluorescent protein having an amino acid sequence derived from any of the following amino acid sequences by substitution from methionine to valine at 40th amino acid residue, substitution from aspartic acid to histidine at 62th amino acid residue, and substitution from isoleucine to methionine at 198th amino acid residue.

- (a) an amino acid sequence shown in SEQ ID NO: 1; or
- (b) an amino acid sequence comprising a deletion, substitution and/or addition of one or several amino acids with respect to the amino acid sequence shown in SEQ ID NO: 1, and having fluorescent properties.

A further aspect of the present invention provides a fluorescent protein having an amino acid sequence derived from any of the following amino acid sequences by substitution from methionine to isoleucine at 10th amino acid residue, substitution from leucine to valine at 12th amino acid residue, substitution from methionine to valine at 40th amino acid residue, substitution from valine to alanine at 60th amino acid residue, substitution from aspartic acid to histidine at 62th amino acid residue, substitution from tyrosine to asparagine at 119th amino acid residue, substitution from proline to serine at 144th amino acid residue, substitution from arginine to leucine at 197th amino acid residue, and substitution from isoleucine to methionine at 198th amino acid residue:

- (a) an amino acid sequence shown in SEQ ID NO: 1; or
- (b) an amino acid sequence comprising a deletion, substitution and/or addition of one or several amino acids with respect to the amino acid sequence shown in SEQ ID NO: 1, and having fluorescent properties.

A further aspect of the present invention provides a fluorescent protein having an amino acid sequence derived from any of the following amino acid sequences by substitution from methionine to isoleucine at 10th amino acid residue, substitution from methionine to valine at 40th amino acid residue, substitution from valine to alanine at 60th amino acid residue, substitution from aspartic acid to histidine at 62th

3

amino acid residue, substitution from lysine to glutamic acid at 70th amino acid residue, substitution from tyrosine to asparagine at 119th amino acid residue, substitution from arginine to glutamine at 197th amino acid residue, and substitution from isoleucine to methionine at 198th amino acid residue:

- (a) an amino acid sequence shown in SEQ ID NO: 1; or
- (b) an amino acid sequence comprising a deletion, substitution and/or addition of one or several amino acids with respect to the amino acid sequence shown in SEQ ID NO: 1, and having fluorescent properties.

A further aspect of the present invention provides a fluorescent protein having an amino acid sequence derived from any of the following amino acid sequences by substitution from valine to alanine at 60th amino acid residue, substitution from aspartic acid to glycine at 62th amino acid residue, substitution from tyrosine to histidine at 63th amino acid residue, substitution from histidine to leucine at 196th amino acid residue, and substitution from isoleucine to threonine at 198th amino acid residue:

- (a) an amino acid sequence shown in SEQ ID NO: 1; or
- (b) an amino acid sequence comprising a deletion, substitution and/or addition of one or several amino acids with respect to the amino acid sequence shown in SEQ ID NO: 1, and having fluorescent properties.

A further aspect of the present invention provides DNA encoding the protein of the present invention.

A further aspect of the present invention provides any of the following DNA:

- (a) DNA encoding the amino acid sequence shown in SEQ ID NO: 1; or
- (b) DNA encoding an amino acid sequence comprising a deletion, substitution and/or addition of one or several amino acids with respect to the amino acid sequence shown in SEQ ID NO: 1, and encoding a fluorescent protein.

A further aspect of the present invention provides DNA having any of the following nucleotide sequences:

- (a) a nucleotide sequence shown in SEQ ID NO: 2; or
- (b) a nucleotide sequence comprising a deletion, substitution and/or addition of one or several nucleotides with respect to the nucleotide sequence shown in SEQ ID NO: 2, and encoding a fluorescent protein.

A further aspect of the present invention provides DNA comprising any of the following nucleotide sequences:

- (a) a nucleotide sequence shown in SEQ ID NO: 13, 15, 17, 19, or 21; or
- (b) a nucleotide sequence comprising a deletion, substitution and/or addition of one or several nucleotides with respect to the nucleotide sequence shown in SEQ ID NO: 13, 15, 17, 19, or 21, and encoding a fluorescent protein.

In another aspect of the present invention, there is provided a recombinant vector having the DNA of the present invention.

In another aspect of the present invention, there is provided a transformant having the DNA or recombinant vector of the present invention.

In another aspect of the present invention, there is provided a fusion fluorescent protein consisting of the fluorescent proteins of the present invention and another protein.

Preferably, said another protein is one that localizes in the cell, and more preferably one specific to an intracellular organelle.

In another aspect of the present invention, there is provided a method for analyzing the localization or dynamics of a protein in cells, characterized in that the fusion protein of the present invention is allowed to be expressed in cells.

4

In another aspect of the present invention, there is provided a fluorescent reagent kit which comprises any of the fluorescent protein, DNA, recombinant vector, transformant or fusion protein of the present invention.

A further aspect of the present invention provides a method for producing a fluorescent protein capable of green-to-red photoconversion in response to irradiation with light, which comprises introducing substitution from an amino acid corresponding to 62th amino acid residue of the fluorescent protein having the amino acid sequence shown in SEQ ID NO: 1 to histidine into a fluorescent protein incapable of green-to-red photoconversion in response to irradiation with light.

A further aspect of the present invention provides a method for producing a fluorescent protein capable of green-to-red photoconversion in response to irradiation with light, which comprises introducing at least one of the following amino acid substitutions into a fluorescent protein incapable of green-to-red photoconversion in response to irradiation with light;

(1) substitution from an amino acid corresponding to 62th amino acid residue of the fluorescent protein having the amino acid sequence shown in SEQ ID NO: 1 to histidine;

(2) substitution from an amino acid corresponding to 10th amino acid residue of the fluorescent protein having the amino acid sequence shown in SEQ ID NO: 1 to isoleucine;

(3) substitution from an amino acid corresponding to 12th amino acid residue of the fluorescent protein having the amino acid sequence shown in SEQ ID NO: 1 to valine;

(4) substitution from an amino acid corresponding to 40th amino acid residue of the fluorescent protein having the amino acid sequence shown in SEQ ID NO: 1 to valine;

(5) substitution from an amino acid corresponding to 60th amino acid residue of the fluorescent protein having the amino acid sequence shown in SEQ ID NO: 1 to alanine;

(6) substitution from an amino acid corresponding to 70th amino acid residue of the fluorescent protein having the amino acid sequence shown in SEQ ID NO: 1 to glutamic acid;

(7) substitution from an amino acid corresponding to 119th amino acid residue of the fluorescent protein having the amino acid sequence shown in SEQ ID NO: 1 to asparagine;

(8) substitution from an amino acid corresponding to 144th amino acid residue of the fluorescent protein having the amino acid sequence shown in SEQ ID NO: 1 to serine;

(9) substitution from an amino acid corresponding to 197th amino acid residue of the fluorescent protein having the amino acid sequence shown in SEQ ID NO: 1 to leucine or glutamine; or

(10) substitution from an amino acid corresponding to 198th amino acid residue of the fluorescent protein having the amino acid sequence shown in SEQ ID NO: 1 to methionine;

A further aspect of the present invention provides a method for producing a fluorescent protein with an increased rate of green-to-red photoconversion and enhanced fluorescence intensity, which comprises introducing at least one of the following amino acid substitutions in a fluorescent protein:

(1) substitution from an amino acid corresponding to 62th amino acid residue of the fluorescent protein having the amino acid sequence shown in SEQ ID NO: 1 to histidine;

(2) substitution from an amino acid corresponding to 54th amino acid residue of the fluorescent protein having the amino acid sequence shown in SEQ ID NO: 1 to phenylalanine;

(3) substitution from an amino acid corresponding to 69th amino acid residue of the fluorescent protein having the amino acid sequence shown in SEQ ID NO: 1 to valine;

5

(4) substitution from an amino acid corresponding to 87th amino acid residue of the fluorescent protein having the amino acid sequence shown in SEQ ID NO: 1 to tyrosine;

(5) substitution from an amino acid corresponding to 93th amino acid residue of the fluorescent protein having the amino acid sequence shown in SEQ ID NO: 1 to methionine;

(6) substitution from an amino acid corresponding to 109th amino acid residue of the fluorescent protein having the amino acid sequence shown in SEQ ID NO: 1 to methionine;

(7) substitution from an amino acid corresponding to 121th amino acid residue of the fluorescent protein having the amino acid sequence shown in SEQ ID NO: 1 to isoleucine;

(8) substitution from an amino acid corresponding to 140th amino acid residue of the fluorescent protein having the amino acid sequence shown in SEQ ID NO: 1 to valine; or (9) substitution from an amino acid corresponding to 160th amino acid residue of the fluorescent protein having the amino acid sequence shown in SEQ ID NO: 1 to valine;

A further aspect of the present invention provides a method for producing a fluorescent protein capable of purple-to-blue photoconversion in response to irradiation with light, which comprises introducing at least one of the following amino acid substitutions in a fluorescent protein incapable of purple-to-blue photoconversion in response to irradiation with light:

(1) substitution from an amino acid corresponding to 60th amino acid residue of the fluorescent protein having the amino acid sequence shown in SEQ ID NO: 1 to alanine;

(2) substitution from an amino acid corresponding to 62th amino acid residue of the fluorescent protein having the amino acid sequence shown in SEQ ID NO: 1 to glycine;

(3) substitution from an amino acid corresponding to 63th amino acid residue of the fluorescent protein having the amino acid sequence shown in SEQ ID NO: 1 to histidine;

(4) substitution from an amino acid corresponding to 196th amino acid residue of the fluorescent protein having the amino acid sequence shown in SEQ ID NO: 1 to leucine; or

(5) substitution from an amino acid corresponding to 198th amino acid residue of the fluorescent protein having the amino acid sequence shown in SEQ ID NO: 1 to threonine;

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 shows the results of measuring the fluorescence emission spectrum and the excitation spectrum of the fluorescent protein (KkG) derived from *favia favius* of the present invention.

FIG. 2 shows the pH dependence of the fluorescent protein (KkG) derived from *favia favius* of the present invention.

FIG. 3 shows a comparison of amino acid sequences (SEQ ID NOS: 22, 23, 1, 12, 14, 16, 18, 20, respectively, in order of appearance), wherein *** represents chromophore-forming amino acids.

FIG. 4 shows changes in spectral properties via light irradiation at 365 nm.

A1: the absorption spectra of KkG

A2: the absorption spectra of KKH after irradiation with light

A3: changes in the absorption spectra of KikGR via irradiation with light

A4: changes in the excitation spectra and in the fluorescence emission spectra of KikGR via irradiation with light

FIG. 5 shows a comparison of fluorescence intensities in *E. coli* and in HeLa cells.

FIG. 5A left: before irradiation at 365 nm

FIG. 5A right: after irradiation at 365 nm

6

FIG. 5B: expression in HeLa cells

7 hours, 12 hours, and 18 hours after irradiation from the above

475AF20/530DF35 exp Isec Dichroic mirror 430DCLP

Xenon 75W, ND 10%T X10 UplanFI NA0.3

FIG. 6 shows green-to-red photoconversion and comparison of brightness (A: change in KikGR fluorescence intensities; B: an image of KikGR-expressing cells; C: change in Kaede fluorescence intensities; and D: an image of Kaede-expressing cells).

Green 475AF20/530DF35 exp 50 ms

Red 550DF30/575ALP exp 100 ms

Violet 400DF 10 exp 100 ms

Dicroic mirror: 430DCLP

Xenon75W ND10% T

Bin4 Cool SNAP HQ

X40 UApo340/NA1.35

FIG. 7 shows changes in the fluorescence emission spectra of KBL2 via ultraviolet irradiation at 270 nm (including the second-order light from the grating), wherein fluorescence at 380 nm is decreased and fluorescence at 450 nm is increased upon ultraviolet irradiation.

FIG. 8 shows the excitation spectra and the fluorescence emission spectra of KBL2 after photoconversion (including the second-order light from the grating).

FIG. 9 shows changes in the absorption spectra via irradiation at 365 nm (fine line: before irradiation; bold line: after irradiation).

PREFERRED EMBODIMENTS OF THE INVENTION

Hereafter, embodiments of the present invention are described in detail.

(1) Fluorescent Protein of the Present Invention

The fluorescent protein of the present invention is derived from *favia favius*, and has the following properties:

(1) an excitation maximum wavelength is 507 nm;

(2) a fluorescence maximum wavelength is 517 nm;

(3) a molar absorption coefficient at 482 nm is 80,000;

(4) a quantum yield is 0.68; and

(5) pH sensitivity of the fluorescence maximum is stable at pH 5 to pH 11.

Favia favius is one type of corals of the Cnidaria class, the Anthozoa class, the *Hexacorallia* subclass, and the Faviidae order.

The fluorescent protein of the present invention exhibits an excitation maximum wavelength of 507 nm and a fluorescence maximum wavelength of 517 nm as described in the following examples. The molar absorption coefficient at 482 nm is 80,000, and the quantum yield is 0.68. The molar absorption coefficient represents a quantity of absorbed photons per fluorescent molecule. The quantum yield is a numerical value indicating the quantity of absorbed photons that can be emitted as fluorescence.

Specific examples of the fluorescent protein of the present invention include those having any of the following amino acid sequences:

(a) an amino acid sequence shown in SEQ ID NO: 1; or

(b) an amino acid sequence comprising a deletion, substitution and/or addition of one or several amino acids with respect to the amino acid sequence shown in SEQ ID NO: 1, and having fluorescent properties.

The scope of "one or several" in the phrase "an amino acid sequence comprising a deletion, substitution and/or addition of one or several amino acids" used herein is not particularly limited in the present specification. For example, it means 1 to

20, preferably 1 to 10, more preferably 1 to 7, further preferably 1 to 5, and particularly preferably 1 to 3.

The term "having fluorescent properties" used herein covers all of the cases where any fluorescence is given. Various properties such as fluorescence intensity, excitation wavelength, fluorescence wavelength or pH sensitivity, may be changed or may remain unchanged, as compared with the protein having an amino acid sequence shown in SEQ ID NO: 1.

The method of obtaining the fluorescent protein of the present invention is not particularly limited. The protein may be either a protein synthesized by chemosynthesis, or recombinant protein produced by a gene recombination technique.

Where a recombinant protein is produced, it is necessary to obtain DNA encoding the protein. Appropriate primers are designed by using information regarding the amino acid sequence shown in SEQ ID NO: 1 of the sequence listing of the present specification and the nucleotide sequence shown in SEQ ID NO: 2. Using these primers, PCR is carried out by using cDNA library derived from *favia favius* as a template, so that DNA encoding the fluorescent protein of the present invention can be obtained. Where a partial fragment of DNA encoding the fluorescent protein of the present invention are obtained by the above-described PCR, the produced DNA fragments are ligated to one another by a gene recombination technique, so that DNA encoding the desired fluorescent protein can be obtained. The fluorescent protein of the present invention can be produced by introducing this DNA into an appropriate expression system. Expression in an expression system will be described later in the present specification.

(2) DNA of the Present Invention

According to the present invention, a gene encoding the fluorescent protein of the present invention is provided.

Specific example of DNA encoding the fluorescent protein of the present invention may include either one of the following DNA:

(a) DNA encoding the amino acid sequence shown in SEQ ID NO: 1; or

(b) DNA encoding an amino acid sequence comprising a deletion, substitution and/or addition of one or several amino acids with respect to the amino acid sequence shown in SEQ ID NO: 1, and encoding a fluorescent protein.

Further example of DNA encoding the fluorescent protein of the present invention may include either one of the following DNA:

(a) DNA having a nucleotide sequence shown in SEQ ID NO: 2; or

(b) DNA having a nucleotide sequence comprising a deletion, substitution and/or addition of one or several nucleotides with respect of the nucleotide sequence shown in SEQ ID NO: 2, and encoding a fluorescent protein.

The DNA of the present invention can be synthesized by, for example, the phosphoamidite method, or it can also be produced by polymerase chain reaction (PCR) using specific primers. The DNA of the present invention or a fragment thereof is produced by the method described above in the specification.

A method of introducing a desired mutation into a certain nucleic acid sequence is known to a person skilled in the art. For example, known techniques such as a site-directed mutagenesis, PCR using degenerated oligonucleotides, or the exposure of cells containing nucleic acid to mutagens or radioactive rays, are appropriately used, so as to construct DNA having a mutation. Such known techniques are described in, for example, Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory, Cold

Spring Harbor, N.Y., 1989; and Current Protocols in Molecular Biology, Supplements 1 to 38, John Wiley & Sons (1987-1997).

(3) Recombinant Vector of the Present Invention

The DNA of the present invention can be inserted into a suitable vector and used. The type of a vector used in the present invention is not particularly limited. For example, it may be either a vector that can autonomously replicate (e.g., a plasmid, etc.), or vector that is incorporated into the genomes of host cells when it is introduced into the host cells and is then replicated together with the chromosome into which it is incorporated.

The vector used in the present invention is preferably an expression vector. In an expression vector, elements necessary for transcription (e.g., a promoter, etc.) are functionally ligated to the DNA of the present invention. The promoter is a DNA sequence which shows a transcriptional activity in host cells, and it is appropriately selected depending on the type of host cells.

Examples of a promoter which can operate in bacterial cells may include a *Bacillus stearothermophilus* maltogenic amylase gene promoter, a *Bacillus licheniformis* alpha-amylase gene promoter, a *Bacillus amyloliquefaciens* BAN amylase gene promoter, a *Bacillus subtilis* alkaline protease gene promoter, a *Bacillus pumilus* xylosidase gene promoter, P_R and P_L promoters of phage rhamda, and lac, trp and tac promoters of *Escherichia coli*.

Examples of a promoter which can operate in mammalian cells may include an SV40 promoter, an MT-1 (metallothionein gene) promoter, and an adenovirus-2 major late promoter. Examples of a promoter which can operate in insect cells may include a polyhedrin promoter, a P10 promoter, an *Autographa californica* polyhedrosis basic protein promoter, a baculovirus immediate-early gene 1 promoter, and a baculovirus 39K delayed-early gene promoter. Examples of a promoter which can be operate in yeast host cells may include promoters derived from yeast glycolytic genes, an alcohol dehydrogenase gene promoter, a TPI1 promoter, and an ADH2-4c promoter.

Examples of a promoter which can operate in filamentous cells may include an ADH3 promoter and a tpiA promoter.

In addition, an appropriate terminator such as a human growth hormone terminator, or a TP11 terminator or ADH3 terminator for fungal cells, may be functionally bound to the DNA of the present invention, as necessary. The recombinant vector of the present invention may further have elements such as a polyadenylation signal (e.g., one derived from SV40 or the adenovirus 5E1b region), a transcription enhancer sequence (e.g., an SV40 enhancer), or a translation enhancer sequence (e.g., one encoding the adenovirus VA RNA).

The recombinant vector of the present invention may further comprise a DNA sequence which enables the replication of the recombinant vector in host cells. SV40 replication origin is an example of such a sequence (when the host cells are mammalian cells).

The recombinant vector of the present invention may further comprise a selective marker. Examples of such a selective marker may include genes, complements of which are absent from host cells, such as a dihydrofolate reductase (DHFR) gene or a *Shizosaccharomyces pombe* TPI gene, and drug resistant genes such as ampicillin, kanamycin, tetracycline, chloramphenicol, neomycin or hygromycin-resistant genes.

A method for ligating the DNA of the present invention, a promoter and, as desired, a terminator and/or a secretory signal sequence to one another and inserting these items into a suitable vector is known to a person skilled in the art.

(4) Transformant of the Present Invention

A transformant can be produced by introducing the DNA or recombinant vector of the present invention into a suitable host.

Any cell can be used as a host cell into which the DNA or recombinant vector of the present invention is introduced, as long as the DNA construct of the present invention can be expressed therein. Examples of such a cell may include bacteria, yeasts, fungal cells, and higher eukaryotic cells.

Examples of bacteria may include Gram-positive bacteria such as *Bacillus* or *Streptomyces*, and Gram-negative bacteria such as *Escherichia coli*. These bacteria may be transformed by the protoplast method or other known methods, using competent cells.

Examples of mammalian cells may include HEK 293 cells, HeLa cells, COS cells, BHK cells, CHL cells, and CHO cells. A method of transforming mammalian cells and expressing the introduced DNA sequence in the cells is also known. Examples of such a method may include the electroporation, the calcium phosphate method, and the lipofection method.

Examples of yeast cells may include those belonging to *Saccharomyces* or *Shizosaccharomyces*. Examples of such cells may include *Saccharomyces cerevisiae* and *Saccharomyces kluyveri*. Examples of a method of introducing a recombinant vector into yeast host cells may include the electroporation, the spheroplast method, and the lithium acetate method.

Examples of other fungal cells may include those belonging to Filamentous fungi such as *Aspergillus*, *Neurospora*, *Fusarium* or *Trichoderma*. Where Filamentous fungi are used as host cells, transformation can be carried out by incorporating DNA constructs into host chromosomes, so as to obtain recombinant host cells. Incorporation of DNA constructs into the host chromosomes is carried out by known methods, and such known methods may include homologous recombination and heterologous recombination.

Where insect cells are used as host cells, both a vector into which a recombinant gene is introduced and a baculovirus are co-introduced into insect cells, and a recombinant virus is obtained in the culture supernatant of the insect cells. Thereafter, insect cells are infected with the recombinant virus, so as to allow the cells to express proteins (described in, for example, Baculovirus Expression Vectors, A Laboratory Manual; and Current Protocols in Molecular Biology, Bio/Technology, 6, 47 (1988)).

The *Autographa californica* nuclear polyhedrosis virus, which is a virus infecting to insects belonging to *Barathra brassicae*, can be used as baculovirus.

Examples of insect cells used herein may include Sf9 and Sf21, which are *Spodoptera frugiperda* ovarian cells [Baculovirus Expression Vectors, A Laboratory Manual, W. H. Freeman & Company, New York, (1992)], and HiFive (manufactured by Invitrogen), which are *Trichoplusia ni* ovarian cells.

Examples of the method of co-introducing both a vector into which a recombinant gene has been introduced and the above baculovirus into insect cells to prepare a recombinant virus may include the calcium phosphate method and the lipofection method.

The above transformant is cultured in an appropriate nutritive medium under conditions enabling the introduced DNA construct to be expressed. In order to isolate and purify the fusion fluorescent protein of the present invention from the culture product of the transformant, common methods of isolating and purifying proteins may be used.

For example, where the protein of the present invention is expressed in a state dissolved in cells, after completion of the

culture, cells are recovered by centrifugal separation, and the recovered cells are suspended in a water type buffer. Thereafter, the cells are disintegrated using an ultrasonic disintegrator or the like, so as to obtain a cell-free extract. A supernatant is obtained by centrifuging the cell-free extract, and then, a purified sample can be obtained from the supernatant by applying, singly or in combination, the following ordinary protein isolation and purification methods: the solvent extraction, the salting-out method using ammonium sulfate or the like, the desalting method, the precipitation method using an organic solvent, the anion exchange chromatography using resins such as diethylaminoethyl (DEAE) sepharose, the cation exchange chromatography using resins such as S-Sepharose FF (manufactured by Pharmacia), the hydrophobic chromatography using resins such as butyl sepharose or phenyl sepharose, the gel filtration method using a molecular sieve, the affinity chromatography, the chromatofocusing method, and the electrophoresis such as isoelectric focusing.

(5) Use of the Fluorescent Protein of the Present Invention and a Fusion Fluorescent Protein Comprising the Same

The fluorescent protein of the present invention can be fused with another protein, so as to construct a fusion fluorescent protein.

A method of obtaining the fusion fluorescent protein of the present invention is not particularly limited. It may be either a protein synthesized by chemosynthesis, or recombinant protein produced by a gene recombination technique.

Where a recombinant protein is produced, it is necessary to obtain DNA encoding the protein. Appropriate primers are designed using the information regarding the amino acid sequence shown in SEQ ID NO: 1 of the sequence listing of the present specification and the nucleotide sequence shown in SEQ ID NO: 2 thereof. Using these primers, PCR is carried out using a DNA fragment containing the gene of the fluorescent protein of the present invention as a template, so as to produce DNA fragments necessary for construction of the DNA encoding the fluorescent protein of the present invention. Moreover, DNA fragments encoding a protein to be fused is also obtained in the same above manner.

Subsequently, the thus obtained DNA fragments are ligated to one another by a gene recombination technique, so that DNA encoding the desired fusion fluorescent protein can be obtained. This DNA is then introduced into an appropriate expression system, so that the fusion fluorescent protein of the present invention can be produced.

The fluorescent protein of the present invention has an extremely high utility value as a marker. This is to say, the fluorescent protein of the present invention is purified as a fusion protein with an amino acid sequence to be tested, and the fusion protein is introduced into cells by methods such as the microinjection. By observing the distribution of the fusion protein over time, targeting activity of the amino acid sequence to be tested can be detected in the cells.

The type of another protein (an amino acid sequence to be tested) with which the fluorescent protein of the present invention is fused is not particularly limited. Preferred examples may include proteins localizing in cells, proteins specific for intracellular organelles, and targeting signals (e.g., a nuclear transport signal, a mitochondrial presequence, etc.). In addition, the fluorescent protein of the present invention can be expressed in cells and used, as well as being introduced into cells by the microinjection or the like. In this case, a vector into which the DNA encoding the fluorescent protein of the present invention is inserted in such a way that it can be expressed, is introduced into host cells.

Moreover, the fluorescent protein of the present invention can also be used as a reporter protein to determine promoter activity. This is to say, a vector is constructed such that DNA

encoding the fluorescent protein of the present invention is located downstream of a promoter to be tested, and the vector is then introduced into host cells. By detecting the fluorescence of the fluorescent protein of the present invention which is emitted from the cells, the activity of the promoter to be tested can be determined. The type of a promoter to be tested is not particularly limited, as long as it operates in host cells.

A vector used to detect the targeting activity of the above amino acid sequence to be tested or to determine promoter activity is not particularly limited. Examples of a vector preferably used for animal cells may include pNEO (P. Southern, and P. Berg (1982) *J. Mol. Appl. Genet.* 1: 327), pCAGGS (H. Niwa, K. Yamamura, and J. Miyazaki, *Gene* 108, 193-200 (1991)), pRc/CMV (manufactured by Invitrogen), and pCDM8 (manufactured by Invitrogen). Examples of a vector preferably used for yeasts may include pRS303, pRS304, pRS305, pRS306, pRS313, pRS314, pRS315, pRS316 (R. S. Sikorski and P. Hieter (1989) *Genetics* 122: 19-27), pRS423, pRS424, pRS425, pRS426 (T. W. Christianson, R. S. Sikorski, M. Dante, J. H. Shero, and P. Hieter (1992) *Gene* 110: 119-122).

In addition, the type of cells used herein is also not particularly limited. Various types of animal cells such as L cells, BalbC-3T3 cells, NIH3T3 cells, CHO (Chinese hamster ovary) cells, HeLa cells or NRK (normal rat kidney) cells, yeast cells such as *Saccharomyces cerevisiae*, *Escherichia coli* cells, or the like can be used. Vector can be introduced into host cells by common methods such as the calcium phosphate method or the electroporation.

The above obtained fusion fluorescent protein of the present invention wherein the fluorescent protein of the present invention is fused with another protein (referred to as a protein X) is allowed to be expressed in cells. By monitoring a fluorescence emitted, it becomes possible to analyze the localization or dynamics of the protein X in cells. That is, cells transformed or transfected with DNA encoding the fusion fluorescent protein of the present invention are observed with a fluorescence microscope, so that the localization and dynamics of the protein X in the cells can be visualized and thus analyzed.

For example, by using a protein specific for an intracellular organella as a protein X, the distribution and movement of a nucleus, a mitochondria, an endoplasmic reticulum, a Golgi body, a secretory vesicle, a peroxisome, etc., can be observed.

Moreover, for example, axis cylinders or dendrites of the nerve cells show an extremely complicated change in strikes in an individual who is under development. Accordingly, fluorescent labeling of these sites enables a dynamic analysis.

The fluorescence of the fluorescent protein of the present invention can be detected with a viable cell. Such detection can be carried out using, for example, a fluorescence microscope (Axiophoto Filter Set 09 manufactured by Carl Zeiss) or an image analyzer (Digital Image Analyzer manufactured by ATTO).

The type of a microscope can be appropriately selected depending on purposes. Where frequent observation such as pursuit of a change over time is carried out, an ordinary incident-light fluorescence microscope is preferable. Where observation is carried out while resolution is emphasized, for example, in the case of searching localization in cells specifically, a confocal laser scanning microscope is preferable. In terms of maintenance of the physiological state of cells and prevention from contamination, an inverted microscope is preferable as a microscope system. When an erecting microscope with a high-powered lens is used, a water immersion lens can be used.

A filter set can be appropriately selected depending on the fluorescence wavelength of a fluorescent protein. In the case of the fluorescent protein of the present invention, a filter having an excitation light between approximately 490 and 510 nm and a fluorescence between approximately 510 and 530 nm can be preferably used.

When viable cells are observed over time using a fluorescence microscope, a high sensitive cooled CCD camera is used, since photography is carried out in a short time. In the case of the cooled CCD camera, CCD is cooled to decrease thermal noise, so that a weak fluorescence image can be clearly photographed by exposure in a short time.

(6) Kit of the Present Invention

The present invention provides a kit for analyzing the localization of intracellular components and/or analyzing physiologically active substances, which is characterized in that it comprises at least one selected from the fluorescent protein, the fusion fluorescent protein, the DNA, the recombinant vector, or the transformant, which are described in the present specification. The kit of the present invention can be produced from commonly used materials that are known per se, by using common methods.

Reagents such as the fluorescent protein or the DNA are dissolved in an appropriate solvent, so that the reagents can be prepared in a form suitable for conservation. Water, ethanol, various types of buffer solution, etc. can be used as such a solvent.

The present invention will be further described in the following examples. However, the present invention is not limited by these examples.

EXAMPLES

Example 1

Isolation of Novel Fluorescent Protein Gene from Coral (Faviidae)

(1) Extraction of Total RNA

The fluorescent protein gene was isolated from fluorescing corals. *Favia favaus* was employed as a material. Faviidae was crushed with a hammer, 15 ml of TRIzol (GIBCO BRL) was added to 11 g (a wet weight) of the crushed Faviidae, and the resultant was agitated, followed by centrifugation at 1,500×g for 10 minutes. Chloroform (3 ml) was added to the supernatant, and the mixture was agitated for 15 seconds, allowed to stand for 3 minutes, and then centrifuged at 7,500×g for 15 minutes. Isopropanol (3.75 ml) was added to the supernatant, and the mixture was agitated for 15 seconds, allowed to stand for 10 minutes, and then centrifuged at 17,000×g for 10 minutes. The supernatant was discarded, 6 ml of 70% ethanol was added, and the mixture was centrifuged at 17,000×g for 10 minutes. The supernatant was discarded, and the precipitate was dissolved in 200 µl of DEPC water. The total RNA dissolved in DEPC water was diluted 100-fold, and OD260 and OD280 thereof were measured to determine the RNA concentration. 20 µg of total RNA was obtained.

(2) Synthesis of First Strand cDNA

cDNA (33 µl) was synthesized from 3 µg of total RNA using a First-strand cDNA synthesis kit "Ready-To-Go" (Amersham Pharmacia).

(3) Degenerated PCR

From 33 µl of the synthesized first strand cDNA, 3 µl thereof was separated, and PCR was carried out using the same as a template. The amino acid sequences of existing fluorescent proteins were analyzed, similar portions were

extracted, the extracted sequences were converted into nucleotide sequences, and primers were designed based thereon. Primers used:

(SEQ ID NO: 3)
5'-GGI WSB GTI AAY GGV CAY DAN TT -3'
(Primer 1)

(SEQ ID NO: 4)
5'-AACTGGAAGAATTGCGGCCGAGGAA -3'
(Primer 2)

R = A or G, Y = C or T, V = A, C, or G, D = A, G or T

Composition of PCR solution

Template (first strand cDNA)	3 μ l
\times 10 taq buffer	5 μ l
2.5 mM dNTPs	4 μ l
100 μ M primer 1	1 μ l
100 μ M primer 2	1 μ l
MilliQ	35 μ l
Taq polymerase (5 U/ μ l)	1 μ l

PCR conditions

94° C. for 1 min (PAD)
94° C. for 30 sec (denaturation)
52° C. for 30 sec (annealing of primer to template)
72° C. for 1 min (primer elongation)
72° C. for 7 min (final elongation)
4° C. (preservation)

The amplification product (1 μ l) obtained in the first PCR cycle was used as a template to carry out PCR under the same temperature conditions, provided that the following primers were used:

(SEQ ID NO: 5)
5'-TGC CWT TTG CIT TIG AYA TIT TG -3'
(Primer 3)

(SEQ ID NO: 6)
5'-GTC ITC TTY TGC ACI ACI GGI CCA TYD GVA GGA
AA -3'
(Primer 4)

A band of an expected size of 350 bp was cleaved via agarose gel electrophoresis and then purified.

(4) Subcloning and Nucleotide Sequencing

The purified DNA fragment was ligated to the pT7-Blue vector (Novagen). The resultant was transformed into the *E. coli* strain (TG1) for blue/white selection, plasmid DNA was purified from *E. coli* white colonies, and the nucleotide sequence of the inserted DNA fragment was determined using a DNA sequencer. The determined nucleotide sequences were compared with the nucleotide sequences of other fluorescent protein genes to determine whether or not the nucleotide sequence of the DNA of interest was derived from a fluorescent protein. As to the nucleotide sequences that had been determined to be a part of a fluorescent protein gene, cloning of a full-length gene was carried out by the 5'-RACE method and by the 3'-RACE method.

(5) 5'-RACE Method

In order to determine the nucleotide sequence on the 5'-side of the DNA fragment obtained via degenerated PCR, the 5'-RACE method was implemented using the 5'-RACE System for Rapid Amplification of cDNA Ends, Version 2.0 (GIBCO BRL). The total RNA (3 μ g) prepared in (1) was used as a template.

The following primers were used for the first amplification of DC-tailed cDNA.

(SEQ ID NO: 7)
5'-GGCCACGCGTCGACTAGTACGGGIIGGGIIGGGIIG-3'
(Primer 5)

(SEQ ID NO: 8)
5'-TTG TCA AGA TAT CGA AAG CGA ACG GCA GAG -3'
(Primer 6)
I = inosine

The following primers were used for the second amplification.

(SEQ ID NO: 9)
5'-GGCCACGCGTCGACTAGTAC-3'

(SEQ ID NO: 10)
5'-GTC CAC CCT CTA CGA CTT TGA GTT CCA TAT-3'

PCR conditions were set in accordance with the protocol included in the kit.

The amplified 700-bp band was cleaved via agarose gel electrophoresis and then purified. The purified DNA fragment was ligated to the pT7-Blue vector (Novagen). The resultant was transformed into the *E. coli* strain (TG1) for blue/white selection, plasmid DNA was purified from *E. coli* white colonies, and the nucleotide sequence of the inserted DNA fragment was determined using a DNA sequencer.

(6) Determination of Full Length Nucleotide Sequences and Protein Expression in *E. coli*

A primer was prepared from a portion corresponding to the N-terminus of the protein obtained in (5), an oligo dT primer was used for the C-terminus, and PCR was carried out using the first strand cDNA prepared in (2) as a template. Primer used:

(SEQ ID NO: 11)
5'-CCC GGA TCC GAT GAG TGT GAT TAC AWC AGA AAT GAA
GAT GGA GC-3'
(Primer 7)

Composition of PCR solution

Template (first strand cDNA)	3 μ l
\times 10 pyrobest buffer	5 μ l
2.5 mM dNTPs	4 μ l
100 μ M primer 7	1 μ l
100 μ M oligo dT primer	1 μ l
MilliQ	35 μ l
Pyrobest polymerase (5 U/ μ l)	1 μ l

PCR conditions

94° C. for 1 min (PAD)
94° C. for 30 sec (denaturation)
52° C. for 30 sec (annealing of primer to template)
72° C. for 1 min (primer elongation)
The above 3 steps were repeated 30 times.
72° C. for 7 min (final elongation)
4° C. (preservation)

The amplified band of approximately 900 bp was cleaved via agarose gel electrophoresis, purified, subcloned into the BamHI/EcoRI site of the pRSET vector (Invitrogen), and then expressed in the *E. coli* strain (JM109-DE3). The plasmid was recovered, and the inserted full length nucleotide sequence was determined. The clone was designated as KkG.

The full-length nucleotide sequence is shown in SEQ ID NO: 2, and the full-length amino acid sequence is shown in SEQ ID NO: 1.

Since the expressed protein was constructed to have a His-tag at its N-terminus, the expressed protein was purified on Ni-Agarose gel (QIAGEN). Purification was carried out in accordance with the protocol included in the kit. Subsequently, the properties of the purified protein were analyzed. (7) Analysis of Fluorescence Properties

The absorption spectrum was measured using a solution of 10 μ M fluorescent protein (KkG) in PBS. The molar absorption coefficient was determined based on the peak value of this spectrum. The absorption peak was observed at 507 nm, the fluorescent protein was diluted with the aforementioned buffer to adjust the absorption at 450 nm to 0.005, and the fluorescence emission spectra after excitation at 450 nm were measured (FIG. 1). Similarly, the fluorescence emission spectrum of EGFP (CLONTECH) was measured in a state where the absorption at 450 nm is 0.005. The quantum yield of EGFP was set at 0.6, and the quantum yield of the protein of the present invention was determined. The results are shown in Table 1.

TABLE 1

	Excitation maximum	Fluorescence maximum	Molar absorption coefficient	Quantum yield	pH sensitivity	Amino acid number
KkG	507 nm	517 nm	80,000 (482 nm)	0.68	Stable at pH 5 to 11	227

(8) Measurement of pH Sensitivity

The fluorescent protein of the present invention was diluted with the following buffers, and the fluorescence emission spectra were measured.

The pH levels of the buffers are as follows.

pH4, 5: Acetate buffer

pH6: MES buffer

pH7: MOPS buffer

pH8: HEPES buffer

pH9, 10: Glycine buffer

pH11: Phosphate buffer

The results of pH dependence measured at the fluorescence maximum wavelength are shown in FIG. 2.

Example 2

Preparation of Various Fluorescent Proteins with Improved Fluorescence Properties

(1) Preparation of Fluorescent Protein Capable of Green-to-red Photoconversion in Response to Irradiation with Light (Ultraviolet Rays and Purple Light)

By substitution from aspartic acid (D) to histidine (H) at 62th amino acid residue of the fluorescent protein KkG that is incapable of green-to-red photoconversion in response to irradiation with light, the properties of this protein could be modified so that it became capable of green-to-red photoconversion in response to irradiation with light (where the resulting protein is referred to as "KKH," the amino acid sequence of which is shown in SEQ ID NO: 12, and the nucleotide sequence of which is shown in SEQ ID NO: 13) (FIGS. 3 and 4). In FIG. 4A2, an arrow indicates an increase in the absorption (583 nm) at a portion that emits red fluorescence upon irradiation with light. By substitution from methionine (M) to valine (V) at 40th amino acid residue, substitution from aspar-

tic acid (D) to histidine (H) at 62th amino acid residue, and substitution from isoleucine (I) to methionine (M) at 198th amino acid residue in the fluorescent protein KkG, a fluorescent protein having photosensitivity higher than that of the fluorescent protein (KKH) that is capable of green-to-red photoconversion in response to irradiation with light, i.e., a fluorescent protein capable of green-to-red photoconversion with a weak light (H8PV, the amino acid sequence of which is shown in SEQ ID NO: 14, and the nucleotide sequence of which is shown in SEQ ID NO: 15), could be obtained (FIG. 3).

By substitution from methionine (M) to isoleucine (I) at 10th amino acid residue, substitution from leucine (L) to valine (V) at 12th amino acid residue, substitution from methionine (M) to valine (V) at 40th amino acid residue, substitution from valine (V) to alanine (A) at 60th amino acid residue, substitution from aspartic acid (D) to histidine (H) at 62th amino acid residue, substitution from tyrosine (Y) to asparagine (N) at 119th amino acid residue, substitution from proline (P) to serine (S) at 144th amino acid residue, substitution from arginine (R) to leucine (L) at 197th amino acid residue, and substitution from isoleucine (I) to methionine (M) at 198th amino acid residue in the fluorescent protein KkG, a fluorescent protein having photosensitivity higher than that of the fluorescent protein (H8PV) capable of green-to-red photoconversion in response to irradiation with light, i.e., a fluorescent protein capable of green-to-red photoconversion with a weak light (H38PVL, the amino acid sequence of which is shown in SEQ ID NO: 16, and the nucleotide sequence of which is shown in SEQ ID NO: 17), could be obtained (FIG. 3).

By substitution from methionine (M) to isoleucine (I) at 10th amino acid residue, substitution from methionine (M) to valine (V) at 40th amino acid residue, substitution from valine (V) to alanine (A) at 60th amino acid residue, substitution from aspartic acid (D) to histidine (H) at 62th amino acid residue, substitution from lysine (K) to glutamic acid (E) at 70th amino acid residue, substitution from tyrosine (Y) to asparagine (N) at 119th amino acid residue, substitution from arginine (R) to glutamine (Q) at 197th amino acid residue, and substitution from isoleucine (I) to methionine (M) at 198th amino acid residue in the fluorescent protein KkG, a fluorescent protein having photosensitivity higher than that of the fluorescent protein (H38PVL) capable of green-to-red photoconversion in response to irradiation with light, i.e., a fluorescent protein capable of green-to-red photoconversion with a weak light (KikGR, the amino acid sequence of which is shown in SEQ ID NO: 18, and the nucleotide sequence of which is shown in SEQ ID NO: 19), could be obtained (FIGS. 3 and 4).

FIG. 4A3 and FIG. 4A4 show that absorption (507 nm) at a portion emitting green fluorescence (517 nm) is gradually decreased upon irradiation with light and absorption (583 nm) at a portion emitting red fluorescence (593 nm) is gradually increased. That is, a fluorescent protein capable of green-to-red photoconversion in response to irradiation with light can be produced by introducing substitution from an amino acid corresponding to 62th amino acid residue of KkG (more specifically, this amino acid refers to X of 3 amino acid residues XYG which forms the chromophore shown in an asterisk portion of FIG. 3, wherein X represents any amino acid, Y generally represents tyrosine and optionally an aromatic amino acid such as phenylalanine, tryptophan, or histidine, and G represents glycine) to histidine in a fluorescent protein KkG incapable of green-to-red photoconversion in response to irradiation with light. Further, when any fluorescent protein incapable of green-to-red photoconversion in

response to irradiation with light which has substitution from an amino acid corresponding to 62th amino acid residue of KkG to histidine (H) further comprises any of the following amino acid substitutions; namely, substitution from an amino acid corresponding to 10th amino acid residue of KkG to isoleucine (I), substitution from an amino acid corresponding to 12th amino acid residue of KkG to valine (V), substitution from an amino acid corresponding to 40th amino acid residue of KkG to valine (V), substitution from an amino acid corresponding to 60th amino acid residue of KkG to alanine (A), substitution from an amino acid corresponding to 70th amino acid residue of KkG to glutamic acid (E), substitution from an amino acid corresponding to 119th amino acid residue of KkG to asparagine (N), substitution from an amino acid corresponding to 144 amino acid residue of KkG to serine (S), substitution from an amino acid corresponding to 197th amino acid residue of KkG to leucine (L) or glutamine (Q), or substitution from an amino acid corresponding to 198th amino acid residue of KkG to methionine (M); a fluorescent protein capable of green-to-red photoconversion in response to irradiation with light can be prepared, and a fluorescent protein with high photosensitivity, i.e., a fluorescent protein capable of green-to-red photoconversion with a weak light, can be obtained.

The fluorescent proteins Kaede and KikGR capable of green-to-red photoconversion in response to irradiation with light were expressed in *E. coli* and then compared. As a result, the fluorescence intensity of KikGR was found to be higher than that of Kaede both before and after photoconversion (FIG. 5A). When the Kaede and KikGR genes were introduced into the HeLa cells and expressed therein, KikGR emitted fluorescence earlier than Kaede (FIG. 5B). When KikGR and Kaede that had been expressed in the HeLa cells were irradiated with light to result in green-to-red photoconversion in the cells, the rate of green-to-red photoconversion of KikGR was apparently faster than that of Kaede, and the fluorescence intensity of KikGR was stronger (FIGS. 6A, 6B, 6C, and 6D). Accordingly, it is easily expected that the rate of green-to-red photoconversion can be increased and the fluorescence intensity can be enhanced by introducing the same amino acid substitution as that of the case of KikGR at a portion of Kaede which was found to be different from that of KikGR upon comparison of amino acid sequences. In particular, the shaded portion in FIG. 3 is considered to significantly affect fluorescence properties, since the amino acid side chain is directed inside the protein when the protein has a three-dimensional structure. Specifically, when any fluorescent protein having substitution from an amino acid corresponding to 62th amino acid residue of KkG to histidine (H) comprises any of the following amino acid substitutions; namely substitution from an amino acid corresponding to 54th amino acid residue of KkG to phenylalanine (F), substitution from an amino acid corresponding to 69th amino acid residue of KkG to valine (V), substitution from an amino acid corresponding to 87th amino acid residue of KkG to tyrosine

(Y), substitution from an amino acid corresponding to 93th amino acid residue of KkG to methionine (M), substitution from an amino acid corresponding to 109th amino acid residue of KkG to methionine (M), substitution from an amino acid corresponding to 121th amino acid residue of KkG to isoleucine (I), substitution from an amino acid corresponding to 140th amino acid residue of KkG to valine (V), or substitution from an amino acid corresponding to 160th amino acid residue of KkG to valine (V); a fluorescent protein with an increased rate of green-to-red photoconversion and enhanced fluorescence intensity can be produced.

(2) Preparation of Fluorescent Protein Capable of Purple-to-blue Photoconversion in Response to Irradiation with Light (Ultraviolet Rays and Purple Light)

The fluorescent protein KkG incapable of purple-to-blue photoconversion in response to irradiation with light was subjected to substitution from valine (V) to alanine (A) at 60th amino acid residue, substitution from aspartic acid (D) to glycine (G) at 62th amino acid residue, substitution from tyrosine (Y) to histidine (H) at 63th amino acid residue, substitution from histidine (H) to leucine (L) at 197th amino acid residue, and substitution from isoleucine (I) to threonine (T) at 199th amino acid residue. Thus, the properties of this protein could be modified so that it became capable of purple (380 nm)-to-blue (450 nm) photoconversion in response to irradiation with light (where the resulting protein is referred to as "Kbl2," the amino acid sequence of which is shown in SEQ ID NO: 20, and the nucleotide sequence of which is shown in SEQ ID NO: 21) (FIGS. 3, 7, 8 and 9). Specifically, a fluorescent protein capable of purple-to-blue photoconversion in response to irradiation with light can be prepared by introducing all or any one of the following substitutions in any fluorescent protein; substitution from an amino acid corresponding to 60th amino acid residue of KkG to alanine (A), substitution from an amino acid corresponding to 62th amino acid residue of KkG to glycine (G), substitution from an amino acid corresponding to 63th amino acid residue of KkG to histidine (H), substitution from an amino acid corresponding to 196th amino acid residue of KkG to leucine (L), and substitution from an amino acid corresponding to 198th amino acid residue of KkG to threonine (T).

Industrial Applicability

The present invention provides a novel fluorescent protein derived from *favia favaus*. The fluorescent protein of the present invention is a novel protein having primary structures that differ from those of existing fluorescent proteins. The fluorescent protein of the present invention has given fluorescence properties and thus is useful in molecular biological analysis. Therefore, use of the fluorescent protein of the present invention enables fluorescence labeling of mammalian cells without causing toxicity. Use of a novel gene that is discovered through the present invention as a starting material enables provision of fluorescent substances exhibiting a wide variety of properties.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 23

<210> SEQ ID NO 1

<211> LENGTH: 227

<212> TYPE: PRT

<213> ORGANISM: *Favia favaus*

-continued

<400> SEQUENCE: 1

Met Ser Val Ile Thr Ser Glu Met Lys Met Glu Leu Arg Met Glu Gly
 1 5 10 15
 Ala Val Asn Gly His Lys Phe Val Ile Thr Gly Lys Gly Ser Gly Gln
 20 25 30
 Pro Phe Glu Gly Ile Gln Asn Met Asp Leu Thr Val Ile Glu Gly Gly
 35 40 45
 Pro Leu Pro Phe Ala Phe Asp Ile Leu Thr Thr Val Phe Asp Tyr Gly
 50 55 60
 Asn Arg Val Phe Val Lys Tyr Pro Glu Glu Ile Val Asp Tyr Phe Lys
 65 70 75 80
 Gln Ser Phe Pro Glu Gly Tyr Ser Trp Glu Arg Ser Met Ser Tyr Glu
 85 90 95
 Asp Gly Gly Ile Cys Leu Ala Thr Asn Asn Ile Thr Met Lys Lys Asp
 100 105 110
 Gly Ser Asn Cys Phe Val Tyr Glu Ile Arg Phe Asp Gly Val Asn Phe
 115 120 125
 Pro Ala Asn Gly Pro Val Met Gln Arg Lys Thr Val Lys Trp Glu Pro
 130 135 140
 Ser Thr Glu Lys Met Tyr Val Arg Asp Gly Val Leu Lys Gly Asp Val
 145 150 155 160
 Asn Met Ala Leu Leu Leu Gln Gly Gly Gly His Tyr Arg Cys Asp Phe
 165 170 175
 Arg Thr Thr Tyr Lys Ala Lys Lys Val Val Gln Leu Pro Asp Tyr His
 180 185 190
 Phe Val Asp His Arg Ile Glu Ile Thr Ser His Asp Lys Asp Tyr Asn
 195 200 205
 Lys Val Lys Leu Tyr Glu His Ala Lys Ala His Ser Gly Leu Pro Arg
 210 215 220
 Leu Ala Lys
 225

<210> SEQ ID NO 2

<211> LENGTH: 684

<212> TYPE: DNA

<213> ORGANISM: Favia favaus

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (1)..(681)

<400> SEQUENCE: 2

atg agt gtg att aca tca gaa atg aag atg gag ctg cgt atg gaa ggc 48
 Met Ser Val Ile Thr Ser Glu Met Lys Met Glu Leu Arg Met Glu Gly
 1 5 10 15
 gct gta aac ggg cac aag ttc gtg att aca ggg aaa gga agt ggc cag 96
 Ala Val Asn Gly His Lys Phe Val Ile Thr Gly Lys Gly Ser Gly Gln
 20 25 30
 cct ttc gag gga ata cag aat atg gac ctg aca gtc ata gag ggc gga 144
 Pro Phe Glu Gly Ile Gln Asn Met Asp Leu Thr Val Ile Glu Gly Gly
 35 40 45
 cct ctt cct ttt gct ttc gat atc ctg aca aca gta ttc gat tac ggc 192
 Pro Leu Pro Phe Ala Phe Asp Ile Leu Thr Thr Val Phe Asp Tyr Gly
 50 55 60
 aac cgg gta ttt gtc aaa tac cca gaa gaa ata gta gac tac ttc aag 240
 Asn Arg Val Phe Val Lys Tyr Pro Glu Glu Ile Val Asp Tyr Phe Lys
 65 70 75 80
 cag tcg ttt cct gag ggt tat tct tgg gaa cga agc atg agt tac gaa 288
 Gln Ser Phe Pro Glu Gly Tyr Ser Trp Glu Arg Ser Met Ser Tyr Glu

-continued

85	90	95	
gac ggg gga att tgc ctc gcc	aca aac aat ata acg atg aag aaa gac		336
Asp Gly Gly Ile Cys Leu Ala Thr	Asn Asn Ile Thr Met Lys Lys Asp		
100	105	110	
ggc agc aac tgt ttt gtc tat	gaa att cga ttt gat ggt gtg aac ttt		384
Gly Ser Asn Cys Phe Val Tyr	Glu Ile Arg Phe Asp Gly Val Asn Phe		
115	120	125	
cct gcc aat ggt cca gtt atg	cag agg aag acc gtc aaa tgg gag cca		432
Pro Ala Asn Gly Pro Val Met	Gln Arg Lys Thr Val Lys Trp Glu Pro		
130	135	140	
tcc act gag aaa atg tat	gtg cgt gat gga gtg ctg aag ggt gat gtt		480
Ser Thr Glu Lys Met Tyr Val	Arg Asp Gly Val Leu Lys Gly Asp Val		
145	150	155	160
aac atg gct ctg ttg ctt caa	gga ggt ggc cat tac cga tgt gac ttc		528
Asn Met Ala Leu Leu Leu Gln	Gly Gly Gly His Tyr Arg Cys Asp Phe		
165	170	175	
aga act act tac aaa gca aag	aag gtt gtc cag ttg cca gac tat cac		576
Arg Thr Thr Tyr Lys Ala Lys	Lys Val Val Gln Leu Pro Asp Tyr His		
180	185	190	
ttc gtg gat cat cga att	gag ata aca agc cat gac aag gat tac aac		624
Phe Val Asp His Arg Ile Glu	Ile Thr Ser His Asp Lys Asp Tyr Asn		
195	200	205	
aag gtt aag ctg tat gag	cat gct aaa gct cat tcc ggg ctg cca agg		672
Lys Val Lys Leu Tyr Glu His	Ala Lys Ala His Ser Gly Leu Pro Arg		
210	215	220	
ctg gcc aag taa			684
Leu Ala Lys			
225			

<210> SEQ ID NO 3
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 primer
 <220> FEATURE:
 <221> NAME/KEY: modified_base
 <222> LOCATION: (3)..(3)
 <223> OTHER INFORMATION: Inosine
 <220> FEATURE:
 <221> NAME/KEY: modified_base
 <222> LOCATION: (9)..(9)
 <223> OTHER INFORMATION: Inosine
 <220> FEATURE:
 <221> NAME/KEY: modified_base
 <222> LOCATION: (21)..(21)
 <223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 3

ggnwsbgtna ayggvcayda ntt 23

<210> SEQ ID NO 4
 <211> LENGTH: 27
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 primer

<400> SEQUENCE: 4

aactggaaga attcgcggcc gcaggaa 27

<210> SEQ ID NO 5
 <211> LENGTH: 23
 <212> TYPE: DNA

-continued

```

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
    primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: Inosine
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (14)..(14)
<223> OTHER INFORMATION: Inosine
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (20)..(20)
<223> OTHER INFORMATION: Inosine

```

```

<400> SEQUENCE: 5

```

```

tgccwtttgc nttngayatn ttg

```

23

```

<210> SEQ ID NO 6
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
    primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Inosine
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (15)..(15)
<223> OTHER INFORMATION: Inosine
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (18)..(18)
<223> OTHER INFORMATION: Inosine
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (21)..(21)
<223> OTHER INFORMATION: Inosine

```

```

<400> SEQUENCE: 6

```

```

gtcntcttyt gcacnacngg nccatydgva ggaaa

```

35

```

<210> SEQ ID NO 7
<211> LENGTH: 36
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
    primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (24)..(25)
<223> OTHER INFORMATION: Inosine
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (29)..(30)
<223> OTHER INFORMATION: Inosine
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (34)..(35)
<223> OTHER INFORMATION: Inosine

```

```

<400> SEQUENCE: 7

```

```

ggccacgcgt cgactagtagc gggnnngggnn gggnnng

```

36

```

<210> SEQ ID NO 8
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

```

-continued

<220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 8

ttgtcaagat atcgaaagcg aacggcagag 30

<210> SEQ ID NO 9
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 9

ggccacgcgt cgactagtag 20

<210> SEQ ID NO 10
 <211> LENGTH: 30
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 10

gtccaccctc tacgactttg agttccatat 30

<210> SEQ ID NO 11
 <211> LENGTH: 44
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 11

cccgatccg atgagtgatga ttacawcaga aatgaagatg gagc 44

<210> SEQ ID NO 12
 <211> LENGTH: 227
 <212> TYPE: PRT
 <213> ORGANISM: Favia fava

<400> SEQUENCE: 12

Met Ser Val Ile Thr Ser Glu Met Lys Met Glu Leu Arg Met Glu Gly
 1 5 10 15

Ala Val Asn Gly His Lys Phe Val Ile Thr Gly Lys Gly Ser Gly Gln
 20 25 30

Pro Phe Glu Gly Ile Gln Asn Met Asp Leu Thr Val Ile Glu Gly Gly
 35 40 45

Pro Leu Pro Phe Ala Phe Asp Ile Leu Thr Thr Val Phe His Tyr Gly
 50 55 60

Asn Arg Val Phe Val Lys Tyr Pro Glu Glu Ile Val Asp Tyr Phe Lys
 65 70 75 80

Gln Ser Phe Pro Glu Gly Tyr Ser Trp Glu Arg Ser Met Ser Tyr Glu
 85 90 95

Asp Gly Gly Ile Cys Leu Ala Thr Asn Asn Ile Thr Met Lys Lys Asp
 100 105 110

Gly Ser Asn Cys Phe Val Tyr Glu Ile Arg Phe Asp Gly Val Asn Phe
 115 120 125

Pro Ala Asn Gly Pro Val Met Gln Arg Lys Thr Val Lys Trp Glu Pro

-continued-

```

130      135      140
Ser Thr Gln Lys Met Tyr Val Arg Asp Gly Val Leu Lys Gly Asp Val
145      150      155      160
Asn Met Ala Leu Leu Leu Gln Gly Gly His Tyr Arg Cys Asp Phe
165      170      175      180
Arg Thr Thr Tyr Lys Ala Lys Lys Val Val Gln Leu Pro Asp Tyr His
185      190      195      200
Phe Val Asp His Arg Ile Gln Ile Thr Ser His Asp Lys Asp Tyr Asn
205      210      215      220
Lys Val Lys Leu Tyr Gln His Ala Lys Ala His Ser Gly Leu Pro Arg
225
Leu Ala Lys
<210> SEQ ID NO 13
<211> LENGTH: 684
<212> TYPE: DNA
<213> ORGANISM: Favia favaus
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(681)
<400> SEQUENCE: 13
48      atg agt gtg atg atc aca tca gaa atg aag atg gag ctg cgt atg gaa ggc
1      Met Ser Val Ile Thr Ser Gln Met Lys Met Gln Leu Arg Met Gln Gly
5      10      15
gct gta aac ggg cac aag ttc gtg atc aca ggg aaa gga agt ggc cag
96      Ala Val Asn Gly His Lys Phe Val Ile Thr Gly Lys Gly Ser Gly Gln
20      25
gag gga ata cag aat atg gac ctg aca gtc ata gag ggc gga
144      Pro Phe Gln Gly Ile Gln Asn Met Asp Leu Thr Val Ile Gln Gly Gly
35      40      45
cct ctt cct ctt gct ttc gat atc ctg aca gta ttc cat tac ggc
192      Pro Leu Pro Phe Ala Phe Asp Ile Leu Thr Val Phe His Tyr Gly
50      55      60
aac cgg gta ttc gtc aaa tac cca gaa gaa ata gta gac tac tcc aag
240      Asn Arg Val Phe Val Lys Tyr Pro Gln Ile Val Asp Tyr Phe Lys
65      70      75      80
cag tgc ttc cct gag ggt tat tct tgg gaa cga agc atg agt tac gaa
288      Gln Ser Phe Pro Gln Gly Tyr Ser Trp Gln Arg Ser Met Ser Tyr Gln
85      90      95
gac ggg gga att tgc ccc aca aac aat ata acg atg aag aaa gac
336      Asp Gly Gly Ile Cys Leu Ala Thr Asn Ile Thr Met Lys Asp
100      105      110
ggc agc aac tgt ttc gtc tat gaa atc cga ttt gat ggt atg aac ttt
384      Gly Ser Asn Cys Phe Val Tyr Gln Ile Arg Phe Asp Gly Val Asn Phe
115      120      125
cct gcc aat ggt cca gtt atg cag agg aag acc gtc aaa tgg gag cca
432      Pro Ala Asn Gly Pro Val Met Gln Arg Lys Thr Val Lys Trp Gln Pro
130      135      140
tcc act gag aaa atg tat gtc cgt gat gga gtc ctg aag ggt gat gtt
480      Ser Thr Gln Lys Met Tyr Val Arg Asp Gly Val Leu Lys Gly Asp Val
145      150      155      160
aac atg gct ctg ttc cta gga ggt ggc cat tac cga tgt gat ttc
528      Asn Met Ala Leu Leu Leu Gln Gly Gly His Tyr Arg Cys Asp Phe
165      170      175
aga act act tac aaa gca aag aag gtt gtc cag ctg cca gac tat cac
576      Arg Thr Thr Tyr Lys Ala Lys Lys Val Val Gln Leu Pro Asp Tyr His
180      185      190
ttc gtg gat cat cga att gag ata aca agc cat gac aag gat tac aac
624

```

-continued

```

Phe Val Asp His Arg Ile Glu Ile Thr Ser His Asp Lys Asp Tyr Asn
    195                200                205
aag gtt aag ctg tat gag cat gct aaa gct cat tcc ggg ctg cca agg    672
Lys Val Lys Leu Tyr Glu His Ala Lys Ala His Ser Gly Leu Pro Arg
    210                215                220

ctg gcc aag taa    684
Leu Ala Lys
225

```

```

<210> SEQ ID NO 14
<211> LENGTH: 227
<212> TYPE: PRT
<213> ORGANISM: Favia favaus

```

```

<400> SEQUENCE: 14

```

```

Met Ser Val Ile Thr Ser Glu Met Lys Met Glu Leu Arg Met Glu Gly
 1          5          10          15
Ala Val Asn Gly His Lys Phe Val Ile Thr Gly Lys Gly Ser Gly Gln
    20          25          30
Pro Phe Glu Gly Ile Gln Asn Val Asp Leu Thr Val Ile Glu Gly Gly
    35          40          45
Pro Leu Pro Phe Ala Phe Asp Ile Leu Thr Thr Val Phe His Tyr Gly
    50          55          60
Asn Arg Val Phe Val Lys Tyr Pro Glu Glu Ile Val Asp Tyr Phe Lys
 65          70          75          80
Gln Ser Phe Pro Glu Gly Tyr Ser Trp Glu Arg Ser Met Ser Tyr Glu
    85          90          95
Asp Gly Gly Ile Cys Leu Ala Thr Asn Asn Ile Thr Met Lys Lys Asp
   100          105          110
Gly Ser Asn Cys Phe Val Tyr Glu Ile Arg Phe Asp Gly Val Asn Phe
   115          120          125
Pro Ala Asn Gly Pro Val Met Gln Arg Lys Thr Val Lys Trp Glu Pro
   130          135          140
Ser Thr Glu Lys Met Tyr Val Arg Asp Gly Val Leu Lys Gly Asp Val
 145          150          155          160
Asn Met Ala Leu Leu Leu Gln Gly Gly Gly His Tyr Arg Cys Asp Phe
   165          170          175
Arg Thr Thr Tyr Lys Ala Lys Lys Val Val Gln Leu Pro Asp Tyr His
   180          185          190
Phe Val Asp His Arg Met Glu Ile Thr Ser His Asp Lys Asp Tyr Asn
   195                200                205
Lys Val Lys Leu Tyr Glu His Ala Lys Ala His Ser Gly Leu Pro Arg
   210                215                220

Leu Ala Lys
225

```

```

<210> SEQ ID NO 15
<211> LENGTH: 684
<212> TYPE: DNA
<213> ORGANISM: Favia favaus
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(681)

```

```

<400> SEQUENCE: 15

```

```

atg agt gtg att aca tca gaa atg aag atg gag ctg cgt atg gaa ggc    48
Met Ser Val Ile Thr Ser Glu Met Lys Met Glu Leu Arg Met Glu Gly
 1          5          10          15
gct gta aac ggg cac aag ttc gtg att aca ggg aaa gga agt ggc cag    96

```

-continued-

144	cct ttc gag gga ata cag gat ggc gac ctc gca gta gac ttc aag ggc gga Pro phe gln gly ile gln asn val asp leu thr val ile gln gly gly	35	40	45	192	cct ctt cct ttc gct ttc gat atc ctc gca aca gta ttc cat tac ggc Pro leu pro phe ala phe asp ile leu thr thr val phe his tyr gly	50	55	60	240	aac cgg gta ttt gtc aaa tac cca gaa gaa ata gta gac tac ttc aag Asn arg val phe val lys tyr pro gln ile val asp tyr phe lys	65	70	75	80	288	cag tgc ttt cct gag ggt tat tct tgg gaa cga agc atg agt tac gaa gln ser phe pro gln gly tyr ser trp gln arg ser met ser tyr gln	85	90	95	336	gac ggg gga att tgc ctc gcc aca aac ata cgy atg aag aaa gac asp gly gly ile cys leu ala thr asn ile thr met lys lys asp	100	105	110	384	ggc agc aac tgt ttc gtc tat gaa atc cga ttt gat ggt gtc aac ttt gly ser asn cys phe val tyr gln ile arg phe asp gly val asn phe	115	120	125	432	cct gcc aat ggt cca gtt atg cag agy aag acc gtc aaa tgg gag cca pro ala asn gly pro val met gln arg lys thr val lys trp gln pro	130	135	140	480	tcc act gag aaa atg tat gtc cgt gat gga gtc atg ctc aag ggt gat gtt ser thr gln lys met tyr val arg asp gly val leu lys gly asp val	145	150	155	528	aac atg gct ctc ctc ctc ctc cca gga ggt ggc cat tac cga tgt gac ttc asn met ala leu leu gln gly gly gly his tyr arg cys asp phe	165	170	175	576	aga act act tac aaa gca aag gtt gtc cag tgc cag tgc cca gac tat cac arg thr thr tyr lys ala lys lys val val gln leu pro asp tyr his	180	185	190	624	ttc gtc gat cat cga atg gaa gta aca agc cat gac aag gat tac aac phe val asp his arg met gln ile thr ser his asp lys asp tyr asn	195	200	205	672	aag gtc aag ctc gat gat gac cat gct aaa gct cat tcc ggg ctc cca agg lys val lys leu tyr gln his ala lys ala his ser gly leu pro arg	210	215	220	684	ctg gcc aag ta	225	225	<210> SEQ ID NO 16 <211> LENGTH: 227 <212> TYPE: PRT <213> ORGANISM: Favita favus <400> SEQUENCE: 16	1	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90	95
-----	--	----	----	----	-----	--	----	----	----	-----	--	----	----	----	----	-----	--	----	----	----	-----	--	-----	-----	-----	-----	--	-----	-----	-----	-----	--	-----	-----	-----	-----	--	-----	-----	-----	-----	--	-----	-----	-----	-----	--	-----	-----	-----	-----	--	-----	-----	-----	-----	--	-----	-----	-----	-----	----------------	-----	-----	--	---	---	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----

-continued

Asp Gly Gly Ile Cys Leu Ala Thr Asn Asn Ile Thr Met Lys Lys Asp
 100 105 110
 Gly Ser Asn Cys Phe Val Asn Glu Ile Arg Phe Asp Gly Val Asn Phe
 115 120 125
 Pro Ala Asn Gly Pro Val Met Gln Arg Lys Thr Val Lys Trp Glu Ser
 130 135 140
 Ser Thr Glu Lys Met Tyr Val Arg Asp Gly Val Leu Lys Gly Asp Val
 145 150 155 160
 Asn Met Ala Leu Leu Leu Gln Gly Gly Gly His Tyr Arg Cys Asp Phe
 165 170 175
 Arg Thr Thr Tyr Lys Ala Lys Lys Val Val Gln Leu Pro Asp Tyr His
 180 185 190
 Phe Val Asp His Leu Met Glu Ile Thr Ser His Asp Lys Asp Tyr Asn
 195 200 205
 Lys Val Lys Leu Tyr Glu His Ala Lys Ala His Ser Gly Leu Pro Arg
 210 215 220
 Leu Ala Lys
 225

<210> SEQ ID NO 17
 <211> LENGTH: 684
 <212> TYPE: DNA
 <213> ORGANISM: Favia favaus
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(681)

<400> SEQUENCE: 17

atg agt gtg att aca tca gaa atg aag atc gag gtg cgt atg gaa ggc	48
Met Ser Val Ile Thr Ser Glu Met Lys Ile Glu Val Arg Met Glu Gly	
1 5 10 15	
gct gta aac ggg cac aag ttc gtg att aca ggg aaa gga agt ggc cag	96
Ala Val Asn Gly His Lys Phe Val Ile Thr Gly Lys Gly Ser Gly Gln	
20 25 30	
cct ttc gag gga ata cag aat gtg gac ctg aca gtc ata gag ggc gga	144
Pro Phe Glu Gly Ile Gln Asn Val Asp Leu Thr Val Ile Glu Gly Gly	
35 40 45	
cct ctt cct ttt gct ttc gat atc ctg aca aca gca ttc cat tac ggc	192
Pro Leu Pro Phe Ala Phe Asp Ile Leu Thr Thr Ala Phe His Tyr Gly	
50 55 60	
aac cgg gta ttt gtc aaa tac cca gaa gaa ata gta gac tac ttc aag	240
Asn Arg Val Phe Val Lys Tyr Pro Glu Glu Ile Val Asp Tyr Phe Lys	
65 70 75 80	
cag tcg ttt cct gag ggt tat tct tgg gaa cga agc atg agt tac gaa	288
Gln Ser Phe Pro Glu Gly Tyr Ser Trp Glu Arg Ser Met Ser Tyr Glu	
85 90 95	
gac ggg gga att tgc ctc gcc aca aac aat ata acg atg aag aaa gac	336
Asp Gly Gly Ile Cys Leu Ala Thr Asn Asn Ile Thr Met Lys Lys Asp	
100 105 110	
ggc agc aac tgt ttt gtc aat gaa att cga ttt gat ggt gtg aac ttt	384
Gly Ser Asn Cys Phe Val Asn Glu Ile Arg Phe Asp Gly Val Asn Phe	
115 120 125	
cct gcc aat ggt cca gtt atg cag agg aag acc gtc aaa tgg gag tca	432
Pro Ala Asn Gly Pro Val Met Gln Arg Lys Thr Val Lys Trp Glu Ser	
130 135 140	
tcc act gag aaa atg tat gtg cgt gat gga gtg ctg aag ggt gat gtt	480
Ser Thr Glu Lys Met Tyr Val Arg Asp Gly Val Leu Lys Gly Asp Val	
145 150 155 160	
aac atg gct ctg ttg ctt caa gga ggt ggc cat tac cga tgt gac ttc	528

-continued

```

Asn Met Ala Leu Leu Leu Gln Gly Gly Gly His Tyr Arg Cys Asp Phe
      165                               170                               175
aga act act tac aaa gca aag aag gtt gtc cag ttg cca gac tat cac      576
Arg Thr Thr Tyr Lys Ala Lys Lys Val Val Gln Leu Pro Asp Tyr His
      180                               185                               190

ttc gtg gat cat cta atg gag ata aca agc cat gac aag gat tac aac      624
Phe Val Asp His Leu Met Glu Ile Thr Ser His Asp Lys Asp Tyr Asn
      195                               200                               205

aag gtt aag ctg tat gag cat gct aaa gct cat tcc ggg ctg cca agg      672
Lys Val Lys Leu Tyr Glu His Ala Lys Ala His Ser Gly Leu Pro Arg
      210                               215                               220

ctg gcc aag taa      684
Leu Ala Lys
225

```

```

<210> SEQ ID NO 18
<211> LENGTH: 227
<212> TYPE: PRT
<213> ORGANISM: Favia fавus

```

```

<400> SEQUENCE: 18

```

```

Met Ser Val Ile Thr Ser Glu Met Lys Ile Glu Leu Arg Met Glu Gly
1      5      10      15
Ala Val Asn Gly His Lys Phe Val Ile Thr Gly Lys Gly Ser Gly Gln
      20      25      30
Pro Phe Glu Gly Ile Gln Asn Val Asp Leu Thr Val Ile Glu Gly Gly
      35      40      45
Pro Leu Pro Phe Ala Phe Asp Ile Leu Thr Thr Ala Phe His Tyr Gly
      50      55      60
Asn Arg Val Phe Val Glu Tyr Pro Glu Glu Ile Val Asp Tyr Phe Lys
      65      70      75      80
Gln Ser Phe Pro Glu Gly Tyr Ser Trp Glu Arg Ser Met Ser Tyr Glu
      85      90      95
Asp Gly Gly Ile Cys Leu Ala Thr Asn Asn Ile Thr Met Lys Lys Asp
      100      105      110
Gly Ser Asn Cys Phe Val Asn Glu Ile Arg Phe Asp Gly Val Asn Phe
      115      120      125
Pro Ala Asn Gly Pro Val Met Gln Arg Lys Thr Val Lys Trp Glu Pro
      130      135      140
Ser Thr Glu Lys Met Tyr Val Arg Asp Gly Val Leu Lys Gly Asp Val
      145      150      155      160
Asn Met Ala Leu Leu Leu Gln Gly Gly Gly His Tyr Arg Cys Asp Phe
      165                               170                               175
Arg Thr Thr Tyr Lys Ala Lys Lys Val Val Gln Leu Pro Asp Tyr His
      180                               185                               190
Phe Val Asp His Gln Met Glu Ile Thr Ser His Asp Lys Asp Tyr Asn
      195                               200                               205
Lys Val Lys Leu Tyr Glu His Ala Lys Ala His Ser Gly Leu Pro Arg
      210                               215                               220
Leu Ala Lys
225

```

```

<210> SEQ ID NO 19
<211> LENGTH: 684
<212> TYPE: DNA
<213> ORGANISM: Favia fавus
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(681)

```

<400> SEQUENCE: 19

1 Met Ser Val Ile Thr Ser Gln Met Lys Ile Gln Leu Arg Met Gln Gly
48 gct gta aac ggg cac aag ttc gtg att aca ggg aaa gga agt ggc cag
96 Ala Val Asn Gly His Lys Phe Val Ile Thr Gly Lys Ser Gly Gln
20
30
144 cct ttc gag gga ata cag aat gtg gac ctg aca gtc ata gag ggc gga
35
40
45
192 cct cct cct ttt gct ttc gat atc ctg aca gca ttc cat tac ggc
50
55
60
240 aac cgg gta ttc gtc gaa tac cca gaa gaa ata gta gac tac ttc aag
65
70
75
288 cag ttc cct gag ggt tat tct tgg gaa cga agc atg agt tac gaa
85
90
336 gac ggg gga att tgc ctc gcc aca aac aat ata cag atg aag aaa gac
100
105
384 ggc agc aac tgt ttt gtc aat gaa att cga ttt gat ggt gtg aac ttt
115
120
125
432 cct gcc aat ggt cca gtc atg cag agg aag acc gtc aaa tgg gag cca
130
135
140
480 tcc act gag aaa atg tat gtc cgt gat gga gtc ctg aag ggt gat gra
145
150
155
480 Ser Thr Gln Lys Met Tyr Val Arg Asp Gly Val Leu Lys Gly Asp Val
160

528 aac atg gct ctg ttc caa gga ggt ggc cat tac cga tgt gac ttc
165
170
576 aga act act tac aaa gca aag aag gtc gtc cag ttc cca gac tac cac
180
185
624 ttc tgc gat cat caa atg gaa ata aca agc cat gac aag gat tac aac
195
200
672 aag gtt aag ctg tat gag cat gct aaa gct cat tcc ggg ctg cca agg
210
215
220
684 Leu Ala Lys
225

<210> SEQ ID NO 20

<211> LENGTH: 227

<212> TYPE: PRT

<213> ORGANISM: PAVIA FAVUS

<400> SEQUENCE: 20

1 Met Ser Val Ile Thr Ser Gln Met Lys Met Gln Leu Arg Met Gln Gly
5
10
15
Ala Val Asn Gly His Lys Phe Val Ile Thr Gly Lys Ser Gly Gln
20
25
30
Pro Phe Gln Gly Ile Gln Asn Met Asp Leu Thr Val Ile Gln Gly Gly
35
40
45

-continued

Pro Leu Pro Phe Ala Phe Asp Ile Leu Thr Thr Ala Phe Gly His Gly
 50 55 60

Asn Arg Val Phe Val Lys Tyr Pro Glu Glu Ile Val Asp Tyr Phe Lys
 65 70 75 80

Gln Ser Phe Pro Glu Gly Tyr Ser Trp Glu Arg Ser Met Ser Tyr Glu
 85 90 95

Asp Gly Gly Ile Cys Leu Ala Thr Asn Asn Ile Thr Met Lys Lys Asp
 100 105 110

Gly Ser Asn Cys Phe Val Tyr Glu Ile Arg Phe Asp Gly Val Asn Phe
 115 120 125

Pro Ala Asn Gly Pro Val Met Gln Arg Lys Thr Val Lys Trp Glu Pro
 130 135 140

Ser Thr Glu Lys Met Tyr Val Arg Asp Gly Val Leu Lys Gly Asp Val
 145 150 155 160

Asn Met Ala Leu Leu Leu Gln Gly Gly Gly His Tyr Arg Cys Asp Phe
 165 170 175

Arg Thr Thr Tyr Lys Ala Lys Lys Val Val Gln Leu Pro Asp Tyr His
 180 185 190

Phe Val Asp Leu Arg Thr Glu Ile Thr Ser His Asp Lys Asp Tyr Asn
 195 200 205

Lys Val Lys Leu Tyr Glu His Ala Lys Ala His Ser Gly Leu Pro Arg
 210 215 220

Leu Ala Lys
 225

<210> SEQ ID NO 21
 <211> LENGTH: 684
 <212> TYPE: DNA
 <213> ORGANISM: Favia fавus
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(681)

<400> SEQUENCE: 21

atg agt gtg att aca tca gaa atg aag atg gag ctg cgt atg gaa ggc Met Ser Val Ile Thr Ser Glu Met Lys Met Glu Leu Arg Met Glu Gly 1 5 10 15	48
gct gta aac ggg cac aag ttc gtg att aca ggg aaa gga agt ggc cag Ala Val Asn Gly His Lys Phe Val Ile Thr Gly Lys Gly Ser Gly Gln 20 25 30	96
cct ttc gag gga ata cag aat atg gac ctg aca gtc ata gag ggc gga Pro Phe Glu Gly Ile Gln Asn Met Asp Leu Thr Val Ile Glu Gly Gly 35 40 45	144
cct ctt cct ttt gct ttc gat atc ctg aca aca gca ttc ggt cac ggc Pro Leu Pro Phe Ala Phe Asp Ile Leu Thr Thr Ala Phe Gly His Gly 50 55 60	192
aac cgg gta ttt gtc aaa tac cca gaa gaa ata gta gac tac ttc aag Asn Arg Val Phe Val Lys Tyr Pro Glu Glu Ile Val Asp Tyr Phe Lys 65 70 75 80	240
cag tcg ttt cct gag ggt tat tct tgg gaa cga agc atg agt tac gaa Gln Ser Phe Pro Glu Gly Tyr Ser Trp Glu Arg Ser Met Ser Tyr Glu 85 90 95	288
gac ggg gga att tgc ctc gcc aca aac aat ata acg atg aag aaa gac Asp Gly Gly Ile Cys Leu Ala Thr Asn Asn Ile Thr Met Lys Lys Asp 100 105 110	336
ggc agc aac tgt ttt gtc tat gaa att cga ttt gat ggt gtg aac ttt Gly Ser Asn Cys Phe Val Tyr Glu Ile Arg Phe Asp Gly Val Asn Phe 115 120 125	384
cct gcc aat ggt cca gtt atg cag agg aag acc gtc aaa tgg gag cca	432

-continued

```

<210> SEQ ID NO 23
<211> LENGTH: 225
<212> TYPE: PRT
<213> ORGANISM: Trachyphyllia geoffroyi

<400> SEQUENCE: 23

Met Ser Leu Ile Lys Pro Glu Met Lys Ile Lys Leu Leu Met Glu Gly
1          5          10          15
Asn Val Asn Gly His Gln Phe Val Ile Glu Gly Asp Gly Lys Gly His
20          25          30
Pro Phe Glu Gly Lys Gln Ser Met Asp Leu Val Val Lys Glu Gly Ala
35          40          45
Pro Leu Pro Phe Ala Tyr Asp Ile Leu Thr Thr Ala Phe His Tyr Gly
50          55          60
Asn Arg Val Phe Ala Lys Tyr Pro Asp His Ile Pro Asp Tyr Phe Lys
65          70          75          80
Gln Ser Phe Pro Lys Gly Phe Ser Trp Glu Arg Ser Leu Met Phe Glu
85          90          95
Asp Gly Gly Val Cys Ile Ala Thr Asn Asp Ile Thr Leu Lys Gly Asp
100         105         110
Thr Phe Phe Asn Lys Val Arg Phe Asp Gly Val Asn Phe Pro Pro Asn
115         120         125
Gly Pro Val Met Gln Lys Lys Thr Leu Lys Trp Glu Ala Ser Thr Glu
130         135         140
Lys Met Tyr Leu Arg Asp Gly Val Leu Thr Gly Asp Ile Thr Met Ala
145         150         155         160
Leu Leu Leu Lys Gly Asp Val His Tyr Arg Cys Asp Phe Arg Thr Thr
165         170         175
Tyr Lys Ser Arg Gln Glu Gly Val Lys Leu Pro Gly Tyr His Phe Val
180         185         190
Asp His Cys Ile Ser Ile Leu Arg His Asp Lys Asp Tyr Asn Glu Val
195         200         205
Lys Leu Tyr Glu His Ala Val Ala His Ser Gly Leu Pro Asp Asn Val
210         215         220

Lys
225
    
```

The invention claimed is:

1. An isolated DNA comprising any of the following nucleotide sequences:

(a) the nucleotide sequence shown in SEQ ID NO: 2, 13, 15, 17, 19, or 21; or

(b) the nucleotide sequence of SEQ ID NO: 2 with 1 to 60 nucleotide substitutions, and encoding a fluorescent protein.

2. A recombinant vector having the DNA of claim 1.

3. A transformant having the DNA of claim 1.

* * * * *