



(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
12.01.2005 Bulletin 2005/02

(51) Int Cl.7: **C12N 9/54**, C12R 1/07
// C12N9:54, C12R1:07,
C12N1:20, C12R1:07

(21) Application number: **99973990.7**

(86) International application number:
PCT/JP1999/006392

(22) Date of filing: **16.11.1999**

(87) International publication number:
WO 2001/016302 (08.03.2001 Gazette 2001/10)

(54) **NOVEL THERMOSTABLE COLLAGEN-DIGESTING ENZYME, NOVEL MICROORGANISM
PRODUCING THE ENZYME AND PROCESS FOR PRODUCING THE ENZYME**

HITZESTABILES COLLAGEN-ABBAUENDES ENZYM, MIKROORGANISMUS UND VERFAHREN
ZU DESSEN HERSTELLUNG

NOUVELLE ENZYME THERMOSTABLE DIGERANT LE COLLAGENE, NOUVEAU
MICRO-ORGANISME PRODUISANT L'ENZYME ET METHODE DE PRODUCTION DE L'ENZYME

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**

(56) References cited:
US-A- 4 315 988

(30) Priority: **30.08.1999 JP 24281699**

(43) Date of publication of application:
22.08.2001 Bulletin 2001/34

(83) Declaration under Rule 28(4) EPC (expert
solution)

(73) Proprietor: **Japan Science and Technology
Agency
Kawaguchi-shi Saitama (JP)**

(72) Inventors:
• **NISHINO, Tokuzo**
Sendai-shi, Miyagi-ken 989-3204 (JP)
• **NAKAYAMA, Toru**
Sendai-shi, Miyagi-ken 981-3202 (JP)
• **TSURUOKA, Naoki**
Sendai-shi, Miyagi-ken 982-0021 (JP)
• **AKAI, Minoru**
Sendai-shi, Miyagi-ken 989-3202 (JP)

(74) Representative: **Cresswell, Thomas Anthony**
J.A. KEMP & CO.
14 South Square
Gray's Inn
London WC1R 5JJ (GB)

- **SAKO Y ET AL:** "An extremely heat-stable extracellular proteinase (aeropyrolysin) from the hyperthermophilic archaeon Aeropyrum pernix K1" **FEBS LETTERS, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 415, no. 3, 6 October 1997 (1997-10-06), pages 329-334, XP004261167 ISSN: 0014-5793**
- **ASDORNITHEE S. ET AL.:** 'Isolation and characterization of a collagenolytic enzyme from bacillus licheniformis N22' **JOURNAL OF FERMENTATION AND BIOENGINEERING vol. 78, no. 4, 1994, pages 283 - 287, XP002946891**
- **KAWAHARA H. ET AL.:** 'Isolation and characterization of new type of collagenase producing bacterium, bacillus alvei DC-1' **BIOSCI. BIOTECH. BIOCHEM. vol. 57, no. 8, 1993, pages 1372 - 1373, XP002935559**
- **MAKINEN K.K. ET AL.:** 'Purification and properties of an extracellular collagenolytic protease produced by the human oral bacterium bacillus cereus (strain soc 67)' **J. BIOL. CHEM. vol. 262, no. 26, 1987, pages 12488 - 12495, XP002935560**
- **KABADJOVA P. ET AL.:** 'Isolation and distribution of streptomyces populations with heat-resistant collagenolytic activity from two protein-rich areas in Bulgarian soil' **FOLIA MICROBIAL. vol. 41, no. 5, 1996, pages 423 - 429, XP002935561**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

FIELD OF THE INVENTION

5 **[0001]** The present invention relates to a thermostable collagen-decomposing enzyme produced by a new microorganism, and said new microorganism and a method for production of said enzyme by said microorganism. Concretely, the present invention relates to a novel thermostable collagen-decomposing enzyme having the highest reactivity (substrate specificity) to collagen, produced by a novel microorganism of the genus *Bacillus*, said novel microorganism and a process for producing of said enzyme by said microorganism.

DESCRIPTION OF THE PRIOR ART

10 **[0002]** Collagen-decomposing and gelatin-decomposing enzymes have been widely used in industry. For example, collagen peptides, which are hydrolytic products from collagen by these enzymes, are useful as material for cosmetics, because of their interesting physiological activities such as moisture-keeping effects or immunity-activation activity. Therefore, collagen peptides are widely used for medical and cosmetic purposes. Further, gelatin, which is a denatured form of collagen, is used as the coating material for photograph films, and gelatin-decomposing enzymes are used for the recycling of the photograph and X-ray films. Many kinds of proteases are known to decompose gelatin; however, the decomposition of collagen by these proteases is still difficult. For hydrolysis of collagen, specific metal proteases, named "collagenase", should be used.

15 **[0003]** Recently, many attempts have been made for the effective use of the organic wastes. For instance, the composting of garbage wastes is one concrete example of the biorecycling of organic wastes. More than 30% of animal protein are composed of collagen, therefore, garbage wastes produced from daily kitchen activities in houses and restaurants and the wastes from meat processing factories should contain large quantities of collagen.

20 **[0004]** Because of specific, highly ordered structure, collagen is generally insoluble in water and difficult to be decomposed, and therefore degradation of collagen proceeds very slowly during composting. Most part of unusable portions produced from livestock industries are composed of collagen, and, therefore, are treated by incineration, causing problems such as anathermal of the earth or the generation of carbon dioxide or dioxin which cause the air pollution.

25 **[0005]** These problems must be solved from the view point to make an effective use of materials.

30 **[0006]** It is well known that the temperature of the organic wastes raises to 50-65°C or higher during the composting process. Therefore, if thermostable enzymes or thermophilic microorganisms which are active even under such high-temperature composting conditions are used, the composting of organic waste should proceed more effectively.

35 **[0007]** Nowadays, industrial collagenases are those from microorganisms (bacteria), and as a concrete example, an actinomycetous collagenase of the genus *Streptomyces* can be mentioned. Other microbial collagenases are also known; for instance, collagenases from *Clostridium histolyticum* (Biochemistry 1984, 23, 3077-3085) and *Cytophaga* sp. (Biosci, Biotech, Biochem., 1993, 57, 1894-1898) are the concrete examples.

40 **[0008]** Concerning the example of enzyme which is industrially used, it is necessary for the enzyme to be thermostable from the view point of treating speed and the subject to be treated.

45 **[0009]** However, all known collagenases are of mesophilic origin and lacks of thermostability, and these circumstances hampers their efficient industrial applications. Until now, a collagen-decomposing enzyme with sufficient thermostability (having high optimum temperature) for the industrial applications has not yet been developed.

50 **[0010]** Usually, it is difficult to use collagen in an industrial scale because there is no thermostable enzyme to act effectively in an industrial scale. Therefore, it is obvious that above mentioned problem can be solved perfectly, if an enzyme having high activity to collagen is developed.

55 **[0011]** As mentioned above, the object of this invention is to find out a thermostable collagen-decomposing enzyme.

[0012] The inventors of this invention have carried out an intensive study to find a microorganism producing a thermostable collagen-decomposing enzyme in nature, and have found a promising thermophilic bacterium belonging to the genus *Bacillus* genus that produces said enzyme in the soil of Sendai, Japan and accomplished the present invention.

[0013] The microorganism, which is used to produce a thermostable collagen-decomposing enzyme of this invention, belongs to the genus *Bacillus*, and is termed strain NTAP-1. This strain has been deposited according to the requirement of deposit based on Budapest treaty in the Biotic Technology Industries Institute of the Agency of Industrial Science of Technology belonging to the Ministry of International Trade and Industry Japan and accepted by the accept number of FERM BP-6926 on November 1, 1999. (This strain is originally deposited on August 27, 1999 under accession number FERM P-17535.) (in the specification, this strain is shortened only as NTAP-1 strain.)

[0014] The inventors have found that the industrially useful enzyme can be obtained by the use of this strain, and the obtained enzyme can be used as the catalyst for bioconversion.

DETAILED DESCRIPTION OF THE INVENTION

[0015] The first important point of this invention is a thermostable collagen-decomposing enzyme obtainable from a bacterium belonging to a Bacillus genus, wherein: (1) the bacterium is Gram-negative or Gram-indefinite; (2) the bacterium has spore forming ability; (3) the bacterium is motile, (4) the bacterium grows at 70°C, does not grow at 30°C or 80°C (5) grows at pH 5 and does not grow at pH 7; (6) the bacterium is rod-shaped; (7) the bacterium is negative to catalase; (8) the bacterium is negative to oxidase; (9) the bacterium is negative to O/F test; (10) the bacterium has acetoin-producing activity; and (11) the bacterium has gelatin-decomposing activity, and said enzyme is further characterized by the following features: (a) the enzyme has greater hydrolysis activity towards collagen and gelatin than casein and albumin; (b) optimum reaction pH is between pH 3.5 and 4.5; (c) optimum reaction temperature is between 65°C and 70°C; (d) the enzyme retains more than 60% of its original activity after heat treatment at 60°C and pH 6.0 for 4 hours; (e) the enzyme is stable between pH 3 to 6; and (f) molecular weight of the enzyme estimated by SDS-polyacrylamide gel electrophoresis is approximately 46,000.

[0016] Accordingly, an excellent action and effect which can be used for the decomposition of collagen at 70°C; or lower temperatures can be expected. And more desirably, said thermostable collagen-decomposing enzyme of this invention is produced by the microorganism belonging to the genus Bacillus or Bacillus sp. strain NTAP-1.

[0017] The second important point of this invention is a producing method of the thermostable collagen-decomposing enzyme of the invention comprising, using a microorganism which has following features, that is, by the following features: (1) the bacterium is Gram-negative or Gram-indefinite, (2) the bacterium has a spore forming ability, (3) the bacterium is motile, (4) the bacterium grows at 70°C, does not grow at 30°C or 80°C and grows at pH 5, does not grow at pH 7, (6) the bacterium is rod-shaped, (7) the bacterium is negative to catalase, (8) the bacterium is negative to oxidase, (9) the bacterium is negative to O/F test, (10) the bacterium has acetoin producing activity and (11) the bacterium has gelatin decomposition activity, purifying and accumulating the thermostable collagen-decomposing enzyme.

[0018] Desirably, the producing method of said thermostable collagen-decomposing enzyme, wherein the microorganism belonging to the genus Bacillus is the Bacillus genus bacteria NTAP-1 strain.

[0019] The third important point of this invention is a new developed microorganism belonging to a Bacillus genus, which produces said thermostable collagen-decomposing enzyme. desirably, said microorganism is the strain titled as Bacillus sp. NTAP-1 and have deposited according to the requirement of deposit based on Budapest treaty in the Biotic Technology Industries Institute of the Agency of Industrial Science of Technology belonging to the Ministry of International Trade and Industry Japan and accepted by the accession number FERM BP-6926 on November 1, 1999 (This strain is originally deposited on August 27, 1999 under accession number FERM P-17535.).

[0020] The inventors of this invention have found that among the microorganism belonging to Bacillus genus there is a novel microorganism which produces thermostable collagen-decomposing enzyme, and have accomplished the present invention.

BRIEF ILLUSTRATION OF THE DRAWINGS

[0021] FIG.1 is a graph showing the heat stability of the thermostable collagen-decomposing enzyme, FIG.2 is a graph showing the pH-stability of said thermostable collagen-decomposing enzyme, FIG.3 is a graph showing the temperature-dependence of the reaction of said thermostable collagen-decomposing enzyme and FIG.4 is a graph showing the pH-dependence of the reaction of said thermostable collagen-decomposing enzyme.

THE BEST EMBODYMENT OF THE INVENTION

[0022] The present invention will be illustrated more in detail.

A. The microbiological features of bacteria used to produce a thermostable collagen-decomposing enzyme are mentioned above. Further, this microorganism can be preserved by freezing method (-80°C around).

B. Growing condition

name of cultivate medium: GGY medium

components of medium: medium containing 1.5% of glucose, 1.5% of gelatin and 0.01 % of yeast extract.

pH of medium: 4.8

sterilizing condition of medium: 20 minutes at 120°C

temperature of medium: 60°C

aerobic condition

C. component of protecting agent: 30% glycerol aqueous solution (not necessary to adjust pH of protecting agent)

EP 1 126 024 B1

not necessary to adjust pH of protecting agent
sterilizing condition of protecting agent: 20 minutes at 120°C

[0023] The characteristics of the thermostable collagen-decomposing enzyme of this invention will be illustrated more minutely with reference to the drawings.

FIG.1 is the graph showing the relative remaining activity of the thermostable collagen-decomposing enzyme after heat treatment at various temperature for 1 hour, and in this graph, the activity after heat treatment at 30°C is taken to be 100%.

FIG.2 is the graph showing the relative remaining activity of the thermostable collagen-decomposing enzyme after treatment at various pHs for 1 hour, and in this graph, the activity after treatment at pH 4.1 is taken to be 100%.

FIG.3 is the graph showing the relative activity of the thermostable collagen-decomposing enzyme at various temperature, and in this graph, the enzyme activity at 60°C that indicates maximum activity is taken to be 100%.

FIG.4 is the graph showing the relative activity of the thermostable collagen-decomposing enzyme at various pHs, and in this graph, the enzyme activity at pH 3.8 that indicates maximum activity value is taken to be 100%.

EXAMPLES

[EXAMPLE 1]

[0024] Various kinds of specimen such as soils, composts, river and lake waters are diluted to 100-10,000 times with 0.85% NaCl, and 0.1ml of said diluted solution was spread on GGY agar-agar medium, then are left for 2 or 3 days at 70°C. The colony grown on medium was isolated and inoculated in 5 ml of GGY liquid medium and cultivated with shaking for 2 or 3 days at 70°C. The collagen-decomposing enzyme activity of several hundred kinds of isolates are evaluated using the supernatant of culture according to the method described in Example 2.

[0025] The strain that indicates the highest collagen-decomposing activity was selected and named it NTAP-1 strain.

[0026] The taxonomical characteristics of NTAP-1 strain can be illustrated as follows.

(1) cell morphology: rod-shaped (0.8×2-3μm), curved and becomes chain form by aging.

(2) Gram's staining: negative or indefinite

(3) spore forming ability: yes

(4) motility: yes

(5) shape and characteristic of colonies: circular, corrugated or slightly convex, having smooth surface and transparent.

(6) growing temperature: grows at 70°C, but does not grow at 80°C.

(7) catalase: negative

(8) oxidase: negative

(9) O/F test: negative

(10) biochemical test :

[0027] Decomposes

glucose, fructose, sorbose, D-arabinose, L-arabinose, ribose, D-xylitol, L-xylitol, D-turanose, L-turanose, D-lyxose, D-tagatose, 5-ketogluconic acid.

[0028] Does not decompose

glycelol, erythritol, adonitol, β-methyl-D-xylose, galactose, mannose, rhamnose, dulcitol, α-methyl-D-mannose, α-methyl-D-glucose, N-acetyl-glucosamine, amidagline, arbutin, aesuculin, salicin, cellobiose, maltose, milk sugar, melibiose, cane sugar, trehalose, inulin, melezitose, raffinose, glycogen, xylitol, gentiobiose, D-fucose, L-fucose, D-arabitol, L-arabitol, 2-ketogluconic acid.

Enzyme activity	
β-galactosidase	negative
arginine dihydrolase	negative
lysine decarboxylase	negative
urease	negative
tryptophan deaminase	negative
gelatinase	positive

Others	
use of citric acid	no
production of H ₂ S	no
production of indole	negative
production of acetoin	positive
reduction of nitrate	positive
anaerobic growth	slightly observed
growth at pH 7	no
growth at pH 5.1	yes
growth at 30°C	no

[0029] From above mentioned taxonomic features, the taxonomic positioning of this bacteria is referred in Bergey's Manual of Systematic Bacteriology, vol 2 p1104-1139, author: S.H. Sneath, editor: P.H. Snrth et al. (publisher: Williams & Willkins).

[0030] This bacteria is a spore-forming rod-shaped bacterium. Although Gram-negative nature of the bacterium is distinct from Gram-positive nature of known species of the genus Bacillus, it is recognized that it is a strain of the genus Bacillus because it grows aerobically.

[0031] Among known species of the genus Bacillus, B. acidocaldarius, B.lichenformis, or B. coagulans are known to be thermophilic and acidophilic. However, this strain should not be B. lichenformis and B. coagulans because B. lichenformis and B. coagulans are catalase-positive and can grow at 40°C but not at 65 °C. Also, it is different from the standard species of B.acidocaldarius because it produces acetoin. Therefore, it is not possible to confirm that whether it is a modified species of B. acidocaldarius or it belongs to a different species; the species of this bacteria can not be specified.

[EXAMPLE 2]

[0032] 5 ml of medium (pH 4.8) containing 1.5% of glucose, 1.5% of gelatin and 0.01% of yeast extract is poured into 5ml test tube and sterilized for 20 minutes at 120°C. NTAP-1 (the shortened name of Bacillus genus NTAP-1 to discriminate the microorganism of this invention) is inoculated on said medium and cultivated with shaking for 4 days. The culture medium is centrifuged for 20 minutes at 8,000 r.p.m., and the activity of the thermostable collagen-decomposing enzyme in the supernatant is measured. Namely, 0.4ml of enzyme liquid is mixed to 0.1 ml of 1M sodium acetate buffer (pH 4.5) and the mixture is pre-incubated for 5 minutes at 60°C. Then 3mg of Azocoll (azo dye-linked collagen powder: product of Sigma Co., Ltd.) is suspended, and enzyme reaction is carried out at 60°C with stirring for 1 hour. After the reaction, the reaction mixture is chilled on ice, and insoluble Azocoll was separated by centrifugation.

[0033] During the enzyme reaction, Azocoll is decomposed by the enzyme and the supernatant turns red. By measuring the absorbance at 518 nm of the supernatant, the activity of the thermostable collagen-decomposing enzyme is estimated. The amount of enzyme which makes the absorbance at 518 nm increases 0.001 by 1 minute under said condition is defined as 1 unit (U). The concentration of enzyme activity of the obtained supernatant liquid of the cultivated liquid is 3.1 U/ml.

[EXAMPLE 3]

[0034] 6 litter of same medium to Example 2 is poured into a jar fermentor of 10-liter vessel. After sterilized for 30 minutes at 120°C, 200 ml of the NTAP-1 culture is inoculated on said medium. The cultivation is carried out at 60°C with 6 liters/min aeration for 4 days. The thermostable collagen-decomposing activity of the culture supernatant is measured. The activity of enzyme of the supernatant liquid is 5.0 U/ml.

[0035] The resultant supernatant was used as the starting material, the purification and concentration of thermostable collagen-decomposing enzyme was carried out according to the following process.

[0036] Ammonium sulfate is added to the supernatant liquid and the precipitate formed by ammonium sulfate 40% saturation was collected and dissolved in 585 ml of 0.01M acetate buffer (pH 5.0). Phenyl-Sepharose (product of AmeshamPharmacia Biotech.) was added to the solution, stirred and mixed for 1 hour, then the mixture was filtrated and the resin is separated. The enzyme activity absorbed to the resin was eluted by washing the resin with 1:1 mixture of 0.01M acetic acid buffer (pH 5.0) and ethylene glycol, and the active fraction was then dialyzed against 0.01 M phosphate buffer (pH 7.0).

[0037] Then, the solution was passed through DEAE-Sephadex (AmeshamPharmacia Biotech) column which is pre-

EP 1 126 024 B1

viously equilibrated with 0.01 M phosphate buffer (pH 7.0). The linear gradient (0-1M) of sodium chloride was used to elute the enzyme activity from the column. The eluate was fractioned into about 90 fractions, and the activity of enzyme of each fractions are measured according to the method described in Example 1.

[0038] Then, ammonium sulfate is added to the active fractions (233 ml) to 20% saturation. The enzyme activity is absorbed to a column by pass the enzyme solution through the phenyl-Sepharose (AmeshamPharmacia Biotech) column which is previously equilibrated with 0.01 M acetic acid buffer (pH 5.0).

[0039] Then, the enzyme activity adsorbed to the column was eluted by washing the column with a linear concentration slope (0-50%) of ethylene glycol in the equilibration buffer. The eluated solution is divided into 90 fractions approximately, and the activity of enzyme of each fraction is measured according to the measuring method described in Example 1. From fractions of eluated solution, active fractions (total 48ml) are collected.

[0040] This solution is dialyzed against 0.01M phosphate buffer (pH 7.0), then applied to a column of MONO-Q (AmeshamPharmacia Biotech) which is previously made equilibrated with the same buffer. The activity was eluted by washing the column with a linear gradient (0-1M) of sodium chloride.

[0041] Active fractions (10.6 ml) were collected and concentrated to 0.5 ml using Centricon (centrifuge concentrator: Amicon Co., Ltd.: said concentrator has an ultrafiltration membrane made of cellulose derivative at the bottom of the container. When mixed solution composed of enzyme and protein is poured into the container and centrifuged by 6000 r.p.m., protein contained in the solution is remained on the membrane, while water or low molecular weight ion passes through the film and recovered as a filtrated liquid.), and divided by a gel filtrating chromatography method, and 1.2 ml of fraction having higher activity is collected. The activity yield of the dissolved fraction from the cultivated liquid is 1.4%, and the concentration of the activated enzyme is 416 U/ml.

[0042] The enzyme solution thus obtained was analyzed by SDS-polyacrylamide gel electrophoresis according to Laemmli procedure (Laemmli, U.K., Nature, 1970, 227, 680-685), and molecular weight of the thermostable collagen-decomposing enzyme is estimated to be approximately 46,000.

[EXAMPLE 4]

(Experiment to investigate the substrate specificity of the thermostable collagen-decomposing enzyme)

[0043] After 5 mg of Azocoll is suspended in 0.4 ml of 0.01 M sodium acetate buffer (pH 4.5), 0.1 ml of enzyme solution, whose enzyme concentration is adjusted properly by dilution, is added and allowed to react for 1 hour at 60°C with constant shaking. After the reaction, the reaction mixture was chilled on ice for 1 hour and then centrifuged. The absorbance at 518 nm of the supernatant is measured. The reaction mixture prepared by same process except using water instead of enzyme solution was used as a blank solution. The absorbance at 518 nm of the reaction mixture whose added Azocoll was perfectly solubilized is measured. The same experiments are carried out on various enzyme concentrations, and the amount of enzyme which gives approximately 50% degradation (to solubilize approximately 2.5 mg Azocoll) under these condition is determined.

[0044] By the same process as mentioned above, 5 mg of collagen, gelatin, casein or cow serum albumin (all are the products Nacalai Tesque, Co.) are respectively suspended (or dissolved) in 0.4 ml of 0.01 M sodium acetate buffering solution (pH 4.5), then 0.1 ml of enzyme solution of previously decided concentration is added and reacted for 1 hour at 60°C with constant shaking. After the reaction, the reaction mixture was kept at 4°C for 20 minutes and centrifuged. In cases which use casein or bovine serum albumin, 0.5 ml of 50% trichloroacetic acid is added to the reaction mixture and was kept at 4°C for 20 minutes then centrifuged. The adsorption at 280 nm of the supernatant was measured. From the adsorption value at 280 nm when each proteins are perfectly solubilized, the decomposing ratio of each proteins are measured. When the decomposing rate of Azocoll is regarded as 100%, the relative solubilizing rate of each proteins are listed in Table 1.

Table 1

Relative solubilizing rate of each proteins, when decomposing rate of Azocoll is regarded as 100%	
collagen	+++
gelatin	+++
casein	±
albumin	±

+++ : has relative activity greater than 80% to Azocoll

± : has relative activity lower than 20% to Azocoll

EP 1 126 024 B1

Table 1 (continued)

Relative solubilizing rate of each proteins, when decomposing rate of Azocoll is regarded as 100%	
keratin	-
remarks	

- : not reacted

[0045] The obtained results are shown in Fig.1. This enzyme is stable up to 60°C. In the meanwhile, the stability of enzymes when enzymes are treated by various pH are investigated. The buffer to be used in specific pH range are listed below.

- pH 2.5 to 3.5 :1M glycine-HCl buffer
- pH 3.5 to 5.5: 1 M sodium acetate buffer
- pH 6.0 to 8.0: 1M sodium phosphate buffer
- pH 8.0 to 9.0: 1M glycine-NaOH buffer
- pH 9.0 to 10.0: 1M sodium phosphate buffer

After 0.0025 ml of these buffers and 0.0025 ml of 1% aqueous solution of Tween 80 are added to 0.02 ml of enzyme liquid, placed at the temperature of 60°C for 1 hour. Then, 0.05 ml of 1 M acetic acid buffer (pH 4.0) is added to this treated enzyme liquid and the activity is measured. The obtained result indicates that this enzyme is stable at the pH range from 3 to 6 (Fig.2).

[EXAMPLE 7]

(Experiment to investigate the of the optimum reaction temperature and optimum reaction pH of thermostable collagen-decomposing enzyme)

[0046] The activity of the thermostable collagen-decomposing enzyme of this invention was measured at 30, 40, 50, 60, 70 and 80°C. The method described in Example 1 was used except changing the reaction temperature. The results showed that this enzyme exhibited the highest activity at 60°C (refer to Fig. 3).

[0047] Secondly, the activity of this enzyme is measured at various (from pH 2.5 to 7.2). The method for activity measurement was based on the method described in Example 1 except changing buffer component to be used in the reaction system as follows: 0.01 M glycine-HCl buffer (pH 2.5 to 7.2), 0.01 M sodium acetate buffer (pH 4 to 6) or 0.01 M potassium phosphate buffer (pH 6 to 8). The results showed that this enzyme showed the highest activity at pH 3.7 to 3.9 (refer to Fig. 4).

POTENTIALS FOR THE INDUSTRIAL USE

[0048] Obviously from the above mentioned Examples, by the present invention, it becomes possible to prepare effectively a thermostable collagen-decomposing enzyme which is excellent at the optimum temperature, optimum pH and collagen substrate specificity by the use of above mentioned novel microorganism. Therefore, the present invention make it possible to utilize the materials which are not utilized in livestock industries, and largely contribute to the production of collagen peptides that has potential applications in medical, pharmaceutical, and food industries.

Claims

1. A thermostable collagen-decomposing enzyme obtainable from a bacterium belonging to a Bacillus genus, wherein: (1) the bacterium is Gram-negative or Gram-indefinite; (2) the bacterium has spore forming ability; (3) the bacterium is motile, (4) the bacterium grows at 70°C, does not grow at 30°C or 80°C (5) grows at pH 5 and does not grow at pH 7; (6) the bacterium is rod-shaped; (7) the bacterium is negative to catalase; (8) the bacterium is negative to oxidase; (9) the bacterium is negative to O/F test; (10) the bacterium has acetoin-producing activity; and (11) the bacterium has gelatin-decomposing activity, and said enzyme is further **characterized by** the following features: (a) the enzyme has greater hydrolysis activity towards collagen and gelatin than casein and albumin; (b) optimum reaction pH is between pH 3.5 and 4.5; (c) optimum reaction temperature is between 65°C and 70°C, (d) the enzyme retains more than 60% of its original activity after heat treatment at 60°C and pH 6.0 for 4 hours, (e) the enzyme is stable between pH 3 to 6; and (f) molecular weight of the enzyme estimated by SDS-polyacry-

lamide gel electrophoresis is approximately 46,000.

2. An enzyme according to claim 1, wherein the bacterium is *Bacillus* genus bacteria NTAP-1 strain.
- 5 3. A method for producing an enzyme according to claim 1 or 2, the method comprising using a bacterium as defined in claim 1 to accumulate the enzyme in a culture medium, purifying the enzyme and collecting it.
4. A method according to claim 3, wherein the bacterium is *Bacillus* genus bacteria NTAP-1 strain.
- 10 5. A bacterium belonging to a *Bacillus* genus which bacterium produces an enzyme according to claim 1.
6. A bacterium according to claim 5 which is *Bacillus* sp. NTAP-1 deposited under the Budapest treaty in the Biotic Technology Industries Institute of the Agency of Industrial Science of Technology belonging to the Ministry of International Trade and Industry Japan and accepted under the accession number FERM BP-6926 on November 1, 1999 (deposited under the original accession number FERM P-17535 on August 27, 1999).
- 15 7. Use of an enzyme according to claim 1 or 2 for decomposing collagen.

20 Patentansprüche

1. Wärmestabiles, Kollagen-abbauendes Enzym, das aus einem Bakterium, das zu einer *Bacillus*-Gattung gehört, erhältlich ist, wobei (1) das Bakterium Gramnegativ oder Gram-undefiniert ist; (2) das Bakterium eine Sporenbildungsfähigkeit aufweist; (3) das Bakterium bewegungsfähig ist; (4) das Bakterium bei 70 °C wächst und bei 30 °C oder 80 °C nicht wächst, (5) bei pH 5 wächst und bei pH 7 nicht wächst; (6) das Bakterium stäbchenförmig ist; (7) das Bakterium gegenüber Katalase negativ ist; (8) das Bakterium gegenüber Oxidase negativ ist; (9) das Bakterium gegen einen O/F-Test negativ ist; (10) das Bakterium eine Acetoin-produzierende Aktivität aufweist; und (11) das Bakterium eine Gelatin-abbauende Aktivität aufweist, und das Enzym ferner durch folgende Merkmale **gekennzeichnet** ist: (a) das Enzym weist höhere Hydrolyse-Aktivität gegenüber Kollagen und Gelatin als gegenüber Kasein und Albumin auf; (b) der optimale pH-Wert der Umsetzung liegt zwischen pH 3,5 und 4,5; (c) die optimale Temperatur der Umsetzung liegt zwischen 65 °C und 70 °C; (d) das Enzym behält mehr als 60 % seiner ursprünglichen Aktivität nach einer vierstündigen Wärmebehandlung bei 60 °C und einem pH von 6,0; (e) das Enzym ist zwischen pH 3 und 6 stabil; und (f) das durch SDS-Polyacrylamidgelelektrophorese abgeschätzte Molekulargewicht des Enzyms beträgt etwa 46 000.
- 25 2. Enzym gemäß Anspruch 1, wobei das Bakterium zu dem NTAP-1-Bakterien-Stamm der Gattung *Bacillus* gehört.
3. Verfahren zur Herstellung eines Enzyms gemäß Anspruch 1 oder 2, wobei das Verfahren das Verwenden eines Bakteriums wie in Anspruch 1 definiert zur Anreicherung des Enzyms in einem Kulturmedium, Reinigen des Enzyms und Sammeln desselben umfasst.
- 30 4. Verfahren gemäß Anspruch 3, wobei das Bakterium zu dem NTAP-1-Bakterien-Stamm der Gattung *Bacillus* gehört.
5. Bakterium, das zu einer *Bacillus*-Gattung gehört, wobei das Bakterium ein Enzym gemäß Anspruch 1 herstellt.
- 45 6. Bakterium gemäß Anspruch 5, das *Bacillus* sp. NTAP-1 ist, das gemäß Budapester Vertrag bei dem Biotic Technology Industries Institute of the Agency of Industrial Science of Technology, das zum Ministry of International Trade and Industry Japan gehört, hinterlegt ist und unter der Hinterlegungsnummer FERM BP-6926 am 1. November 1999 (hinterlegt am 27. August 1999 unter der ursprünglichen Hinterlegungsnummer FERM P-17535) angenommen wurde.
- 50 7. Verwendung eines Enzyms gemäß Anspruch 1 oder 2 zum Abbauen von Kollagen.

55 Revendications

1. Enzyme thermostable décomposant le collagène, pouvant être obtenue à partir d'une bactérie appartenant à un genre *Bacillus*, dans laquelle : (1) la bactérie est Gram-négative ou Gram- indéfinie; (2) la bactérie a une capacité

EP 1 126 024 B1

à former des spores; (3) la bactérie est mobile; (4) la bactérie croît à 70°C, ne croît pas à 30°C ou 80°C; (5) croît à un pH de 5 et ne croît pas à un pH de 7; (6) la bactérie est en forme de bâton; (7) la bactérie est négative à la catalase; (8) la bactérie est négative à l'oxydase; (9) la bactérie est négative au test O/F; (10) la bactérie a une activité de production d'acétoïne; et (11) la bactérie a une activité de décomposition de la gélatine, et ladite enzyme est en outre **caractérisée par** les particularités suivantes : (a) l'enzyme a une plus grande activité d'hydrolyse à l'égard du collagène et de la gélatine qu'à l'égard de la caséine et de l'albumine; (b) le pH de réaction optimal est entre pH 3,5 et 4,5; (c) la température de réaction optimale est entre 65°C et 70°C; (d) l'enzyme conserve plus de 60 % de son activité d'origine après traitement thermique à 60°C et pH 6,0 pendant 4 heures; (e) l'enzyme est stable entre pH 3 à 6; et (f) le poids moléculaire de l'enzyme estimé par électrophorèse sur gel de SDS-polyacrylamide est d'approximativement 46 000.

2. Enzyme selon la revendication 1, dans laquelle la bactérie est une souche de bactérie NTAP-1 du genre Bacillus.

3. Procédé pour la production d'une enzyme selon la revendication 1 ou 2, le procédé comprenant l'utilisation d'une bactérie telle que définie dans la revendication 1 pour accumuler l'enzyme dans un milieu de culture, la purification de l'enzyme et son recueil.

4. Procédé selon la revendication 3, dans lequel la bactérie est une souche de bactérie NTAP-1 du genre Bacillus.

5. Bactérie appartenant à un genre Bacillus, laquelle bactérie produit une enzyme selon la revendication 1.

6. Bactérie selon la revendication 5, qui est Bacillus sp. NTAP-1 déposée conformément au traité de Budapest auprès du Biotic Technology Industries Institute de l'Agency of Industrial Science of Technology appartenant au Ministère japonais du Commerce International et de l'Industrie et acceptée sous le numéro d'accession FERM BP-6926 le 1er novembre 1999 (déposée sous le numéro d'accession d'origine FERM P-17535 le 27 août 1999).

7. Utilisation d'une enzyme selon la revendication 1 ou 2 pour la décomposition du collagène.

Fig. 1




Fig. 2




Fig. 3




Fig. 4

