



US 20080313752A1

(19) **United States**

(12) **Patent Application Publication**

Miyamoto et al.

(10) **Pub. No.: US 2008/0313752 A1**

(43) **Pub. Date: Dec. 18, 2008**

(54) **METHOD FOR DIFFERENTIATING MESENCHYMAL STEM CELLS INTO STEROID-PRODUCING CELLS**

(30) **Foreign Application Priority Data**

Mar. 3, 2004 (JP) 2004-058406

(76) Inventors: **Kaoru Miyamoto, Fukui (JP); Takashi Yazawa, Fukui (JP); Akihiro Umezawa, Tokyo (JP)**

Publication Classification

(51) **Int. Cl.**
A01K 67/027 (2006.01)
C12N 5/06 (2006.01)
C12P 33/00 (2006.01)

(52) **U.S. Cl.** 800/13; 435/377; 435/52

(57) **ABSTRACT**

To control the differentiation of mesenchymal stem cells into steroid hormone producing cells. Mesenchymal stem cells can be differentiated into steroid hormone-producing cells by being stimulated by a transcriptional factor (SF-1), preferably by the transcriptional factor (SF-1) and cAMP. The present invention is a method for differentiating mesenchymal stem cells into steroid hormone producing cells, comprising stimulating the mesenchymal stem cells by the transcriptional factor (SF-1). The mesenchymal stem cells may be further stimulated by cAMP.

Correspondence Address:

**JENKINS, WILSON, TAYLOR & HUNT, P. A.
Suite 1200 UNIVERSITY TOWER, 3100 TOWER
BLVD.,
DURHAM, NC 27707 (US)**

(21) Appl. No.: **10/591,530**

(22) PCT Filed: **Feb. 18, 2005**

(86) PCT No.: **PCT/JP05/02548**

§ 371 (c)(1),
(2), (4) Date: **Sep. 1, 2006**




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10




Figure 11




Figure 12




Figure 13





Figure 14



METHOD FOR DIFFERENTIATING MESENCHYMAL STEM CELLS INTO STEROID-PRODUCING CELLS

FIELD OF THE INVENTION

[0001] The present invention relates to a method for differentiating mesenchymal stem cells into steroid-producing cells.

PRIOR ART

[0002] It is well known that mesenchymal stem cells have multiple differentiation potential in vivo beyond germ layers (references 1-3). It is expected that the mesenchymal stem cells will play an important role in the fields of development and regeneration, since these stem cells are easily obtained from adults and are easily subjected to established cell lines after differentiation in contrast to other embryo stem cells. The method for differentiating mesenchymal stem cells into adipocytes, chondrocytes or osteoblasts has been examined already (reference 4).

[0003] reference 1: J. Clin. Invest. 10, 697-705, 1999

[0004] reference 2: Exp. Cell. Res. 288, 51-9, 2003

[0005] reference 3: Biochem. Biophys. Res. Commun. 295, 354-61, 2002

[0006] reference 4: WO2002/022788

PROBLEMS TO BE SOLVED BY THE INVENTION

[0007] The present inventors discovered a method for controlling the differentiation of mesenchymal stem cells derived from bone marrow into steroid-producing cells. The inventors believe that the method of the present invention will become very useful method in the field of development and regeneration.

MEANS TO SOLVE THE PROBLEMS

[0008] The present inventors discovered that mouse bone marrow mesenchymal stem cells were differentiated into Leydig cells, steroid hormone-producing cells, when transplanted into testis (Example 4). Furthermore, it was found that a limited portion of mouse bone marrow-derived mesenchymal stem cells KUM9 were spontaneously differentiated into steroid-producing cells during culturing, and at the same time, that a transcriptional factor SF-1 was expressed (Example 5). Therefore, the inventors prepared a KUM9-derived strain, wherein SF-1 is expressed constitutively, and added a stimulation by cAMP, then the inventor found that the KUM9-derived cells stably produced steroid hormone (Examples 1 and 2). Moreover, it was confirmed that not only a gonadal steroid hormone-producing enzymes but also adrenal steroid hormone-producing enzymes are induced in a human cell line (Example 3). It is expected that these results are fundamental data in using differentiation of stem cells for medical treatment of human steroid hormone deficiency and provide a useful method for regeneration treatment.

[0009] From the above results, the inventors discovered that stimulation of mesenchymal stem cells by a transcriptional factor (SF-1), preferably with SF-1 and cAMP, enabled to differentiate the mesenchymal stem cells into steroid-producing cells and accomplished the present invention.

[0010] Namely, the present invention is a method for differentiating bone marrow-derived mesenchymal stem cells into steroid hormone-producing cells, comprising stimulat-

ing the mesenchymal stem cells by a transcriptional factor (SF-1). Furthermore, said method may comprise further stimulating the mesenchymal stem cells by cAMP.

[0011] The mesenchymal stem cells are preferably derived from bone marrow and said mesenchymal stem cells are preferably human cells.

[0012] Still furthermore, the present invention is a method for producing steroid-producing cells, comprising producing steroid producing cells by implementing said method in vitro or by transplanting the mesenchymal stem cells into a mammalian reproductive organ.

[0013] Moreover the present invention is steroid-producing cells or non-human animals containing said steroid-producing cells obtained by these method.

ADVANTAGES OF THE PRESENT INVENTION

[0014] The present invention provides, for the first time, a method for differentiating mesenchymal stem cells into steroid hormone-producing cells.

[0015] It will be possible to obtain a large amount of differentiated cells by transplanting stem cells into a genital gland of a non-human animal such as swine, if immune difficulties were overcome. The most important issue in regeneration medicine and others is to maintain transplanted cells keeping the differentiated state and the present invention may be applied as a tool for the development of said method. Said differentiated cells may be used as model cells for the development of regeneration medicine.

BRIEF DESCRIPTION OF THE DRAWINGS

[0016] FIG. 1 shows pIRES-puro2 vector.

[0017] FIG. 2 shows KUM9 cells (A) and KUM9 cells stably transfected with parent vector (B) and with SF-1 expressing vector (C).

[0018] FIG. 3 shows the western blotting showing SF-1 protein detected in various cell lines. SF-7 and SF-9 are the 7th and 9th cell lines, respectively, which are established by transduction of SF-1.

[0019] FIG. 4 shows the result of RT-PCR for RNA isolated from KUM9 cells, which are stably transduced with parent vector (pIRES) and SF-1 (SF-9) and are cultured for 7 days in the presence of cAMP. Days shown on the upper row are days until extraction of RNA after the addition of cAMP. HSD3b (3b-HSD) expresses on the 3rd day and P450c17 and P450c11b1 express on the 5th day.

[0020] FIG. 5 shows the western blotting for proteins extracted from the cells cultured under a condition similar to that described in FIG. 4. Proteins from MA10-derived cells originated from mouse Leydig cells are used as a positive control.

[0021] FIG. 6 shows quantification by RIA of secreted steroid hormone, which is obtained by the recovery of supernatant of the medium used for the cells cultured under a condition similar to that described in FIG. 4. The amount of hormone produced within 24 hrs was determined by changing culture medium to fresh medium on the day before the each indicated day.

[0022] FIG. 7 shows immunostaining by anti-3 β HSD I antibody of SF-9 cells, which are fixed before adding cAMP (-) or after culture for 7 days in the presence of cAMP (+). DAPI (4',6-Diamidino-2-phenylindole, dihydrochloride) is a dye marker for DNA and shows the intracellular localization of nucleus.

8[0023] FIG. 8 shows the result of RT-PCR for RNA extracted from hMSC E6/E7 cells, which is derived from human bone marrow mesenchymal stem cells, are stably transduced with SF-1, and are cultured for 7 days in the presence of cAMP.

[0024] FIG. 9 shows testis transplanted with mesenchymal stem cells derived from bone marrow. ST shows seminiferous tubule.

[0025] FIG. 10 shows pEGFP-1 vector.

[0026] FIG. 11 shows pPUR vector.

[0027] FIG. 12 shows the charts of flow cytometry for a GFP positive clone.

[0028] FIG. 13 shows immunostaining of P450scc in GFP positive cells.

[0029] FIG. 14 shows the results of RT-PCR of various enzymes in GFP positive (scc+) and negative (scc-) cell fractions.

DETAILED DESCRIPTION OF THE INVENTION

[0030] Mesenchymal stem cells are derived from various interstitial cells and are able to differentiate into various cells. Mesenchymal stem cells are present not only in bone marrow but also interstitial and other tissues (e.g. in kidney, articular membrane, amnion and cord blood) of mesoderm-derived tissues. All these mesenchymal stem cells can be used in the present invention.

[0031] Mesenchymal stem cells derived from bone marrow are interstitial cells, which present in marrow stroma and maintain hematopoiesis, and said stem cells from even adult bear pluripotency of differentiating into various cells and tissues. Mesenchymal stem cells derived from bone marrow may be isolated by inserting an injection needle into an epiphysis for example, by feeding PBS into the epiphysis, by pulling out bone marrow cells, by seeding and subjecting the bone marrow cells to adhere to a plastic dish and by repeating the subculture.

[0032] The transcriptional factor (SF-1), an inducing factor used in the method of the present invention, is an orphan intranuclear receptor, which is expressed in genital and adrenal gland-type steroid hormone-producing cells, and has been known to control the transcription of steroid hormone producing-enzymes (Endocrine Reviews vol. 18, No. 3, 361-377 (1997); The FASEB Journal vol. 10 1569-1577 (1996)). Even if SF-1 is derived from different animal species, SF-1 binds to a common target DNA sequence in mesenchymal stem cells and the factor is expected to provide the same result.

[0033] cAMP exists ubiquitously in all living organism, whose intra-cellular concentration is 10^{-6} to 10^{-7} M. cAMP participates in generation of specific enzymes and metabolic control in target cells and also participates in growth and

differentiation of cells. cAMP is a second messenger of LH and ACTH, which induces the expression of steroid hormone production-related enzymes in genital and adrenal glands and enhances the production of steroid hormones.

[0034] The mean to stimulate by the transcriptional factor (SF-1) and cAMP may include the direct contact of these factors with mesenchymal stem cells or the use of a vector expressing these factors.

[0035] To differentiate mesenchymal stem cells into steroid hormone-producing cells, mesenchymal stem cells may be stimulated by an inducing factor in vitro or said cells may be transplanted into a mammalian reproductive organ. In the case of in vitro induction, for example, human mesenchymal stem cells may be cultured in DMEM (Dulbecco's Modified Eagle's Medium) containing 10% FBS and mouse stem cells may be cultured in IMDM (Iscove's Modified Dulbecco's Medium) or in DMEM containing 10% FBS. The above cells are cultured in an incubator with 5% CO₂ at 37° C. as usual. The concentration of SF-1 is about 0.1~10 µg, preferably at 1 µg/1×10⁵ cells, and that of cAMP is about 0.5~2.0 mM, preferably at 1 mM/1×10⁵ cells.

[0036] When mesenchymal stem cells were transplanted into a mammalian reproductive organ, a certain factor activates the cell (in the case of organ transplantation, interaction with the surrounding cells) to express SF-1, which is considered to make these stem cells to differentiate into steroid hormone-producing cells.

[0037] In the differentiation of mesenchymal stem cells, the role of cAMP is subsidiary. Furthermore, since cAMP is necessarily present in all kind of cells with a large variety of concentrations, intracellular concentration of cAMP might be raised by some reason without adding exogenous cAMP.

[0038] As described above, steroid-producing cells may be obtained accordingly, then steroid hormone may be obtained from the steroid-producing cells.

[0039] Steroid-producing cells derived from mesenchymal stem cells include adrenal cortical cells, ovarian granulosa cells, ovarian capsular cells, testis Leydig cells, testis, Sertoli cells and others.

[0040] Obtained steroid hormones include any steroid hormone and the like, which are synthesized from cholesterol, such as progestin, androgen, estrogen, glucocorticoid and mineralcorticoid.

[0041] The following examples further illustrate the present invention, but it is not intended to limit the scope of the present invention.

[0042] Gene expression of steroid hormone synthesizing enzymes was examined by RT-PCR using the primers shown in Table 1.

TABLE 1

Symbol	Enzyme	Primer	SEQ ID NO:
StAR	steroidogenic acute regulatory protein	F-GAAGGAAAGCCAGCAGGAGAACG	1
		R-CTCTGATGACACCCTCTGCTCC	2
P450scc	cholesterol side chain cleavage enzyme	F-CGCACAGTCCAGAACAACAAGCA	3
		R-CGGTTAGAGAAGGCAGGATAGAG	4
HSD3b	3-β hydroxysteriod dehydrogenase	F-GCAGACCATCCTAGATGTCAAT	5
		R-TCATCATAGCTTTGGTGAGG	6

TABLE 1-continued

Symbol	Enzyme	Primer	SEQ ID NO:
P450c17	17-alpha-hydroxylase/ 17,20 lyase	F-AAAATAATAACACTGGGGAAGGC	7
		R-TGGGTGTGGGTGTAATGAGATGG	8
P450c21	steroid 21-hydroxylase	F-AGAGGATCCGCTTGGGGCTGC	9
		R-GGAGAATTCCTTATGGATGGC	10
P4501 1b1	steroid 11-beta- hydroxylase	F-TCACCAAATGTATCAAGAATGTGT	11
		R-CCATCTGCACATCCTCTTTCTCTT	12
P4501 1b2	aldosterone synthase	F-CCAACAGATGTATCTGGGAGGTGC	13
		R-CCATCTGCACATCCTCTTGCCTCA	14
GAPDH	glyceraldehyde phosphate dehydrogenase	F-ACCACAGTCCATGCCATCAC	15
		R-TCCACCACCTGTGCTGTA	16

EXAMPLE 1

[0043] The coding region (Nucleotides 1~1389 of SEQ ID NO: 17) of rat SF-1 gene (Gene Bank NM_053344, SEQ ID NO:17) was integrated in pIRES-puro2 vector (Clontech, FIG. 1). KUM9 cells (obtained from Department of Reproductive Biology, National Research Institute for Child Health and Development), an established adult mouse-derived mesenchymal stem cell line, were transfected with said vector after linearized with Nru I and were selected in a medium containing puromycin (2 µg/ml) at 48 hrs after the transfection. The resistant colonies remained for 2 weeks in the medium were picked up and were cloned.

[0044] Those clones transfected with parent vector pIRES-puro (FIG. 2B) showed similar morphology to that of KUM9 without transfection (FIG. 2A). In contrast, a large number of fat droplets, conceivably containing cholesterol, i.e. a precursor of steroid hormones, were observed intracellularly in clones (SF-7 and SF-9) expressing SF-1 (FIG. 2C).

[0045] Proteins were extracted from each clone (SF-7 and SF-9), were examined the expression of SF-1 protein by Western blotting and SF-1 protein was detected (FIG. 3).

[0046] Furthermore, gene expression of steroid hormone synthesizing enzymes in these cells was examined by RT-PCR and the expression of P450sc was confirmed.

EXAMPLE 2

[0047] The cloned cells (SF-9), wherein expression of SF-1 was observed in example 1, were cultured for 7 days in the presence of cAMP (1 mM, Sigma). On 1, 3, 5 and 7 days after addition of cAMP, RNA was extracted from said cells and was examined the gene expression by RT-PCR. For comparison, the same procedures were performed for cloned cells using the parent vector pIRES as a negative control. The results show that the expression of p450sc was induced after 1 day, HSD3b1 was induced after 3 days and furthermore p450c17 was induced after 5 days (FIG. 4).

[0048] The result of Western blotting shows that the expression of these enzyme proteins was induced similar to that of mRNA (FIG. 5).

[0049] Moreover, according to these results, those cells without transfection with SF-1 (pIRES-puro) do not express steroid hormone-producing enzymes by the addition of cAMP and do not produce steroid hormone. These facts could be interpreted as that only cAMP could not induce differentiation into steroid hormone-producing cells.

[0050] The interpretation depicts that SF-1 is an essential factor to differentiate mesenchymal stem cells into steroid hormone producing cells and cAMP is a cofactor.

[0051] The output of steroid hormone was measured by radioimmuno assay (RIA) after recovery of culture medium on the 3rd day and 7th day. According to the results, progesterone and androgen were detected on the 3rd day and 7th day, respectively, in parallel to the results of expression of mRNA and proteins (FIG. 6) in the group of cells added with cAMP. Not only the induction of steroid synthesizing enzyme but also the synthesis of androstendione, a kind of androgenic hormone, was confirmed.

[0052] Immunostaining of HSD3b1 shows that all cells are positive (FIG. 7).

[0053] The results of examples 1 and 2 show that forced expression of SF-1 leads mesenchymal stem cells derived from bone marrow to differentiate into steroid hormone-producing cells.

EXAMPLE 3

[0054] Induction of steroid hormone-producing cells for human cells was examined in an experiment similar to example 2 using human derived cells. Human bone marrow mesenchymal stem cells hMSC6/E7 (gifted from Field of Clinical Application, Institute for frontier Medical Sciences, Kyoto University) were transfected with said pIRES-puro-SF-1 vector by calcium phosphate method. At 48 hrs after transfection, cells were selected in a medium containing puromycin (0.3 µg/ml), puromycin resistant colonies were picked up after 2 weeks and they were cloned. After 7 days culture in the presence of cAMP, RNA was isolated and gene expression was examined by RT-PCR.

[0055] hMSC cells stably expressing pIRES-puro-SF-1 did not show morphological change as observed for KUM9 cells (FIG. 2A-C). However, analysis of gene expression by RT-PCR shows the induction of expression of a group of steroid hormone synthesizing genes such as StaR, P450sc and 38-HSD. Furthermore, addition of cAMP induced the expression of P450 c21 and P450 11b1, glucocorticoid producing enzymes in adrenal cortex (FIG. 8).

[0056] These results show that human bone marrow-derived mesenchymal stem cells were differentiated into steroid hormone-producing cells.

EXAMPLE 4

[0057] In this example, bone marrow-derived mesenchymal stem cells were transplanted into testis. Bone marrow

cells were isolated from epiphyseal region of os longum from limbs of 5 w.o. male Green rat by insertion of a 26-gauge needle for syringe and by filling of PBS. After red blood cells were disrupted by osmotic shock, said cells were seeded at $5 \times 10^7/100$ ml dish containing α -MEM medium fortified with 20% FBS. After 24 hrs, the medium was changed to remove non-adherent cells. After 3 days, cells were trypsinized by 0.25% trypsin, subcultured, and cultured for further 1 week. These procedures lead to concentrate mesenchymal stem cells among various bone marrow-derived cells. After confluent cells were trypsinized, 1×10^6 cells were suspended in 50 μ l PBS and were injected into testis of 3 w.o. male Sprague Dawley rat (Japan Charles River Co.).

[0058] After 3 weeks, the testis was removed, observed under inverted microscopy and fixed by 4% para-formaldehyde. Frozen sections were prepared and used for immunohistochemical method. Double staining was performed using rabbit anti-rat P450scc antibody (CHEMICON) and mouse anti-GFP antibody (MBL) for the first antibody and Cy3-conjugated anti-rabbit IgG antibody (SIGMA) and FITC-conjugated anti-mouse IgG (SIGMA) as the second antibody.

[0059] GFP positive cells, possibly derived from Green rat bone marrow, were observed easily under inverted fluorescent microscopy and were detected in testis at 3 weeks after transplantation (FIG. 9A). Therefore, it was found that transplanted cells adhered to host testis. To examine whether these cells were differentiated into steroid hormone producing cells (Leydig cells), double immunostaining using anti-GFP (FIG. 9B) and anti-P450scc (FIG. 9C) was performed. Since most of P450scc positive cells were also GFP positive, Green rat bone marrow-derived cells were found to be differentiated into Leydig cells.

EXAMPLE 5

[0060] Stem cells could not completely be cloned by the method of example 4 because of the contamination of macrophage cells. Therefore, an experiment was performed to examine whether KUM9 cells, an established adult mouse-derived mesenchymal stem cells, has a potency to be differentiated in vitro into steroid hormone-producing cells.

[0061] P450scc (a side chain cleavage enzyme) is an enzyme, which is localized on mitochondrial inner membrane and catalyzes a reaction degrading cholesterol into pregnenolone. Since the above reaction is a necessary procedure in all steroid hormone production, P450scc is expressed in all steroid hormone-producing cells. The group of Chung et al. prepared and used transgenic mice with a visual reporter gene Lac Z and showed that 2.3 Kbp of 5' upstream region in human P450scc gene (Gene Bank M60421) controls the expression of P450scc gene in steroid hormone-producing cells in genital and adrenal gland (Endocrinology, 140:5609 18, 1999). Therefore, preparation of stem cells, wherein the visual reporter gene expresses by the control of the said 5' upstream region, conceivably enables to detect the differentiation into steroid hormone-producing cells.

[0062] Therefore, P450scc gene (Gene Bank M60421, SEQ ID NO: 18) was integrated into the upstream of EGFP protein coding region of pEGFP-1 vector (FIG. 10, Clontech) and the vector was fused with pPUR vector (FIG. 11, Clontech) coding puromycin resistant gene.

[0063] Said synthetic vector, after linearized with ApaI, was transfected into KUM9 cells (Gifted from Department of Reproductive Biology, National Research Institute for Child Health and Development), derived from mouse bone marrow mesenchymal stem cells by lipofection. Since fluorescent GFP expresses at the time of expression of P450scc in these cells, spontaneously differentiated KUM9-derived steroid-producing cells could be separated by flow cytometer using fluorescence as a probe.

[0064] At 48 hrs after the separation, cells were selected in the medium containing puromycin (2 μ g/ml) and resistant colonies after 2 weeks was cloned by a limiting dilution method.

[0065] GFP positive cells were detected in 2 clones out of isolated 250 clones during growing phase of these cells under inverted fluorescence microscopic observation. GFP positive and negative cells were separated by flow cytometer for said 2 clones, wherein a lot of GFP positive cells were found, and GFP positive cells were 2~5% (FIG. 12).

[0066] Furthermore, immunostaining of P450scc for GFP positive cells confirmed that GFP positive cells were also P450scc positive (FIG. 13).

[0067] Moreover, GFP positive (scc+) and negative (scc-) cells were separated by a cell sorter, total RNA was extracted from the separated cells and cDNA was prepared by reverse transcription reaction using reverse transcriptase. RT-PCR was performed using a specific primer for each gene. The expression of mRNA of P450scc gene was observed in GFP positive cells (scc+) (FIG. 14). Additionally, the transcriptional factor SF-1 was expressed in said cells isolated. Still furthermore, examination of the expression of genes involved in steroid hormone production revealed that the expression of 3 β -HSD gene of both genital and adrenal gland type (HSD3b1) and placental type (HSD3b6) was observed.

[0068] From these results, it was interpreted that the transplantation induced the expression of SF-1 in stem cells, and that these stem cells differentiated into steroid hormone-producing cells on contact with SF-1. a transcriptional factor SF-1 was expressed (Example 5). Therefore, the inventors prepared a KUM9-derived strain, wherein SF-1 is expressed constitutively, and added a stimulation by cAMP, then the inventor found that the KUM9-derived cells stably produced steroid hormone (Examples 1 and 2). Moreover, it was confirmed that not only a gonadal steroid hormone-producing enzymes but also adrenal steroid hormone-producing enzymes are induced in a human cell line (Example 3). It is expected that these results are fundamental data in using differentiation of stem cells for medical treatment of human steroid hormone deficiency and provide a useful method for regeneration treatment.

[0069] From the above results, the inventors discovered that stimulation of mesenchymal stem cells by a transcriptional factor (SF-1), preferably with SF-1 and cAMP, enabled to differentiate the mesenchymal stem cells into steroid-producing cells and accomplished the present invention.

[0070] Namely, the present invention is a method for differentiating mesenchymal stem cells into steroid hormone-producing cells, comprising stimulating the mesenchymal stem cells by a transcriptional factor (SF-1). Furthermore, said method may comprise further stimulating the mesenchymal stem cells by cAMP.

[0071] The mesenchymal stem cells are preferably derived from bone marrow and said mesenchymal stem cells are preferably human cells.

[0072] Still furthermore, the present invention is a method for producing steroid-producing cells, comprising producing steroid producing cells by implementing said method in vitro or by transplanting the mesenchymal stem cells into a mammalian reproductive organ.

[0073] Moreover the present invention is steroid-producing cells or non-human animals containing said steroid-producing cells obtained by these method.

ADVANTAGES OF THE PRESENT INVENTION

[0074] The present invention provides, for the first time, a method for differentiating mesenchymal stem cells into steroid hormone-producing cells.

[0075] It will be possible to obtain a large amount of differentiated cells by transplanting stem cells into a genital gland of a non-human animal such as swine, if immune difficulties were overcome. The most important issue in regeneration medicine and

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 18

<210> SEQ ID NO 1

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 1

gaaggaaagc cagcaggaga acg

23

<210> SEQ ID NO 2

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 2

ctctgatgac accactctgc tcc

23

<210> SEQ ID NO 3

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 3

cgcacagtcc agaacaacaa gca

23

<210> SEQ ID NO 4

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 4

cggttagaga aggcaggata gag

23

<210> SEQ ID NO 5

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 5
gcagaccatc ctagatgtca at 22

<210> SEQ ID NO 6
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 6
tcacatagc tttggtgagg 20

<210> SEQ ID NO 7
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 7
aaaataataa cactggggaa ggc 23

<210> SEQ ID NO 8
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 8
tgggtgtggg tgtaatgaga tgg 23

<210> SEQ ID NO 9
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 9
agaggatccg cttggggctg c 21

<210> SEQ ID NO 10
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 10
ggagaattcc ttatggatgg c 21

<210> SEQ ID NO 11
<211> LENGTH: 24

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 11

tcaccaaagt tatcaagaat gtgt 24

<210> SEQ ID NO 12
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 12

ccatctgcac atcctctttc tctt 24

<210> SEQ ID NO 13
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 13

ccaacagatg tatctggaag gtgc 24

<210> SEQ ID NO 14
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 14

ccatctgcac atcctcttgc ctca 24

<210> SEQ ID NO 15
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 15

accacagtcc atgccatcac 20

<210> SEQ ID NO 16
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 16

tccaccacc tgttgtgta 20

- continued

```

<210> SEQ ID NO 17
<211> LENGTH: 1389
<212> TYPE: DNA
<213> ORGANISM: Rattus norvegicus

<400> SEQUENCE: 17
atggactatt cgtacgacga ggacctggac gagctgtgtc cagtgtgtgg tgacaagggtg    60
tcgggctacc actacgggct gctcacgtgc gagagctgca agggcttctt caagcgcaca    120
gtccagaaca acaagcatta cacgtgcacc gagagtcaga gctgcaaaat cgacaagacg    180
cagcgtaaag gctgtccctt ctgccgcttc cagaagtgcc tgacgggtggg catgctgcctg    240
gaagctgtgc gtgctgatcg aatgcggggc gcccggaaca agtttggggc catgtacaag    300
agagaccggg ccttgaagca gcagaagaaa gcacagattc gggccaatgg cttcaaactg    360
gagaccggac caccgatggg ggttcccccg ccacccccctc cccacccgga ctacatgtta    420
ccccctagcc tgcattgcacc ggagcccaag gccctgggtct ctggcccacc cagtggggccg    480
ctgggtgact ttggagcccc atctctgccc atggccgtgc ctggctccca cgggcctctg    540
gctggctacc tctatctgct cttctctaac cgcaccatca agtctgagta tccagagccc    600
tacgccagcc cccctcaaca gccaggggcca ccctacagct atccggagcc cttctcagga    660
gggccaatg taccagagct catattgcag ctgctgcaac tagagccaga ggaggaccag    720
gtgctgtgct gcatcgtggg ctgcctgcag gagccagcca aaagccgcc tgaccagcca    780
gcgcccttca gcctcctctg caggatggcg gaccagacct ttatctccat tgtcgactgg    840
gcacgaaggt gcatgggtatt taaggagctg gaggtggctg accagatgac actgctgcag    900
aactgctgga gtgagctgct ggtgctggac cacatctacc gccaggtcca gtacggcaag    960
gaagacagca tcttctgctg cactggacag gaggtggagc tgagcacggt ggctgtgcag    1020
gctggctccc tgctgcacag cctggtgctg cgggcacagg agttggtgct gcagctgcat    1080
gccctgcaac tggaccgcca ggagtttgtc tgtctcaagt tccctatcct cttcagcctc    1140
gatgtgaaat tccctaacaa ccacagcctg gtaaaggacg cccaggagaa ggccaacgcc    1200
gccctgctgg attacacctt gtgtcactac ccacactgct gggacaaatt ccagcagttg    1260
ctattgtgct tggtggaggt gcggggcactg agcatgcagg ccaaggagta tctgtacctat    1320
aagcatttgg gcaacagatg gcccccgaac aaccttctca ttgagatgct gcaggccaag    1380
cagacttga                                     1389

```

```

<210> SEQ ID NO 18
<211> LENGTH: 2393
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 18
aagcttcagg gatggcagcc gcttgtgaga aacctgagc atgagccact cagccaacca    60
gccaccagct caccagacca cccagccacc cagtcacca gaaaagtgc tcctgggtgc    120
tgcaccctcg gaagctgtga gataataaac atttattgtt ttaagccact aaatthtggg    180
ataatthtgt aagcagcagt aaacagctaa tacattcagc cttgtttgga gtgagtgatg    240
tgthttctgga agctctttca gagaagtgag ggagctattc tcccagaagc cacagcaaac    300
ctttccctgt gtttcattgg cccaaaactg atcggctggc ctatgctgtg atgtgacctat    360
ggcgattgga gaggatgagg caataacctc cagcctgggc cacttctggg gagggcggtca    420

```


- continued

gtgcccacaa cactggggga ggtgcgagg cctgaacgga agttggggtg gctgccaaga	480
ggaccacaag ttcttccatg ccacatcgat tagggctcct tctgaggag gaatgtgggg	540
ctgcgtagaa caatgggatt gactttaagt cagaaagta taaatgtcac ctcagtgctg	600
agacccttgg aggaaaaact agtccttga agactgcttt tottgtggaa gctcatcacc	660
ctgccgctgc tcgtgagaca ctgccttcct tggctgatgt cattocaggc tcaaggtcat	720
catggaggca aaacaggctt tctcactact tctttatcag aaggttcatg actgatgagg	780
tagtgggtcac tccagcggga agagcaacaa ccaactctga taagtacttt tttttttct	840
tctaaaaact gttgctctaa atttgttga agtggttcaa cagtattgga gtctggggtc	900
aagtggctgt gtaaggacaa cttttgcat tgtgggtgat ctatgggctg acacatacaa	960
cagaagaggc caggaggatg tcactcgtgt gtgtgtgtgt gtgtgtgtgt gtgttggtaa	1020
cagctctatt gagatataat tcacacacca tacattcacc catttaaagt atgcaattca	1080
atggctttta gtgtatagag agttgttcaa taattaccac atatcttag aatattttca	1140
ttttcatcat cctaagaatc cctacacact ttatgtccca tcctctaatt cctctatccc	1200
cctagctcta agcaaccacc agtctatfff ctgtctgta gatgtgtgtt ttaaagtcc	1260
tatttgccag aaaataagat tttggggaaa aaaagcaaga agtacctcc aatgataagt	1320
ataaggata gtgtagattg ctgtcatgaa atgctctaca tggcacatgt atgttttctg	1380
acctacacat ttttccaag tgggctgtaa ttttgttga atgacttga aactctacag	1440
gtgactgacc cttgttgttt gggaaagtgg taggtgcagg gtgatggggg gtgggtgtgg	1500
gggctgacct gggctggaag gatgggtctg gggatatgat atgatgggag ggggatgga	1560
agcagctctg aggggtttgc agcacagatc aagtgtgtgg acaggaaga gctgacatcc	1620
tgagttccgg atggcaacca gatttgccaa ggtcttagag tgtgtccaga gtggagcctg	1680
accacagacc tcagctcaag ggaccagag cccctctgag tcagctgtac tgaattacag	1740
ccccaaatct gggcaactg gggagagacg acgaggatta gggttccaag gtgaaactgt	1800
gccattgcgc tccagcctgg gcaacaagaa tgaaactctc taaaaataa ataaaaataa	1860
atagcctaag gatgcatttc tcagaactta tccctgtgt tcaatgatgt gtgtctatac	1920
agtggggcca taactaagac gtatgttgc caagctgca agatagctct gacctctct	1980
tgggcccctc atttccccca aacacagggt gtctgcagtc ttgaccaatg gctgccaggg	2040
catggactcc gctgcagggg ccagtgagg gcccagctc aggcataaagc acaggcagat	2100
atctcaggag tctgctaggg ctggcactga gggcagagac agaggggtct ccctgtcctt	2160
tggagaacct cacgctcag aaattccaga ctgaaccttg ataccgagta ggggaggagc	2220
tgtctgctgg tttgagcctg cagcaggagg aaggacgtga acattttatc agcttctggt	2280
atggccttga gctggtagtt ataactctgg ccctggggc ccagggtctac agtcatccta	2340
gcagtccccg ctgaagtgga gcaggctacag tcacagctgt ggggacagca atg	2393

1. A method for differentiating mesenchymal stem cells into steroid hormone-producing cells, comprising stimulating the mesenchymal stem cells by a transcriptional factor (SF-1).

2. A method of claim 1 comprising further stimulating the mesenchymal stem cells by cAMP.

3. A method of claim 1, wherein the mesenchymal stem cells are derived from bone marrow.

4. A method of claim 3, wherein the mesenchymal cells are derived from human.

5. A method for producing steroid hormone-producing cells, comprising implementing the method of claim 1 in vitro.

6. A method for producing steroid hormone-producing cells, comprising implementing the method of claim 1 by transplanting the mesenchymal stem cells into a mammalian reproductive organ.

7. (canceled)

8. A method for producing steroid hormone, comprising culturing the steroid hormone-producing cells obtained by the method of claim 5 and recovering steroid hormone from the culture medium.

9. The method of claim 8, wherein said steroid hormones are progestin, androgen, estrogen, glucocorticoid or mineral-corticoid.

10. Non-human animals carrying the steroid-producing cells obtained by the method of claim 5.

* * * * *