



(11)

EP 1 584 678 B1

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention of the grant of the patent:
31.08.2016 Bulletin 2016/35

(21) Application number: **03781013.2**

(22) Date of filing: **22.12.2003**

(51) Int Cl.:
C12N 15/00 (2006.01)

(86) International application number:
PCT/JP2003/016496

(87) International publication number:
WO 2004/058964 (15.07.2004 Gazette 2004/29)

(54) **METHOD OF INDUCING HOMOLOGOUS RECOMBINATION OF SOMATIC CELL**
INDUKTIONSVERFAHREN FÜR HOMOLOGE REKOMBINATION EINER KÖRPERZELLE
METHODE D'INDUCTION DE RECOMBINAISON HOMOLOGUE D'UNE CELLULE SOMATIQUE

(84) Designated Contracting States:
CH DE FR GB LI

(30) Priority: **26.12.2002 JP 2002376555**

(43) Date of publication of application:
12.10.2005 Bulletin 2005/41

(73) Proprietors:
• **RIKEN**
Wako-shi, Saitama 351-0198 (JP)
• **Chiome Bioscience, Inc.**
Shibuya-ku
Tokyo 151-0071 (JP)

(72) Inventors:
• **OHTA, Kunihiro**
Wako-shi, Saitama 351-0198 (JP)
• **SEO, Hidetaka**
Wako-shi, Saitama 351-0198 (JP)
• **SHIBATA, Takehiko**
Wako-shi, Saitama 351-0198 (JP)

(74) Representative: **Ahner, Philippe et al**
BREVALEX
95, rue d'Amsterdam
75378 Paris Cedex 8 (FR)

(56) References cited:
• **CHAUHAN S S ET AL: "CONSTRUCTION OF A NEW UNIVERSAL VECTOR FOR INSERTIONAL MUTAGENESIS BY HOMOLOGOUS RECOMBINATION" GENE, ELSEVIER, AMSTERDAM, NL, vol. 120, 1992, pages 281-285, XP002911851 ISSN: 0378-1119**

- **NICKOLOFF J A: 'Transcription enhances intrachromosomal homologous recombination in mammalian cells' MOL. CELL BIOL. vol. 12, no. 12, 1992, pages 5311 - 5318, XP008018567**
- **BUERSTEDDE J, TAKEDA S: 'Increased ratio of targeted to random integration after transfection of chicken b cell lines' CELL vol. 67, no. 1, 1991, pages 179 - 188, XP002976450**
- **CHEN J, ET AL: 'Mutations of the intronic igh enhancer and its flanking sequences differentially affect accessibility of the jh locus' EMBO J vol. 12, no. 12, 1993, pages 4635 - 4645, XP002976451**
- **FORRESTER W C, ET AL: 'Dependence of enhancer-mediated transcription of immunoglobulin mu gene on nuclear matrix attachments regions' SCIENCE vol. 265, 1994, pages 1221 - 1225, XP002976452**
- **BACHL J, WABL M: 'Enhancers of hypermutation' IMMUNOGENETICS vol. 45, no. 1, 1996, pages 59 - 64, XP002962724**
- **BETZ A G, ET AL: 'Element regulation somatic hypermutation of an immunoglobulin kappa gene: critical role for the intron enhancer/matrix attachment region' CELL vol. 77, no. 2, 1994, pages 239 - 248, XP002976453**
- **BACHL J, WABL M: 'An immunoglobulin mutator targets g.c base pairs' PROC. NATL. ACAD. SCI. USA vol. 93, 1996, pages 851 - 855, XP002976454**
- **LAHTI J M: 'Use of gene knockouts in cultured cells to study apoptosis' METHODS vol. 17, no. 4, 1999, pages 305 - 312, XP004466980**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 584 678 B1

- SLEBOS R, TAYLOR J A: 'A novel host cell reactivation assay to assess homologous recombination capacity in human cancer cell lines' **BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS** vol. 281, no. 1, 2001, pages 212 - 219, XP002227693
- ARAKAWA H, ET AL: 'Requirement of the activation-induced deaminase (aid) gene' **SCIENCE** vol. 295, 2002, pages 1301 - 1306, XP002976455
- J A NICKOLOFF ET AL: 'TRANSCRIPTION STIMULATES HOMOLOGOUS RECOMBINATION IN MAMMALIAN CELLS' **MOL. CELL. BIOL.**, [Online] vol. 10, no. 9, 01 September 1990, pages 4837 - 4845, XP055086555 DOI: 10.1128/MCB.10.9.4837 Retrieved from the Internet:
<URL:<http://mcb.asm.org/content/10/9/4837.full.pdf#page=1&view=FitH>> [retrieved on 2013-11-04]

Description

Technical Area

[0001] The present invention generally relates to an art for inducing somatic cell homologous recombination in a cell of a eukaryotic organism, and more specifically concerns a method for inducing somatic cell homologous recombination at an arbitrary genetic locus in a somatic cell, and a cell wherein somatic cell homologous recombination is induced by said method.

[0002] Additionally, the present invention also discloses a method for producing a novel gene having a plurality of genetic information, by using the aforementioned somatic cell homologous recombination induction method.

[0003] Further, the present invention discloses a novel protein encoded by the abovementioned novel gene.

[0004] Still further, the present invention discloses a vector used for, and having been constructed in a structure optimal for, inducing somatic cell homologous recombination.

Background Art

[0005] For eukaryotic organisms, the homologous recombination of somatic cells is one of the most important DNA metabolic reactions for obtaining a diversity of genes, and as a result, for producing diverse protein activity. Therefore, the control of homologous recombination of somatic cells is one of the extremely important problems for obtaining a diversity of genes.

[0006] Conventionally, as a method for obtaining a diversity of genes having new functions or attributes, there is the art of DNA shuffling (see, e.g., Crameri et al., 1998). This art simulates homologous recombination by mixing a plurality of homologous genetic sequences, appropriately digesting with DNaseI and using the resulting small fragments as primer, and conducting PCR with the original gene as a template. However, the analysis of recombination products is generally done after amplified fragments are linked to expression vectors, and transformed into bacteria, and it is thought that the direct analysis of what characteristics the products would have in higher eukaryotic organisms would be difficult. Additionally, checking expression within an animal cell requires that expression compatibility due to transfer into a new vector and codon usage be confirmed successively.

[0007] On the other hand, as systems that induce genetic recombination within animal cells, there are systems that activate recombination within cells, such as systems using the site specific recombination enzyme Cre-lox (see, e.g., DiSant et al., 1995), and systems using the sequence specific endonuclease I-SceI (see, e.g., Rouet et al., 1994). The Cre-lox system uses Cre, a 38 kDa site specific recombinase obtained from the bacteriophage P1, and induces recombination between specific sites called loxP sites. Additionally, the I-SceI system utilizes the activity of I-SceI, an endonuclease derived

from budding yeast, of cleaving a DNA double strand at a recognition site thereof, and inducing DNA homologous recombination. However, in these systems, because recombination is restricted to being between specific sequences, and the recombination event is one-shot, the recombinants that could be obtained were in principle only one type. Additionally, since it was necessary to introduce a sequence relating to the control of the recombination into a chromosome beforehand to cause recombination enzymes such as Cre or I-SceI to be expressed within a cell, the induction of recombination of chromosomes was not easy.

[0008] In 1992, Chauhan and Gottesman disclose homologous recombination between a targeting gene on a pSSC-9 vector and a chromosomal gene. In addition, the targeting gene inserted on the pSSC-9 vector at a restriction site flanking the neo gene is not under the control of a promoter.

[0009] Nickoloff and Reynolds have shown in 1990 that transcription stimulate both reciprocal and non reciprocal events between duplicated regions in yeast. The influence of transcription on homologous intrachromosomal recombination between direct and inverted repeats (Neo wt gene and Neo mt gene) using CHO cells is studied by Nickoloff, 1992. In the gene constructs implemented therein, the transcription promoter (MMTV) is located 5' to the Neo mt gene and the Neo wt gene is located 3' to the Neo mt gene and also 3' to MMTV.

[0010] Lahti proposes to use the avian DT40 cell system which undergoes homologous recombination at very high frequencies as a novel method to generate loss of function mutations in vertebrate cells.

Disclosure of the Invention

[0011] The inventors of the present invention, having considered the abovementioned state of affairs, conducted penetrating research into whether or not a method exists for inducing somatic cell homologous recombination under controlled conditions, and as a result, found to their surprise that by controlling transcription at genetic loci where recombination is desired, it is possible to induce somatic cell homologous recombination between similar base sequences.

[0012] Accordingly, in general, the present invention has the objective of providing a method for inducing somatic cell homologous recombination at a genetic locus in a somatic cell, as defined in claims 1-9.

[0013] Additionally, the present invention discloses a cell wherein somatic cell homologous recombination is induced by the abovementioned method.

[0014] Further, the present invention discloses a novel gene having a plurality of genetic information and a method for producing said novel gene, by using somatic cell homologous recombination induced within a cell.

[0015] Still further, the present invention discloses a novel protein encoded by said novel gene.

[0016] Additionally, the present invention discloses a

vector with a structure constructed in order to desirably control transcription and to be used to induce somatic cell homologous recombination.

[0017] Further, the present invention provides an induction method for somatic cell homologous recombination characterized by somatic cell homologous recombination of a eukaryotic organism cell wherein DNA homologous recombination can occur at an arbitrary genetic locus, whereby, by controlling the transcription of a gene at said genetic locus, DNA homologous recombination is induced between said gene and a similar base sequence. Additionally, a cell is also disclosed wherein somatic cell homologous recombination is induced by said method.

[0018] Additionally, the present invention discloses a method for obtaining a novel gene having a plurality of genetic information, by somatic cell homologous recombination of a eukaryotic organism cell wherein DNA homologous recombination can occur at an arbitrary genetic locus, whereby, by controlling the transcription of a gene at said genetic locus, DNA homologous recombination is induced between said gene and a similar base sequence. Additionally, a novel gene produced by said method is also disclosed.

[0019] Still further, a novel protein encoded by said novel gene is also disclosed.

[0020] Additionally, the present invention discloses a constructed vector with the structure necessary to desirably control transcription, used in order to induce somatic cell homologous recombination.

[0021] By using the art disclosed in the specification of the present invention, since DNA homologous recombination is induced continuously at various locations, it is possible to increase diversity of the obtained recombinants. Additionally, by using the art disclosed in the present specification, since the objective genes having genetic diversity are spontaneously formed within objective cells, without producing a DNA library for recombination required for DNA shuffling technology, screening can be done within said cells for the production of the desired novel gene. Therefore, a diversity of novel genes is obtainable in a time shorter than for the conventional art.

[0022] For the eukaryotic organism cells, it is thought that any cell wherein somatic cell homologous recombination may occur at an arbitrary genetic locus is usable, but chicken derived DT40 cultured cells is used.

[0023] A gene for which somatic cell homologous recombination is induced is an exogenous gene. Additionally, among base sequences similar to said gene are included, for example, genetic sequences that exist in nature, base sequences that exist in nature which do not have the form of a gene, and man-made sequences.

[0024] In the case of an exogenous gene, any gene is utilizable, as long as it is constructed on an appropriate vector with a structure including a transcription promoter upstream of a gene that produces homologous recombination, and a sequence in the vicinity and further up-

stream similar to said gene.

[0025] Here, it is sufficient for the base sequences similar to the objective gene to be located in the region approximately 1 bps to 100 kbps upstream from the transcription initiation location of the objective gene, but this should more preferably be located in a region approximately 500 bps to 50 kbps upstream, and even more preferably approximately 3 kbps to 25 kbps upstream.

[0026] As an exogenous gene, there are no restrictions, but fluorescent protein genes (e.g., GFP genes, CFP genes and the like), drug resistance genes (e.g., hygromycin resistance gene, or puromycin resistance gene) and the like are included.

[0027] The "base sequence similar to a gene for which somatic cell homologous recombination is induced" used in the present invention is a base sequence that is similar to the entirety of said gene. Here, similar means having 60% or greater sequence identity, preferably 70% or greater sequence identity, and more preferably 80% to 99%, including 80%, 82%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, and 99% sequence identity with the entirety of a gene for which somatic cell homologous recombination is induced.

[0028] For the transcription control method used in the present invention, any method well-known to those skilled in the art is usable, but a method using an endogenous transcription promoter, or a method of control using an endogenous inducible transcription promoter is preferable. Additionally, in order to promote transcription activity, an enhancer and a nuclear matrix attachment region (MAR) appropriate for the objective gene may be used. In order to control transcription, it is necessary to align the transcription promoter, the enhancer, and the nuclear matrix attachment region (MAR) in an action-capable manner. Here, "action-capable" means that the transcription promoter, enhancer, and nuclear matrix attachment region (MAR) can realize each of their functions, and that the transcription of the objective gene is controlled as desired. For example, in the case of a transcription promoter, there are no particular restrictions, but it should be located in a region within approximately 100 bps upstream of the transcription initiation location of the objective gene, and in the case of the enhancer, there are no restrictions, but normally, it may exist 100 bps or more, or several thousand bps or more from the transcription initiation point, and may be on the 5' side or the 3' side of the objective gene. Additionally, the nuclear matrix attachment region (MAR) is located adjacent to the enhancer, and may be on the 3' side or the 5' side of the enhancer.

Brief Explanation of the Drawings

[0029]

Figure 1 shows an outline of a method for constructing on a vector a cis-acting region necessary for in-

ducing transcription.

Figure 2 shows the change over time of fluorescent intensity under FACS. Gate E is thought to be ECFP, and a group with even stronger fluorescent intensity was gated to F.

Figure 3 shows the change over time of fluorescent intensity under FACS. Quantification was done by taking the ratio of the fluorescent intensity of E and the fluorescent intensity of F.

Figure 4 shows an observed image of cells wherein homologous recombination was induced between the ECFP gene and the EGFP gene under a fluorescent microscope.

[0030] This is a fluorescent microscope image 5 days after inducing transcription of ECFP. Among cells (A) wherein CFP fluoresces, cells (B) were observed (C) wherein GFP is thought to fluoresce.

[0031] Figure 5 shows analysis results of a novel genetic sequence. As a result of analysis of a sequence directly under TRE of a cell group with a strong fluorescent intensity, four novel types of recombinants, type1, type2, type3 and type4 were found. Further, ECFP and EGFP sequences are shown above and below the four types of sequences. Further, other than ECFP, only sequences differing for EGFP and ECFP are noted.

Best Modes for Embodying the Invention

[0032] Since the method of homologous recombination of the Enhanced Cyan Fluorescent Protein (ECFP) gene utilizes in part the induction method for somatic cell homologous recombination according to the present invention, herebelow, the method of homologous recombination of the enhanced cyan fluorescent protein (ECFP) gene shall be explained in detail.

[0033] As mentioned above, in the homologous recombination method for enhanced cyan fluorescent protein (ECFP) genes of the present invention, cells wherein DNA homologous recombination is occurring are selected and cultivated, and when inducing homologous recombination, DNA homologous recombination of said cells is regulated by an inducible promoter, to induce homologous recombination with enhanced green fluorescent protein (EGFP) gene sequences having a base sequence similar to ECFP.

[0034] Therefore, herebelow, selection and cultivation of cells, construction of a vector for induced transcription control, confirmation of expression of a novel gene wherein homologous recombination is induced, and confirmation of a novel gene sequence wherein homologous recombination is induced shall be explained in order.

1. Selection of cells wherein somatic cell homologous recombination is induced

[0035] "Eukaryotic cells" in the present invention refer to cells wherein somatic cell homologous recombination occurs with some frequency, preferably being human, mouse, sheep, rat, rabbit, or chicken B cells or cell lines thereof, more preferably being human Burkitt's lymphoma derived Ramos cell lines, or chicken B cell derived DT40 cell lines. The eukaryotic cells used in the present invention are DT40 cell lines.

[0036] The "DT40 cell lines" used in the present invention include derived cell lines and sublines wherein the chromosomes of said cells have been modified in some manner (e.g., recombination, insertion, deletion, and the like of a specific gene).

[0037] The cultivation conditions used for the present invention are conducted in a manner publicly known in said technical field, but needless to say, this shall be performed in a medium and cultivation conditions (cultivation temperature, CO₂ concentration) appropriate for the selected cells. As a medium, the generally used RPMI1640 medium or DMEM medium, or media wherein fetal bovine serum has been added to these media, and the like, may be given as examples. Cultivation is normally done in the presence of 5% CO₂, for 1 to 30 days at the appropriate temperature (e.g., 25 degrees Celsius to 40 degrees Celsius) for the utilized cells. During cultivation, antibiotics such as kanamycin and penicillin may be added to the medium as needed during cultivation. Additionally, in cases where transcription control is done using an inducible promoter, a drug for inducing the transcription may be added (e.g., tetracycline). Thus, in cases where the selected immune cells are DT40 cells, for example, IMDM (Invitrogen) is used as the medium, the cultivation temperature is, for example, 39.5 degrees Celsius, and cultivation is done in a 5% CO₂ concentration.

2. Construction of a vector for transcription control (1)
Cis-acting region

[0038] As the method for transcription control used in the present invention, any method may be used as long as it is a well-known art for those skilled in the art, but preferably a method is used wherein a cis-acting region for transcription control is constructed on an appropriate vector, and by introducing said vector into a cell, incorporating it into a specific genetic locus on the chromosome. Here, the "cis-acting region" refers to a DNA sequence necessary for the transcription control of a specific gene, being a region that exists adjacent to said gene. A "cis-acting region" includes transcription promoters, enhancers, nuclear matrix attachment regions (MAR), and other arbitrary sequences that are necessary for controlling transcription activity.

[0039] For the transcription promoter, either constitutive promoters or inducible promoters are usable, but from the fact that transcription control is easy, inducible

promoters are preferable. Inducible promoters include promoters that respond to α -interferon, heat shock, heavy metal ions, steroids such as glucocorticoids (Kaufman, 1990), and tetracycline. Other desirable inducible promoters include those which respond within a cell when the activation factor for a promoter is added from the outside. Preferably, for example, a tetracycline inducible promoter or the like is usable.

[0040] Any enhancer and nuclear matrix attachment region (MAR) is usable, as long as they function for the control of transcription activity within selected cells, but in cases where DT40 cells are used, it is preferable to use a chicken antibody light chain gene enhancer (3' enhancer) and a chicken-derived nuclear matrix attachment region.

[0041] Among the components constituting a cis-acting region, in cases where the sequences of the inducible promoter, enhancer, and nuclear matrix attachment region (MAR) are publicly known, it is possible to produce a primer in order to amplify said sequence by the PCR method or the like based on this sequence, and to obtain them directly from appropriate cDNA libraries or genome libraries. It is preferable for the primer used here to be designed so that, for convenience when cloning, a restriction enzyme site that exists on the vector for cloning is added. The obtained enhancer and nuclear matrix attachment region (MAR) may be cloned using an appropriately added restriction enzyme site, on a vector whereon are constructed a gene for which homologous recombination is to be induced, and a DNA sequence similar to said gene, in the order of a promoter between said gene and a similar DNA sequence, a nuclear matrix attachment region (MAR), and an enhancer, in the 3' direction downstream from said gene, or in the opposite order.

[0042] Additionally, the target gene, base sequence similar to said gene, enhancer, nuclear matrix attachment region (MAR), and the like may be cloned onto commercially available vectors, for example, pTRE2hyg (Clontech) on which inducible promoters have been pre-cloned.

(2) Obtaining the gene (ECFP)

[0043] The gene for which homologous recombination is to be induced may be an endogenous gene, but it may also be an isolated exogenous gene. Because of ease of transcription control, exogenous genes are preferable.

[0044] The genes for which homologous recombination is to be induced are cloned by methods well-known to those skilled in the art, but said genes may be amplified by using a primer set (e.g., a primer pair set designed so that said gene is amplifiable on the 5' end and the 3' end of said gene) usable for amplifying said gene, and a DNA polymerase with a highly accurate DNA polymerase (e.g., Pyrobest DNA Polymerase (TaKaRa)).

[0045] The obtained gene can be cloned into an appropriate cloning vector (e.g., pBlueScriptII, pUC19 and

the like).

[0046] In cases where the gene for which homologous recombination is induced is an ECFP gene, cloning may be done with a commercially available vector (pECFP-C1).

[0047] By a similar method, a base sequence similar to the entirety or a portion of the ECFP gene is obtainable. Any sequence similar to the ECFP gene is usable, but, for example, the EGFP gene sequence and the like are preferable.

[0048] As mentioned above, the obtained ECFP gene and the sequence similar to the ECFP gene are inserted in an action-capable manner on a vector whereupon a "cis-acting region" is constructed, on the upstream side of the ECFP gene (that is, the 5' side), with a transcription promoter in between. The direction in which the similar DNA sequence is inserted may be in the ordinal direction or the opposite direction. The "ordinal direction" refers to the naturally existing direction if the objective DNA sequence occurs naturally, and in cases where the sequence is man-made, either direction may be selected to be the "ordinal direction."

3. Construction of a Transcription Factor Expression Vector

[0049] In order to induce the somatic cell homologous recombination of a gene, the expression of a transcription factor necessary for the promotion or suppression of transcription promoter activity is necessary. For example, in the case of a tetracycline induced promoter, a transcription factor (e.g., a VP16 transcription activation domain fused to a Tet repressor) that regulates transcription by binding to a tetracycline response region. Vectors for expressing these transcription factors may include, for example, promoters linked action-capably relative to the expression of said transcription factors (e.g., human cyto megalovirus immediate early promoter), and appropriate selective markers (e.g., a neomycin resistance marker, a hygromycin resistance marker, and the like).

[0050] Additionally, as vectors for controlling promoters, commercially available vectors, such as pTet-Off or pTet-On vector (Clontech) may be used.

4. Confirming expression of a novel gene for which homologous recombination has been induced

(1) Induction of homologous recombination of a selected gene

[0051] In order to induce homologous recombination of a selected gene (here, an ECFP gene), a factor that further regulates the activity of a transcription factor for a promoter action-capably linked to an ECFP gene is needed. Said factor is determined depending on the transcription promoter selected, and this can be selected by those skilled in the art without an excessive amount of experiments and the like. For example, in cases where

the selected promoter is a tetracycline inducible promoter, tetracycline or doxycycline and the like are usable as said factor.

[0052] Additionally, it is necessary to introduce a vector for transcription control and a transcription factor expression vector into the selected cells. As a method for introducing the vectors into the cells, any method well-known to those skilled in the art may be used, but for example, the calcium phosphate method (Chen and Okayama, 1988), or the cationic lipid method (Elroy-Stein and Moss, 1990), the electroporation method (Neumann et al., 1982), and the like may be used.

[0053] After having introduced the aforementioned two types of vectors into the cells, the objective transcription promoter is activated by adding or removing the factor necessary for activation of the transcription promoter (e.g., tetracycline, doxycycline, and the like) to the medium. By doing so, homologous recombination of the selected gene is induced.

(2) Confirming Expression of a Novel Gene for which Homologous Recombination is Induced

[0054] The expression of a novel gene that is created by the induction of homologous recombination may be confirmed by using methods well-known to those skilled in the art, according to the characteristics of the genetic products prior to the induction of recombination. For example, in cases where the selected gene has a specific drug resistance, by comparing, for cells prior to the induction of homologous recombination, and cells after induction, tolerable concentrations and the like of a specific drug, the expression of a novel gene can be predicted. Additionally, in the case of ECFP genes, the expression of a novel gene may be confirmed by changes in the wavelength of the fluorescent light emitted, and the like.

5. Sequence analysis of a novel gene induced by homologous recombination

[0055] The creation of a novel gene can further be confirmed by conducting sequence analysis on the novel gene for which homologous recombination was done, and by the actual occurrence of homologous recombination.

[0056] The genomic DNA of a cell whereon confirmation of homologous recombination is to be done is extracted by a known method, and the objective gene is amplified by the PCR method using specific DNA primers (e.g., in order to include the entire objective genetic region, design a primer in the forward direction in the vicinity of the 5' side of said gene, and design a primer in the reverse direction on the 3' side of said gene) usable for the amplification of the gene to be confirmed.

[0057] A commercially available DNA polymerase may be used for amplification, but it is preferable to use one that is highly accurate. The conditions for conducting amplification depend on the annealing temperature of the

DNA primer used, and the properties of the DNA polymerase used, but for example, after reacting for 2 minutes at 98 degrees Celsius, 26 cycles of 30 seconds at 98 degrees Celsius, 30 seconds at 58 degrees Celsius, 1 minute at 72 degrees Celsius, and further reacting at 72 degrees Celsius for 5 minutes. The amplification products after reacting are separated by agarose gel electrophoresis, DNA bands containing the objective genes are cut out, and DNA is recovered, then incorporated into a vector for sequencing. The vector for sequencing may be any vector used in said technical area, but, for example, pCR2.1-TOPO (Invitrogen) may be used. The DNA sequence of a gene in the abovementioned prepared vector for sequencing can be determined according to an established method.

[0058] Herebelow, embodiments shall be described, but the present invention is not restricted to these.

Embodiment 1: Construction of a vector for induced transcription control

[0059] In order to introduce a promoter for inductively controlling the expression of a gene into a cell, a cis-acting region necessary for transcription induction was constructed on a vector (figure 1).

[0060] An ECFP gene was attached downstream of a tetracycline inducible promoter (TRE and human cytomegalovirus mini-promoter) and a MAR and an enhancer (3' enhancer) were attached further downstream. Upstream, a spacer sequence of approximately 3 kb (Arg4 yeast gene, 3 kb PstI fragment) was placed, and further, EGFP was attached in the reverse direction to ECFP.

[0061] The PCR method was used for the cloning of the MAR and 3' enhancer. The Exp and High Fidelity PCR system, manufactured by Roche, was used for PCR, and this was conducted with a primer designed in the following way, using DT40 genomic DNA as a template.

40 MAR:

Forward direction primer
5'-GCTGCAGTGCCTTGGGGGTGAAAT-
TCAG-3' (Sequence Number: 1)

45

Reverse direction primer
5'-GCTCTAGAACTGCCCCATTA AAAACTT-
TC-3' (Sequence Number: 2)

50 3' enhancer:

Forward direction primer
5'-GCTCTAGAAAGGCACAGCGCTGTCAG-
GGTGC-3' (Sequence Number: 3)

55

Reverse direction primer
5'-CCGCGGCCGCGTGGTGGGAGCGGGCAG-
GGG-3' (Sequence Number: 4)

[0062] After PCR products were once TA cloned to pCR2.1 vectors, and subsequently digested with an appropriate restriction enzyme (MAR: PstI, XbaI, 3' enhancer: XbaI, NotI), they were re-cloned into pBluescriptII. For the tetracycline inducible promoter, an XhoI EcoRI fragment containing TRE, from pTRE2hyg (Clontech), was cloned into pBlueScriptII. The ECFP gene was inserted into this plasmid containing TRE. For the ECFP gene, NheI, MluI fragments containing ECFP from pECFP-C1 plasmids (Clontech) were blunt-end ligated into pHS305 digested with EcoRI (pHS344). Further, MARs cloned into pBluescriptII were inserted into a downstream portion of pHS344 ECFP (pHS345). As mentioned above, an ARG4 yeast gene was used as a spacer sequence. This is a 3 kb fragment created by the PstI attached to pUC119, but EGFP fragments (NheI, BspEI fragments of pEGFP-C1) were blunt-end ligated to SnaBI fragments of these plasmids. From here, sequences containing EGFP and ARG4 genes were cut out from PstI, and by blunt-end ligation of XhoI fragments of pHS345, pHS346 was obtained. Further, pHS347 was created by inserting 3' enhancer cloned into pBlueScriptII into pHS346 digested with XbaI and NotI.

Embodiment 2: Preparation of cells wherein DNA homologous recombination is induced

(1) Cell cultivation

[0063] DT40 cells were cultivated in 5% CO₂ and 39.5 degrees Celsius in a CO₂ thermostatic oven. IMDM medium (Invitrogen) was used as the medium, with 10% FBS, 1% chicken serum, 100 units per ml of penicillin, 100 µg/ml of streptomycin, and 55 µM 2-mercapto ethanol added.

(2) Transfection

[0064] 30 µg of an expression vector pTA-Hyg (clontech) expressing tetracycline inducible transcription factor was linearized with HindIII, and additionally, 60 µg of pHS347 was linearized with XmnI, and transfected into DT40 cells according to a known method (Buerstedde et al., 1991). At that time, 2.5 mg/ml hygromycin and 100 ng/ml doxycycline were added to the medium and selection was conducted. As a result among hygromycin resistant clones, clones which express ECFP in the absence of doxycycline were able to be obtained (HS101 cell line).

Embodiment 3: Confirmation of novel gene products

[0065] HS101 was cultivated in the absence of doxycycline, and after measuring the fluorescent intensity with an EPICS ELITE ESP (manufactured by Beckman Coulter), a group emitting fluorescent light thought to be ECFP was observed (figure 2: gate E). When cultivation was continued, a group with an even stronger fluorescent in-

tensity than ECFP was observed (figure 2: gate F). When the ratio of E to F was calculated, it was observed that the existence ratio of F increased depending on the cultivation time from the start of induction (figure 3). Further, when cells on the fifth day of cultivation after induction were observed with a fluorescent microscope, among cells emitting ECFP fluorescent light, cells emitting fluorescent light thought to be GFP were observed (figure 4).

10 Embodiment 4: Analysis of novel genetic sequence

(1) Extraction of genomic DNA

[0066] Cells in the vicinity of gate F in FACS were sorted, and 5000 cells were gathered in a silicone coated 1.5 ml tube, with an EPICS ELITE ESP (manufactured by Beckman Coulter). The cells suspended in the sheath fluid were recovered by centrifuge (1000 g, 10 min), 100 µl of genome extraction buffer was directly added to the pellet (100 mM Tris-HCl (pH 8.0), 5 mM EDTA, 0.2% SDS, 200 mM NaCl, and 100 µg/ml of proteinase K), and digestion was done overnight at 50 degrees Celsius. The next day, 250 µl of ethanol was added, and mixing was done by gently shaking up and down and inverting. The genome was recovered by centrifuge (1000 g, 10 min), washed with 70% ethanol, and air-dried. 100 µl of TE (10 mM Tris-HCl (pH 8.0), 1 mM EDTA) was added, and after letting stand for 30 minutes at 50 degrees C, this was dissolved at 4 degrees Celsius overnight.

(2) Analysis of sequence downstream of TRE

[0067] From the extracted genome, a sequence directly downstream of tetracycline inducible promoter was amplified by PCR (Perkin Elmer 9600) by TA cloning, and the sequence was analyzed.

[0068] With 5 µl of genomic DNA fluid (corresponding to 5000 cells) as a template, 10 pmol respectively of upstream (5'-CCATAGAAGACACCGGGACCGATCC-3') (Sequence Number: 5) and downstream primer (5'-TGCACGCTGCCGTCCTCGATGTTG-3') (Sequence Number: 6) was used. Using Pyrobest DNA Polymerase (manufactured by TaKaRa), a reaction was done at the 50 µl scale. The reaction conditions were, after 2 minutes at 98 degrees Celsius, 26 cycles were done of 30 seconds at 98 degrees Celsius, 30 seconds at 58 degrees Celsius, 1 minute at 72 degrees Celsius, and reaction was finally done for 5 minutes at 72 degrees Celsius. Subsequently, 1 µl of ExTaq DNA Polymerase (manufactured by TaKaRa) was added, and after reacting for 15 minutes at 72 degrees Celsius, the entire reaction fluid was separated by agarose gel electrophoresis. The bands corresponding to a light chain gene variable region were cut out, and after DNA was recovered by a Gel Extraction kit (manufactured by Qiagen), this was incorporated into a pCR2.1-TOPO vector, and transformed into *e. coli*. Plasmids were extracted, and sequenced using an ABI PRISM 377 DNA Sequencer (manufactured

by Perkin Elmer). As a result, as shown in figure 5, it is clear that a chimera of ECFP and EGFP has been created. Among the four types of chimeras, 36 clones of type1, 1 clone of type2, 4 clones of type3, and 1 clone of type4 were respectively obtained. Additionally, 18 clones of ECFP were also obtained that are thought to be contaminants from sorting.

Industrial Applicability

[0069] In the present invention, since objective genes having genetic diversity are formed spontaneously without the need for the production of a library, screening is possible immediately thereafter within an animal cell, so that it becomes possible to greatly reduce the amount of time until the objective gene is obtained.

[0070] Additionally, in the present invention, since homologous recombination is continuously induced in a variety of locations, it is in principle possible to increase the diversity of obtained recombinants, and this is applicable as an art that creates genes having novel activity and functions.

Citations

[0071]

Buerstedde, J.M. and Takeda, S. 1991. Cell 67:179-88
Chen, C. and Okayama, H. 1988. Bio-Techniques 6:632-638.

Chauhan, S.S. and Gottesman, M.M. 1992. Gene 120:281-186
Cramer et al., 1998. Nature 391:288-291

DiSanto et al., 1995. Proc. Natl. Acad. Sci. USA 92:377-3 81

Elroy-Stein, O. and B. Moss. 1990. Proc. Natl. Acad. Sci. USA. 87:6743-6747.

Kaufman, R.J. 1990. Methods Enzymol. 185:487-511.

Lahti, J.M. 1999. Methods. 17:305-312.

Neumann, E., M. et al., 1982. EMBO J. 1:841-845.

Nickoloff, J.A. 1992. Mol. Cell. Biol. 12:5311-5318.

Nickoloff, J.A. and Reynolds, R.J. 1990. Mol. Cell. Biol. 10:4837-4845.

Rouet et al., 1994. Proc. Natl. Acad. Sci. USA 91:6064-60 68

SEQUENCE LISTING

[0072]

<110> RIKEN Japan Science and Technology Agency

<120> Conditional regulation of recombination between homologous DNA sequences

<130> RFH15-067 ; A 132-14US (PCT)

<150> JP2002-376555

<151> 2002-12-26

<160> 6

<170> PatentIn version 3.1

<210> 1

<211> 29

<212> DNA

<213> Gallus gallus

<220>

<221> primer_bind

<222> (8)..(29)

<223>

<400> 1

gctgcagtgt cctgggggt gaaattcag 29

<210> 2

<211> 30

<212> DNA

<213> Gallus gallus

<220>

<221> primer_bind

<222> (9)..(30)

<223>

<400> 2

gctctagaac tgccccatt aaaaacttc 30

<210> 3

<211> 30

<212> DNA

<213> Gallus gallus

<220>

<221> primer_bind

<222> (8)..(30)

<223>

<400> 3

gctctagaag gcacagcgct gtcagggtgc 30

<210> 4

<211> 30

<212> DNA

<213> Gallus gallus

<400> 4

ccgcgccgc gttgtgggag cgggcagggg 30

<210> 5

<211> 25

<212> DNA
 <213> Escherichia coli

<220>
 <221> primer_bind
 <222> (1)..(25)
 <223>

<400> 5
 ccatagaaga caccgggacc gatcc 25

<210> 6
 <211> 24
 <212> DNA
 <213> Escherichia coli

<220>
 <221> primer_bind
 <222> (1)..(24)
 <223>

<400> 6
 tgcacgctgc cgctctcgat gttg 24

Claims

1. A method for inducing homologous recombination between a gene and a base sequence having 60%, notably 70%, in particular, 80% and, more particularly, 90% or greater sequence identity to the entirety of the gene for which homologous recombination is induced in a DT40 cell, comprising controlling transcription of said gene by a transcriptional promoter in the DT40 cell, wherein said transcriptional promoter is operably linked to said gene, wherein said base sequence is located on the upstream 5' side of said gene, wherein said base sequence is located on the upstream 5' side of said transcriptional promoter, and wherein said gene is an exogenous gene.
2. A method recited in Claim 1, **characterized in that** a cis-acting region for a transcription control contains either one or both of an enhancer and a nuclear matrix attachment region (MAR).
3. A method recited in Claim 1 or 2, wherein said gene and said base sequence are placed on a vector and wherein said transcriptional promoter is located between said base sequence and said gene.
4. A method recited in Claim 3, **characterized in that** either one or both of an enhancer and a nuclear matrix attachment region (MAR) are inserted onto the aforementioned vector and are operably linked to the aforementioned transcription promoter.

5. A method recited in Claim 3 or 4, **characterized in that** the aforementioned transcription promoter is an inducible promoter.
6. A method recited in Claim 5, **characterized in that** the aforementioned inducible promoter is a tetracycline inducible promoter.
7. A method recited in any one of Claims 3 through 6, **characterized in that** the aforementioned gene is an enhanced cyan fluorescent protein (ECFP) gene.
8. A method recited in any one of Claims 3 through 7, **characterized in that** the base sequence is an enhanced green fluorescent protein (EGFP) genetic sequence.
9. A method recited in any one of Claims 2 through 8, **characterized in that** the aforementioned enhancer is a chicken antibody light chain gene enhancer (3' enhancer), and the aforementioned nuclear matrix attachment region (MAR) is chicken derived.

25 Patentansprüche

1. Verfahren zum Induzieren einer homologen Rekombination zwischen einem Gen und einer Basensequenz, die eine Sequenzidentität von 60 %, insbesondere 70 %, bevorzugt 80 % und mehr bevorzugt 90 % oder höher zu der Gesamtheit des Gens, für das eine homologe Rekombination induziert wird, aufweist, in einer DT40-Zelle, umfassend das Steuern der Transkription des Gens durch einen Transkriptionspromotor in der DT40-Zelle, wobei der Transkriptionspromotor mit dem Gen operativ verknüpft ist, wobei die Basensequenz stromaufwärts der 5'-Seite des Gens angeordnet ist, wobei die Basensequenz stromaufwärts der 5'-Seite des Transkriptionspromotors angeordnet ist, und wobei das Gen ein exogenes Gen ist.
2. Verfahren nach Anspruch 1, **dadurch gekennzeichnet, dass** eine cis-wirkende Region für eine Transkriptionskontrolle entweder eine oder alle beide aus einem Enhancer und einer Kernmatrixanheftungsregion (MAR) enthält.
3. Verfahren nach Anspruch 1 oder 2, wobei das Gen und die Basensequenz auf einem Vektor angeordnet sind und wobei der Transkriptionspromotor zwischen der Basensequenz und dem Gen angeordnet ist.
4. Verfahren nach Anspruch 3, **dadurch gekennzeichnet, daß** entweder eine oder alle beide aus einem Enhancer und einer Kernmatrixanheftungs-

region (MAR) in den zuvor genannten Vektor inseriert und mit dem zuvor genannten Transkriptionspromotor operativ verknüpft sind.


5. Verfahren nach Anspruch 3 oder 4, **dadurch gekennzeichnet, daß** der zuvor genannte Transkriptionspromotor ein induzierbarer Promotor ist.
6. Verfahren nach Anspruch 5, **dadurch gekennzeichnet, daß** der zuvor genannte induzierbare Promotor ein Tetracyclin-induzierbarer Promotor ist.
7. Verfahren nach einem der Ansprüche 3 bis 6, **dadurch gekennzeichnet, daß** das zuvor genannte Gen ein Enhanced Cyan Fluorescent Protein (ECFP)-Gen ist.
8. Verfahren nach einem der Ansprüche 3 bis 7, **dadurch gekennzeichnet, daß** die Basensequenz eine genetische Sequenz eines Enhanced Green Fluorescent Protein (EGFP) ist.
9. Verfahren nach einem der Ansprüche 2 bis 8, **dadurch gekennzeichnet, daß** der zuvor genannte Enhancer ein Antibody Light Chain Gen-Enhancer aus Huhn (3'-Enhancer) ist, und die zuvor genannte Kernmatrixanheftungsregion (MAR) aus Huhn abgeleitet ist.

Revendications

1. Procédé pour induire une recombinaison homologue entre un gène et une séquence de bases ayant 60 %, notamment 70 %, en particulier 80 % et, plus particulièrement, 90 % ou plus d'identité de séquence par rapport à l'intégralité du gène pour lequel la recombinaison homologue est induite dans une cellule DT40, comprenant le contrôle de la transcription dudit gène par un promoteur transcriptionnel dans la cellule DT40, dans lequel ledit promoteur transcriptionnel est lié de façon opérationnelle audit gène, dans lequel ladite séquence de base est située sur le côté 5' en amont dudit gène, dans lequel ladite séquence de base est située sur le côté 5' en amont dudit promoteur transcriptionnel, et dans lequel ledit gène est un gène exogène.
2. Procédé selon la revendication 1, **caractérisé en ce qu'**une région agissant en cis pour contrôler la transcription contient l'un ou les deux d'un amplificateur et d'une région d'attachement à la matrice nucléaire (MAR).
3. Procédé selon la revendication 1 ou 2, dans lequel ledit gène et ladite séquence de bases sont placés

sur un vecteur, et dans lequel ledit promoteur transcriptionnel est situé entre ladite séquence de base et ledit gène.

4. Procédé selon la revendication 3, **caractérisé en ce que** l'un ou les deux d'un amplificateur et d'une région d'attachement à la matrice nucléaire (MAR) sont insérés sur le vecteur susmentionné et sont liés de façon opérationnelle au promoteur de transcription susmentionné.
5. Procédé selon la revendication 3 ou 4, **caractérisé en ce que** le promoteur de transcription susmentionné est un promoteur inductible.
6. Procédé selon la revendication 5, **caractérisé en ce que** le promoteur inductible susmentionné est un promoteur inductible par la tétracycline.
7. Procédé selon l'une quelconque des revendications 3 à 6, **caractérisé en ce que** le gène susmentionné est un gène de la protéine fluorescente cyan amplifiée (ECFP).
8. Procédé selon l'une quelconque des revendications 3 à 7, **caractérisé en ce que** la séquence de bases est une séquence génétique de la protéine fluorescente verte amplifiée (EGFP).
9. Procédé selon l'une quelconque des revendications 2 à 8, **caractérisé en ce que** l'amplificateur susmentionné est un amplificateur de gène de chaîne légère d'anticorps de poulet (amplificateur 3'), et la région d'attachement à la matrice nucléaire (MAR) susmentionnée est dérivée du poulet.






FIG. 2





FIG. 3

FIG. 4






FIG. 5

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- JP 2002376555 A [0072]

Non-patent literature cited in the description

- **BUERSTEDDE, J.M. ; TAKEDA, S.** *Cell*, 1991, vol. 67, 179-88 [0071]
- **CHEN, C. ; OKAYAMA, H.** *BioTechniques*, 1988, vol. 6, 632-638 [0071]
- **CHAUHAN, S.S. ; GOTTESMAN, M.M.** *Gene*, 1992, vol. 120, 281-186 [0071]
- **CRAMERI et al.** *Nature*, 1998, vol. 391, 288-291 [0071]
- **DISANTO et al.** *Proc. Natl. Acad. Sci. USA*, 1995, vol. 92, 377-3 [0071]
- **ELROY-STEIN, O. ; B. MOSS.** *Proc. Natl. Acad. Sci. USA.*, 1990, vol. 87, 6743-6747 [0071]
- **KAUFMAN, R.J.** *Methods Enzymol.*, 1990, vol. 185, 487-511 [0071]
- **LAHTI, J.M.** *Methods*, 1999, vol. 17, 305-312 [0071]
- **NEUMANN, E., M. et al.** *EMBO J.*, 1982, vol. 1, 841-845 [0071]
- **NICKOLOFF, J.A.** *Mol. Cell. Biol.*, 1992, vol. 12, 5311-5318 [0071]
- **NICKOLOFF, J.A. ; REYNOLDS, R.J.** *Mol. Cell. Biol.*, 1990, vol. 10, 4837-4845 [0071]
- **ROUET et al.** *Proc. Natl. Acad. Sci. USA*, 1994, vol. 91, 6064-60 [0071]