



(11) **EP 2 098 246 A1**

(12) **EUROPEAN PATENT APPLICATION**
published in accordance with Art. 153(4) EPC

(43) Date of publication:
09.09.2009 Bulletin 2009/37

(21) Application number: **07851040.1**

(22) Date of filing: **21.12.2007**

(51) Int Cl.:
A61K 45/00 ^(2006.01) **A61K 39/00** ^(2006.01)
A61K 39/395 ^(2006.01) **A61P 37/04** ^(2006.01)
C12N 5/10 ^(2006.01) **C12N 15/09** ^(2006.01)
C12P 21/08 ^(2006.01) **C12Q 1/02** ^(2006.01)
G01N 33/15 ^(2006.01) **G01N 33/50** ^(2006.01)

(86) International application number:
PCT/JP2007/074634

(87) International publication number:
WO 2008/078673 (03.07.2008 Gazette 2008/27)

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IS IT LI LT LU LV MC MT NL PL PT RO SE
SI SK TR**

(30) Priority: **26.12.2006 JP 2006349458**

(71) Applicant: **Japan Science and Technology Agency
Kawaguchi-shi,
Saitama 332-0012 (JP)**

(72) Inventors:
• **TSUBATA, Takeshi
Tokyo 167-0051 (JP)**

• **ONODERA, Taishi
Koganei-shi
Tokyo 184-0014 (JP)**

(74) Representative: **Webber, Philip Michael et al
Frank B. Dehn & Co.
St Bride's House
10 Salisbury Square
London
EC4Y 8JD (GB)**

(54) **METHOD FOR PROMOTING IMMUNE RESPONSE COMPRISING INHIBITING CD22 FUNCTION IN B CELLS**

(57) The purpose of the present invention is to elucidate a relationship between deregulation of signaling by CD22 and a rapid response of B cells by IgG-BCR and the like, and to provide a method capable of inducing a rapid immune response and defending against infection instead of vaccine.

The present invention relates to a method for promoting an immune response causing such a strong pro-

liferation of clones and production of a large amount of antibody-producing cells as those seen in the memory immune response even in the naive B cells expressing IgM-BCR and IgD-BCR by inhibiting The CD22 function in B cells; and to a method for screening a substance capable of promoting the immune response based on a change in the CD22 function in B cells.

EP 2 098 246 A1

Description

FIELD OF THE INVENTION

[0001] The present invention relates to a method for promoting immune response comprising inhibiting CD22 function and the like.

BACKGROUND OF THE INVENTION

[0002] IgM and IgD-positive naive B cells, which reacted with a normal T cell-dependent antigen, will move to splenic red pulp and bridging channel and proliferate there to form extrafollicular foci in an early stage of immune response. Plasmablast will proliferate in the extrafollicular foci and undergo differentiation into antibody-producing cells so as to eliminate an antigen. Most of the antibody-producing cells generated in the early stage will have a short life span, while a part of them will induce class-switching into IgG class and the like. A part of B cells that has reacted with an antigen will proliferate in follicle and form a germinal center. B cells will accumulate point mutations in their immunoglobulin V-region during the germinal center reaction so as to increase their affinity for the antigen. Most of B cells in the germinal center will induce the class-switching. The antibody-producing cells that have gone through the above germinal center reaction will bear the production of antibodies thereafter. A part of B and T lymphocytes that were activated by antigen-stimulation in a first immune response will grow into memory lymphocytes. Memory B cells will be generated from a part of the germinal center B cells. The memory lymphocytes will survive for a long period of time. When being exposed to the same antigen, the memory lymphocyte will proliferates more rapidly proliferate than the naive B cells and form a huge foci of plasmablasts and plasma cells in the splenic red pulp and bridging channel, resulting in a rapid and large-scale production of antibodies. Vaccine will defend against infection by inducing immune memory.

[0003] B cell antigen-receptor (BCR) consists of a membrane-type immunoglobulin and signal component Ig α /Ig β molecules. Upon the reaction with the antigen, BCR will intracellularly transduce a signal through the Ig α /Ig β molecules (class II antigen). Coreceptors of BCR such as CD19, CD22, CD72 and Fc γ R2B modulate an appropriate threshold for signaling by positively or negatively regulating BCR signal transduction (signaling), so that they will have an influence on determination of the fate of B cells such as proliferation, differentiation into the antibody-producing cells, non-response or apoptosis. Among them, CD22 is known as a molecule that will downmodulate the BCR signaling. Three immune receptor tyrosine-based inhibition motifs (ITIM) existing in an intracellular region of CD22 will be phosphorylated directly after BCR crosslinking has occurred, and CD22 will recruit a phosphatase, Src-homology domain 2 (SH2)-containing protein tyrosine phosphatase-1 (SHP-

1) near BCR so as to activate it and and negatively regulate the BCR signaling.

[0004] The memory B cells will be generated through the differentiation mainly from cells that have been class-switched into IgG, and express the membrane-type IgG as the BCR. IgG-BCR will transduce a signal through the Ig α /Ig β molecules like IgM-BCR and IgD-BCR expressed by the naive B cells. Recently, a finding has been obtained that the IgG-BCR is involved in the rapid response of the memory B cells (Wakabayashi, C., et al., Science, 2002, 298(5602): p.2392-5). Goodnow et al. made a transgenic mouse in which almost all of the B cells expressed IgG specific for HEL, and analyzed an immune response of IgG-positive naive B cells. They revealed that the IgG-positive naive B cells would rapidly proliferate upon the antigen stimulation similarly in the case of the memory immune response, so that they would form a huge foci of the plasmablast and produce a large amount of antibodies (Martin, S.W. and C.C. Goodnow, Nat Immunol. 2002, 3(2): p.182-8). These results show that the IgG-BCR has a function different from the IgM-BCR or IgD-BCR, and is involved in a rapid activation of B cells in the memory response

[0005] On the other hand, the present inventors found that while the IgM-BCR or IgD-BCR were negatively regulated by CD22, the IgG-BCR would not be regulated by CD22 so that it could induce an effective signaling (Wakabayashi, C., et al., Science, 2002. 298(5602): p. 2392-5).

[Non Patent Document 1] Nitschke, L. and T. Tsubata, Trends Immunol, 2004. 25(10): p.543-50

[Non Patent Document 2] Tedder, T.F., et al., Annu Rev Immunol, 1997. 15: p.481-504

DISCLOSURE OF THE INVENTION

PROBLEMS TO BE SOLVED BY THE INVENTION

[0006] However, it has been unknown how much the release of signal regulation by CD22 is involved in the rapid response of B cells through IgG-BCR, or whether or not the regulation of CD22 function could induce such a rapid antibody-response as that in the memory response.

[0007] In order to solve the above problems, the present inventors analyzed in detail kinetics in vivo of NP (nitrophenol)-reactive and CD22-deficient B cells in immune response, which cells were obtained by breeding a CD22-deficient mouse with a QM mouse most of whose B cells were reactive with a hapten, NP. As a result, it has been revealed that there will occur such a strong proliferation of clones and production of a large amount of antibody-producing cells as those seen in the memory immune response even in the naive B cells expressing IgM-BCR and IgD-BCR due to the deficiency of CD22, leading to the completion of the present invention.

MEANS FOR RESOLVING PROBLEMS

[0008] The present invention comprises the following aspects:

[Aspect 1] A method for promoting an immune response comprising inhibiting CD22 function in B cells.

[Aspect 2] A method according to Aspect 1, wherein the B cells are IgM and IgD positive naive B cells.

[Aspect 3] A method according to Aspect 1 or 2, wherein the promotion of the immune response comprises the increase of proliferation, division and/or survival of the B cells in an early stage of the immune response.

[Aspect 4] A method according to Aspect 3, wherein the B cells comprises IgG1+B cells, IgM⁺ B cells and germinal center B cells.

[Aspect 5] A method according to Aspect 1 or 2, wherein the promotion of the immune response comprises class switching and/or differentiation of the B cells in an earlier stage.

[Aspect 6] A method according to Aspect 5, wherein the class switching is that to the IgG1⁺ positive cells, and/or the differentiation is that into antibody-producing cells or the germinal center B cells.

[Aspect 7] A method according to any one of Aspects 2 - 6, wherein the early stage in the immune response means from the third to fifth days after the immunization.

[Aspect 8] A method according to Aspect 1, wherein the promotion of the immune response comprises increase in a number of the antibody-producing cells and in an amount of antibodies produced, and/or the production of antibodies in an earlier stage.

[Aspect 9] A method according to Aspect 8, wherein the antibodies produced in the earlier stage are IgG and/or IgM.

[Aspect 10] A method according to any one of Aspects 1 - 9, wherein the inhibition of the CD22 function is caused by its dysfunction due to deficiency or mutation of CD22 gene.

[Aspect 11] A method according to any one of Aspects 1 - 9, wherein the inhibition of the CD22 function is caused by repression of the expression of CD22 gene.

[Aspect 12] A method according to Aspect 11, where-

in the repression of the expression of CD22 gene is done at a transcriptional level.

[Aspect 13] A method according to Aspect 12, wherein gene knockdown by means of siRNA specific for CD22 gene is used.

[Aspect 14] A method according to Aspect 13, wherein the siRNA is a double strand RNA consisting of an oligonucleotide having 21 - 23 bases and its complementary oligonucleotide, which comprises a continuous base sequence of a part of CD22 gene.

[Aspect 15] A method according to any one of Aspects 1 - 9, wherein the inhibition of the CD22 function is related to the inhibition of signal transduction of B cell receptor.

[Aspect 16] A method according to any one of Aspects 1 - 15, wherein the B cells and/or the CD22 or CD22 gene is derived from a mouse.

[Aspect 17] B cells whose CD22 function is inhibited.

[Aspect 18] B cells according to Aspect 17, which are isolated from a knockout animal wherein CD22 gene is knocked out.

[Aspect 19] A method for the production of an antibody using the method according to any one of Aspects 1 - 15 or the B cells whose CD22 function is inhibited according to Aspect 17 or 18.

[Aspect 20] A method for the production according to Aspect 19, wherein the antibody is specific for a T cell-dependent antigen.

[Aspect 21] A method for screening of a substance capable of promoting the immune response based on a change in the CD22 function in the B cells.

[Aspect 22] A method for screening according to Aspect 21, comprising the following steps:

- (a) a step of contacting a substance to be screened with the B cells;
- (b) a step of detecting the CD22 function in said cells; and
- (c) a step of selecting a substance which inhibits said function.

[Aspect 23] A method for screening according to any one of Aspect 19 - 22, wherein the B cells and/or the CD22 or CD22 gene is derived from a mouse.

[Aspect 24] A screening kit used in a method according to any one of Aspects 21 - 23.

ADVANTAGES OF THE INVENTION

[0009] It has been first revealed by the present invention that the immune response can be promoted by inhibiting or repressing the CD22 function in the B cells. And, according to the present invention, such a strong proliferation of clones and a mass production of the antibody-producing cells as those seen in the memory immune response can be effected even in the naive B cells expressing IgM-BCR and IgD-BCR, by inhibiting the CD22 function. These results suggest that a rapid activation of the B cells will occur in the memory immune response since the signaling mediated by IgG-BCR expressed in the memory B cells is not inhibited by CD22, showing the possibility to defend against infection by inducing the rapid immune response instead of a vaccine.

BRIEF DESCRIPTION OF THE DRAWINGS

[0010]

Fig.1 are figures showing the occurrence of burst of extrafollicular plasmablast in CD22-deficient B cells in the early stage of immune response. (A) Flow cytometry analysis of isolated B cells, showing the percentage of CD23⁺ and NIP-reactive BCR⁺ B cells (left) and an expression amount of NIP-reactive BCR in a histogram (right). (B) A protocol of adopted transfer. (C) Spleen tissue sections observed with a confocal fluorescence microscope, showing that the burst of the extrafollicular plasmablast occurred in the CD22-deficient B cells in the early stage of the immune response.

Fig.2 shows the results obtained with flow cytometry analysis. Spleen cells were collected from the recipient mice that were transplanted with B cells of CD22^{+/+}QM (lower polygonal line) and B cells of CD22^{-/-}QM (upper polygonal line), respectively; on the third, fifth and seventh days after the immunization. The expression of B220 (A-F), Ly5.2 (A-F), IgM (C) and IgG1 (D), and NIP and PNA reactivity (E, F) were analyzed. Dead cells were stained with DAPI and excluded from analysis. In spleen cells (DAPI, 2x 10⁵) of the recipient mice, the numbers of the B cells derived from Donor (B220⁺, Ly5.2⁺) (A, B in Fig.2), the IgM⁺ B cells derived from Donor (B220⁺, Ly5.2⁺, IgM⁺) (C in Fig.2), the IgG1⁺ B cells derived from Donor (B220⁺, Ly5.2⁺, IgG1⁺) (D in Fig.2), and the germinal center B cells derived from Donor (B220⁺, Ly5.2⁺, reactive with NIP and PNA) (E, F in Fig.2) were counted, respectively. An average from 4 individuals in each group is shown as a bar.

Fig. 3 shows the results of the proliferation of the Donor B cells, class-switching and differentiation into the germinal center B cells, obtained with flow cy-

tometry analysis. (A) Division of the B cells. As a negative control, only PBS was injected after the transportation of the B cells, and analyzed on the second day after the injection (the lowest figures). Dead cells were stained with DAPI and excluded from analysis, and the results of analysis of spleen cells (DAPI, 2x 10⁵) of each recipient mouse were shown. The experiment was carried out twice giving the same results. (B) Ig class-switching dependent on cell division and differentiation into the germinal center B cells. Percentages of the IgG1⁺ B cells (upper and middle) and the germinal center B cells (lower) were shown, respectively. Dead cells were stained with DAPI and excluded from analysis, and the results of analysis of spleen cells (DAPI, 2x 10⁵) of each recipient mouse were shown. The experiment was carried out twice giving the same results.

Fig. 4 (A, B) shows the number of antibody-producing cells in spleen. Spleen cells were collected from the recipient mice that were transplanted with the B cells of CD22^{+/+}QM (lower polygonal line) and the B cells of CD22^{-/-}QM (upper polygonal line), respectively, on the third, fifth and seventh days after immunization. Among all of the spleen cells (1x 10⁶), the numbers of cells derived from the Donor B cells, which were producing IgM^a antibody (A) and IgG1^a antibody (B) specific for NP, were determined by ELISPOT method. An average from 4 individuals in each group is shown as a bar. (C, D) Titer of NP-specific antibody derived from the Donor B cells in serum. A production amount (μg/ml) of IgM^a antibody (C) and IgG1^a antibody (D) specific for NP in serum of the recipient mice, which were derived from the B cells of CD22^{+/+}QM (lower polygonal line) and the B cells of CD22^{-/-}QM (upper polygonal line), respectively, was determined by EILSA method. An average from 4 individuals in each group is shown as a bar.

BEST MODE FOR CARRYING OUT THE INVENTION

[0011] According to the present invention, the promotion of the immune response was effected by inhibiting (or, repressing) the CD22 function in the B cells. Although the term "B cells" used in the present specification usually means IgM and IgD positive naive B cells (mature B cells), it is not limited to these types and comprises B cells that potentially have capability of differentiating into antibody-producing cells in response to the T antigen-dependent antigen. Furthermore, in the present specification, various kinds of cells including the B cells and the CD22 (gene) may be derived from any animal cells known for those skilled in the art, including mammals such as human and mouse.

[0012] The promotion of the immune response according to the present invention comprises any reaction known for those skilled in the art, for example, the in-

crease in proliferation and survival of the B cells in an early stage of the immune response, especially, the increase of in the proliferation, division and/or survival of the B cells, such as the increase in numbers of the IgG1⁺B cells, the IgM⁺ B cells and the germinal center B cells by a factor of two to three compared to the CD22^{+/+} B cells, the class switching of the B cells into the IgG1⁺ positive cells, and/or differentiation in an earlier stage into, for example, the antibody-producing cells or the germinal center B cells, the increase in a number of cells producing antibodies such as IgG and/or IgM in immune organs such as spleen, and in an amount of the antibodies produced in serum (about three to five times more than that of CD22^{+/+} B cells), and/or the production of these antibodies in an earlier stage, and the promotion of signal transduction through the B cell receptor.

[0013] The "early stage in an immune response" usually means few days such as from the third to the fifth days after the immunization (stimulation or sensitization by an antigen), with a variation of about one or two days depending on the kind of an antigen used as an immunogen, aspects in the promotion of the immune response and the like.

[0014] The inhibition of the CD22 function in the B cells may be carried out by any genetic (genomic) engineering technique or method known for those skilled in the art. As long as the immune response is significantly promoted, it is not necessary to completely inhibit the CD22 function.

[0015] The inhibition of the CD22 function may be caused by any action on the CD22 gene per se, such as by dysfunction due to deficiency or mutation of the CD22 gene. Such action on the gene per se may be effected by, for example, knockout of the CD22 gene by means of gene targeting technique known for those skilled in the art, or introduction of mutation into the gene by means of various methods for that so as to express CD22 variant not having a prescribed function.

[0016] The inhibition of the CD22 function may be caused by repression of the expression of the CD22 gene, which may be done at any optional level such as a transcriptional level.

[0017] For example, the repression of the expression of the gene at the transcriptional level may be done gene by knockdown by means of a nucleic acid sequence inducing RNA interference (RNAi) against the CD22 gene (DNA or mRNA) (a single or double stranded RNA having a sequence specific for a partial sequence of said gene, such as siRNA) ; antisense RNA; or various ribozymes.

[0018] The nucleic acid sequence inducing the RNA interference against the CD22 gene may be optionally designed and prepared by those skilled in the art, based on a base sequence of the CD22 gene registered in any known public database, such as accession No.L16928 GenBank/EMBL/DDBJ for mouse CD22 gene. The siRNA is a double strand RNA consisting of an oligonucleotide having 15-30, especially 21-23 bases and its complementary oligonucleotide, which comprises a continu-

ous base sequence of a part of the CD22 gene. The siRNA may comprise one that will be intracellularly generated upon digestion by Dicer.

[0019] It is also possible to inhibit the CD22 function at the transcriptional level by manipulating a transcriptional-regulating region of the CD22 gene so as to inhibit its normal transcription.

[0020] Furthermore, it is possible to inhibit the CD22 function by expressing a substance that will inhibit the CD22 function at any stage with any transformation method known for those skilled in the art.

[0021] The CD22 function may be inhibited in the B cells in vivo or in vitro. In vivo, such inhibition may be effected by producing a knock-out animal such mouse whose CD22 gene has been knocked out.

[0022] Accordingly, the present invention also relates to B cells (population) whose The CD22 function is inhibited. Such cells may be prepared by isolating CD23⁺ follicle cells by means of any sorting technique known for those skilled in the art from the spleens cells derived from the above knock-out animal or mammal such as the knock-out mouse, or a mammal such as mouse that is genetically related to them (for example, animals such a QM mouse as described in Example in the present specification, which are obtained by being bred with other animals such as a mouse that will react with a particular antigen (hapten)). An example of the above B cells is a spleen cell (IgH^a, Ly5.2) of CD22^{-/-}QM mouse prepared in Example 1. Alternatively, it is possible to easily prepare the B cells (population) whose The CD22 function is inhibited by using any one of the above genetic engineering technique. Furthermore, it is possible to obtain the B cells whose CD22 function is inhibited by knocking out the CD22 gene in various stem cells such as ES cell and differentiating them into B cells.

[0023] The above B cells whose CD22 function is inhibited may be used not only in the method according to the present invention, but also utilized effectively in methods for the analysis of the immune response or for screening a substance capable of promoting an immune response.

[0024] The present invention is also related to a method for the production of an antibody using any one of the above methods or the above B cells whose CD22 function is inhibited. The antibody may be prepared by using any method or technique known for those skilled in the art for any antigen such as a T cell-dependent antigen.

[0025] Since the immune response will be promoted by inhibiting the CD22 function as already described, it will be possible to screen a substance capable of promoting the immune response, for example, based on a change in the CD22 function in the B cells. Accordingly, the present invention also relates to such screening method and a screening kit used in the method.

[0026] One example of the screening method according to the present invention comprises the following steps:

- (a) a step of contacting a substance to be screened with the B cells;
- (b) a step of detecting the CD22 function in said cells; and
- (c) a step of selecting a substance which inhibits said function.

[0027] The change (inhibition) of the CD22 function may be directly detected by determining a degree of inhibition of the expression of the CD22 gene at any stage such as transcription and translation with any method known for those skilled in the art.

[0028] An amount of the expression of the CD22 gene may be determined by an amount of its mRNA in the above screening method. The determination of mRNA may be carried out by any methods known for those skilled in the art such as various quantitative PCR methods such as RT-PCR and micro array (DNA chip) using primers designed optionally based on a base sequence of the gene encoding CD22 protein. Alternatively, the change (inhibition) of the CD22 function may be indirectly detected by detecting various aspects involved in the promotion of the immune response by the inhibition of the CD22 function as already described.

[0029] The cells such as the B cells and/or the CD22 or CD22 gene used in the screening method according to the present invention may be derived from any known animal including mammals such as a mouse and human. The screening may be carried out in vivo or in vitro. In vitro, for example, the B cells may be contacted with the substance to be screened by culturing the cells in the presence of said substance. In vivo, the B cells may be contacted with the substance to be screened in a body by administrating said substance into a experiment animal with a suitable method.

[0030] The kit used in the screening method according to the present invention may be optionally constituted depending on specific mechanism on detection and determination of the CD22 function in B cells and the like. The kit may comprises, for example, an agent comprising an antibody specifically recognizing CD22 protein, primers for amplification of the CD22 gene and a probe for hybridization, which may have a suitable length, for example, of a base sequence having continuous 10 - 100 bases depending on their use.

[0031] The above primers, probe and antibody comprised in the above kit as a constituent may be labeled with a suitable labeling substance such as a radioactive material, fluorescent material, pigment and the like. The kit may further comprise any other elements or components known for those skilled in the art such as various agents, enzymes, buffer, a reaction plate, etc. depending on its structure and purpose for use.

[0032] As already described, the cells such as the B cells and/or the CD22 or CD22 in the present specification may be derived from any known animal including mammals such as a mouse and human. The information about these genes is known for those skilled in the art,

and may be easily obtained from various public database (cf. GenBank/EMBL/DDBJ, accession No.X59350).

[0033] DNAs (genomic DNA and cDNA) encoding CD22 according to the present invention include in addition to DNA having the above specific base sequence (nucleotide sequence) a DNA hybridizing under stringent conditions with a DNA consisting of a base sequence complementary with that of the above DNA, and a DNA consisting of a base sequence showing identity (homology) of about 80% or more, preferably about 95% or more with that of the above DNA, which encode a protein substantially having at least one of the CD22 functions.

[0034] The hybridization may be performed in accordance with a method known in the art, for example, that described in Molecular cloning third ed. (Cold Spring Harbor Lab. Press 2001) or in Current protocols in molecular biology (edited by Frederick M. Ausubel et al., 1987). When a commercially available library is used, the hybridization may be done according to instructions attached to it.

[0035] The term "stringent conditions" means in this specification, for example, those of sodium concentration of 15-900 mM, preferably 15-600 mM, more preferably 15-150 mM, pH of 6-8 at 60°C-68°C.

[0036] Thus, the DNA that may hybridize with DNA consisting of the base sequence complementary with that of the DNA consisting of the above base sequence may include, for example, a DNA comprising a base sequence having identity of about 80% or more, preferably of about 95% or more, and more preferably of about 99% or more on a total average with the whole base sequence of the DNA of interest.

[0037] The above DNAs may be obtained by PCR, for example, using suitable primers prepared on the information about the base sequence disclosed in Non Patent Document 1 or in the present specification. The DNA according to the present invention may be also prepared by chemical synthesis.

[0038] The CD22 protein according to the present invention may include a polypeptide having an amino acid sequence wherein one or a few amino acids are deleted, substituted or added with respect to an amino acid sequence encoded by the above base sequence, which substantially has at least one of the CD22 functions.

[0039] In order to determine the identity of sequences between two amino acid sequences or two nucleotide (base) sequences, the sequences will be pre-treated to become in an optimum condition for comparison. For example, gaps may be inserted into one of the sequences so that alignment with the other sequence will be optimized. After such pre-treatment, amino acid residues or bases in each region will be compared. When amino acid residues or bases located at a certain position in a first sequence are the same as those of a second sequence located at a position corresponding to the above position in the first sequence, these two sequences will be considered as the same at the position. Identity between the two sequences will be shown by percentage of the

number of the amino acid residues or bases that are the same in the two sequences for the number of the whole amino acid residues or bases.

[0040] According to the above principle, identity between the two amino acid sequences or nucleotide sequences may be determined, for example, by algorithm of Karlin and Altschul (Proc. Natl. Acad. Sci. USA 87:2 264-2268, 1990; and Proc. Natl. Acad. Sci. USA 90: 5873-5877, 1993). BLAST or FASTA programs based on the above algorithms may be used for searching a sequence having a high identity with a given sequence from database. Such programs are available at a Web site of National Center for Biotechnology Information on the Internet.

[0041] The DNA showing the above identity in the amino acid sequence or nucleotide sequence may be obtained by resorting to the above hybridization, or may be easily found in public databases or a group of functionally unidentified DNAs obtained with analysis of genome sequences by means of methods conventionally used by those skilled in the art, for example, by searching with the above BLAST software.

[0042] The present invention will be further explained by the following examples, which do not limit the scope of the present invention. The genetic procedures in the examples are done in accordance with any standard techniques in gene engineering and molecular biology such as those described in Sambrook and Maniatis, in Molecular Cloning - A laboratory Manual, Cold Spring Harbor Laboratory Press, New York, 1989; Ausbel, F. M. et al., Current Protocols in Molecular Biology, John Wiley & sons, New York 1995. The disclosure cited in the present specification as a whole will constitute a part of the present specification.

Example 1

Burst of extrafollicular plasmablast of CD22-deficient B cells in an early stage of immune response

[0043] In order to reveal kinetics of the CD22-deficient B cells that have reacted with an antigen in the immune response, the CD22-deficient QM mouse (CD22^{-/-}QM mouse) was prepared by breeding the QM mouse, most of whose B cells will react with nitrophenol (NP), with the CD22-deficient mouse provided Duke University. Among the B cells, follicle B cells are involved in the reaction against the T cell-dependent antigen. Accordingly, spleen cells (IgH^a, Ly5.2) of the CD22^{-/-}QM mouse and the CD22^{+/+}QM (QM mouse without deficiency of CD22) were stained with PE-labeled anti-CD23 antibody (Cat No. 12-0232-85:e-Bioscience) and Magnet beads-labeled anti-PE antibody (Mat. No.120-000-294:Miltenyl biotec), followed by positive sorting with Auto MACS to obtain NP-reactive CD23⁺ follicle B cells (Fig.1A). An amount of the surface immunoglobulin (BCR) specific for NP expressed in the CD22^{-/-} B cells is half that in the CD22^{+/+} B cells (Fig.1A). These results coincide with the

report that the expression amount of BCR would decrease in CD22^{-/-} B cells. The CD22-deficient mouse is available from Core F of Consortium for Functional Glycomics (CFG) or Jackson Laboratory (600 Main Street Bar Harbor, Marine 04609 USA), whose original publication name is mSiglec-2; Colony Code Name:CD22.

[0044] The follicle B cells (4 x 10⁵ per mouse) prepared in the above were administered by an intravenous injection into C57BL/6 Ly5.1 congenic mice (IgH^b, Ly5.1) primed in advance with CGG (100 μg) and CFA adjuvant in order to activate helper T cells (Day -1). In the next day (Day 0), the mice were immunized with the T cell-dependent antigen, NP-CGG (50 μg) and alum adjuvant. On the third, fifth and seventh days after the immunization, spleen tissue sections from the recipients transplanted with the CD22^{-/-} QM B cells and the CD22^{+/+} QM B cells, respectively, were stained with an anti-CD38 antibody (Blue) (Cat.No.12-0381-81:e-Bioscience), an anti-IgM^a antibody (Cat.No.553515:BD PharmMingen) ((a)-(f)), an anti-IgG1^a antibody (Green)(Cat.No.553500: BD Pharmingen) ((g)-(l)), or a NIP-APC (Red), and observed with the confocal fluorescence microscope (Fig. 1C). The follicle B cells originated in the recipients were stained only with the anti-CD38 antibody, while the B cells derived from Donors were stained with the anti-IgM^a antibody, the anti-IgG1^a antibody and the NIP-APC. Burst derived from the B cells of CD22^{-/-} Donor was found in the Bridging Channel (arrow) and splenic red pulp (arrowhead).

[0045] The results showed that while a small number of the Donor-derived B cells were scattered in the boundary between the follicle B cells and T cell region in the mice transplanted with CD22^{+/+} B cells on the third day after the immunization, proliferation of a large number of the B cells was observed in the same region of the mice transplanted with the CD22^{-/-} B cells, most of which were IgM-positive B cells. While only a small amount of plasmablast was increased in the CD22^{+/+} B cells in Bridging Channel on the fifth day after the immunization, the CD22^{-/-} B cells showed an explosive increase of plasmablast both in the Bridging Channel and splenic red pulp, most of which were found to be class-switched into IgG1. In either the CD22^{-/-} B cells or the CD22^{+/+} B cells, most of the B cells were found in the follicle B cells or the germinal center on the seventh day after the immunization, with only a few plasma cells in the splenic red pulp but no a cluster of such plasmablast as found on the fifth day after the immunization.

[0046] The above result showed that the CD22^{-/-} B cells could produce a large amount of plasmablast in such an early stage as on the fifth day after the immunization. This is very similar to the phenomenon that memory B cells would produce a large amount of plasmablast in an early stage of the secondary immune response, and in common with extrafollicular proliferative burst observed on the fifth day in the immune response of IgG transgenic B cells, as reported by Goodnow et al. These results showed that deregulation of the signaling by

CD22 would induce such an explosive proliferation as found in the memory immune response. Four mice used in each group were analyzed to give the same result.

Example 2

Rapid clone-proliferation and sustention of survival of CD22^{-/-} B cells after reaction with an antigen

[0047] The mechanism of the proliferation of a large amount of plasmablast found in the early stage of the immune response of CD22^{-/-} B cells was then studied. It was known that B cells that had reacted with a hapten NP during the immune response would induce class-switching from IgM mainly into IgG1, and that the germinal center B cells would be produced in response to the T cell-dependent antigen. Accordingly, after the CD22^{-/-} B cells and the CD22^{+/+} B cells (4×10^5 cells each) were transplanted into the recipient mice, and the numbers of the total B cells, the IgM-positive B cells, the IgG1-positive B cells or the germinal center B cells, which were derived from the Donor, were counted with the flow cytometry.

[0048] It was found that in both the CD22^{-/-} and CD22^{+/+} QM B cells, all of the total B cells, the IgM-positive B cells, the IgG1-positive B cells and the germinal center B cells had been increased until the fifth day after the immunization, and then decreased on the seventh day after the immunization (Fig.2). There was a tendency that the IgG1-positive B cells would increase a little bit more late than the IgM-positive B cells would (Fig.2D). At a peak time on the fifth day after the immunization, the number of the CD22^{-/-} QM B cells increased about twice or three times as much as that of the CD22^{+/+} QM B cells with respect to all of the total B cells, the IgM-positive B cells, the IgG1-positive B cells and the germinal center B cells (Fig.2E, F). On the other hand, the number of the CD22^{-/-} QM B cells more drastically dropped on the seventh day after the immunization than that of the CD22^{+/+} QM B cells did, down to the same level as the CD22^{+/+} QM B cells. Thus, it was revealed that the CD22^{-/-} QM B cells increased more than the CD22^{+/+} QM B cells did in the early stage of the immune response in each of the sub-populations of the IgM⁺ B cells, the IgG1⁺ B cells and the germinal center B cells.

[0049] In order to further study the mechanism of increase of the CD22^{-/-} QM B cells, CD23 positive CD22^{+/+} QM B cells and CD22^{-/-} QM B cells were stained with CFSE, and C57BL/6Ly5.1 mice were transplanted with 4×10^6 cells of each group and immunized with NP-CGG. The proliferation of the Donor B cells, the class-switching and differentiation into the germinal center B cells was observed with the flow cytometry (Fig.3). The proliferation of Donor B cells was detected by a fluorescence intensity of CFSE of IgM⁺ B cells (B220⁺, Ly5.2⁺, IgM⁺) derived from the donor on the first, second and third days after the immunization. The class-switching was analyzed based on a fluorescence intensity of CFSE and IgG1 production of the B cells (B220⁺, Ly5.2⁺) derived

from the donor on the second and third days after the immunization. The differentiation into the germinal center B cells was analyzed based on a fluorescence intensity of CFSE and CD38 expression of the B cells (B220⁺, Ly5.2⁺) derived from the donor on the third day after the immunization.

[0050] On the first day after the immunization, cells that had decreased in CFSE fluorescence were not detected either in the CD22^{-/-} B cells or the CD22^{+/+} B cells, showing that no cell proliferation occurred. On the second day after the immunization, many cells that had decreased in CFSE fluorescence were observed in both the CD22^{-/-} B cells and the CD22^{+/+} B cells, showing that division of these cells had occurred. However, while the CD22^{+/+} B cells had suspended in the fourth cell division stage, the CD22^{-/-} B cells had already passed fifth or sixth cell division stage on the second day. Furthermore, while a peak of cell division of the CD22^{+/+} B cells occurred in the third cell division stage, that of the CD22^{-/-} B cells occurred in the fourth cell division stage, demonstrating that the cell division of the CD22^{-/-} B cells occurred rapidly in the early stage of the immune response. Furthermore, height of the peak in each cell division stage was higher in the CD22^{-/-} B cells than in the CD22^{+/+} B cells, showing the increase in survival of the CD22^{-/-} B cells, too. On the third day after the immunization, CFSE fluorescence intensity was decreased more significantly in the CD22^{-/-} B cells than in the CD22^{+/+} B cells, showing that the CD22^{-/-} B cells were dividing more rapidly than the CD22^{+/+} B cells were. Since no cell division was observed in the mice that had not immunized after the transplantation of the donor B cells, it was clear that the above cell proliferation was an antigen-specific reaction.

[0051] The emergence of the IgG1-positive cells and differentiation into the germinal center B cells were analyzed together with the detection of CFSE fluorescence. It was found that the IgG1-positive cells started to show up in the third cell division stage in both the CD22^{-/-} B cells and the CD22^{+/+} B cells. On the third day after the immunization, the IgG1-positive cells derived from the CD22^{-/-} B cells had increased about twice as much as that derived from the CD22^{+/+} B cells. As both the CD22^{-/-} B cells and the CD22^{+/+} B cells had shifted into the germinal center B cells (CD38^{lo}) from the seventh or eighth cell division stage, the germinal center B cells derived from the CD22^{-/-} B cells had increased about three times as much as that derived from the CD22^{+/+} B cells (Fig. 3B). No difference was found with respect to the number of cell division that is necessary for differentiation. It may be conceived that more IgG1-positive cells and germinal center B cells were generated from the CD22^{-/-} B cells than from the CD22^{+/+} B cells since the cell division was increased in the CD22^{-/-} B cells.

Example 3

Production of a large amount of antibody-producing cells and antibodies by CD22^{-/-} B cells in the early stage of the immune response

[0052] As a significant burst of plasmablast of the CD22^{-/-} B cells was observed in the section of spleen tissue in the early stage of the immune response, antibody production was studied with a mouse transplanted with the CD22^{-/-} B cells. The number of cells derived from the Donor B cells and producing an antibody specific for NP in the spleen of the recipient mice was determined by ELISPOT method on the third, fifth and seventh days after immunization (Fig.4A, B). From the third day after the immunization, about five times or more as many cells producing antibody of either IgM or IgG1 class were detected in the CD22^{-/-} B cells as in the CD22^{+/+} QM B cells. Even on the fifth day after the immunization, five times or more as many cells producing the antibodies of IgG1 class were detected in the CD22^{-/-} B cells as in the CD22^{+/+} QM B cells. However, the number of the cells derived from the CD22^{-/-} B cells and producing the antibody of either IgM or IgG1 class was decreased on the seventh day after the immunization. On the other hand, the number of the above cells had gradually increased in the case of the CD22^{+/+} B cells, showing difference in their kinetics. These results demonstrated that the antibody-producing cells emerged in the earlier stage in the CD22^{-/-} B cells than in the CD22^{+/+} B cells.

[0053] Titer of NP-specific antibody derived from Donor B cells in serum was then determined (Fig.4C,D). Being related to the increase of the antibody-producing cells in spleen, a specific IgM antibody has been detected from the third day from the immunization in the mice transplanted with the CD22^{-/-} B cells, and about three times as many specific antibodies were detected in the fifth day from the immunization in said mice transplanted with the CD22^{-/-} B cells as those in the mice transplanted with the CD22^{+/+} B cells. With respect to the kinetics, a peak was reached on the fifth day in the CD22^{-/-} B cells and on the seventh day in the CD22^{+/+} B cells, respectively, showing that the production of the antibodies occurred in the earlier stage in the case of the CD22^{-/-} B cells (Fig.4C). Furthermore, the production of the IgG1 class antibodies in the CD22^{-/-} B cells increased more significantly than in the CD22^{+/+} B cells, and the extension of such increase was higher than that with respect to the IgM class antibody. Thus, the IgG1 antibodies had been produced about five times as much in the CD22^{-/-} B cells as in the CD22^{+/+} B cells from the fifth day to tenth day after the immunization (Fig.4D). From the above results, it was clarified that the CD22^{-/-} B cells underwent differentiation into the antibody-producing cells in the earlier stage so as to produce a larger amount of antibodies than the CD22^{+/+} B cells did.

INDUSTRIAL APPLICABILITY

[0054] Once memory immune has been established by vaccine, for example, a rapid immune response including the production of antibodies will occur against invasion of pathogenic microorganisms so as to eliminate them and defend against infection prior to the occurrence of life-threatening infection. If medicine that can induce such an early immune response as the memory immune will is developed, it can be used to defend against infection caused by various microorganisms instead of vaccine. Since unlike in the case of vaccine it is not necessary to develop the above medicine for each microorganism, it will be especially useful for unknown infectious diseases. The present invention has demonstrated that the antibodies can be produced in the early stage by inhibiting or repressing the CD22 function, so that it will be possible to develop a universal medicine for defending against infection by targeting CD22.

Claims

1. A method for promoting an immune response comprising inhibiting CD22 function in B cells.
2. A method according to Claim 1, wherein the B cells are IgM and IgD positive naive B cells.
3. A method according to Claim 1 or 2, wherein the promotion of the immune response comprises the increase of proliferation, division and/or survival of the B cells in an early stage of the immune response.
4. A method according to Claim 3, wherein the B cells comprises IgG1⁺B cells, IgM⁺ B cells and germinal center B cells.
5. A method according to Claim 1 or 2, wherein the promotion of the immune response comprises class switching and/or differentiation of the B cells in an earlier stage.
6. A method according to Claim 5, wherein the class switching is that to the IgG1⁺ positive cells, and/or the differentiation is that into antibody-producing cells or the germinal center B cells.
7. A method according to any one of Claims 2 - 6, wherein the early stage in the immune response means from the third to fifth days after the immunization.
8. A method according to Claim 1, wherein the promotion of the immune response comprises increase in a number of the antibody-producing cells and in an amount of antibodies produced, and/or the production of antibodies in an earlier stage.

9. A method according to Claim 8, wherein the antibodies produced in the earlier stage are IgG and/or IgM.
10. A method according to any one of Claims 1 - 9, wherein the inhibition of the CD22 function is caused by its dysfunction due to deficiency or mutation of CD22 gene. 5
11. A method according to any one of Claims 1 - 9, wherein the inhibition of the CD22 function is caused by repression of the expression of CD22 gene. 10
12. A method according to Claim 11, wherein the repression of the expression of CD22 gene is done at a transcriptional level. 15
13. A method according to Claim 12, wherein gene knockdown by means of siRNA specific for CD22 gene is used. 20
14. A method according to Claim 13, wherein the siRNA is a double strand RNA consisting of an oligonucleotide having 21 - 23 bases and its complementary oligonucleotide, which comprises a continuous base sequence of a part of CD22 gene. 25
15. A method according to any one of Claims 1 - 9, wherein the inhibition of the CD22 function is related to the inhibition of signal transduction of B cell receptor. 30
16. A method according to any one of Claims 1 - 15, wherein the B cells and/or the CD22 or CD22 gene is derived from a mouse. 35
17. B cells whose CD22 function is inhibited.
18. B cells according to Claim 17, which are isolated from a knockout animal wherein CD22 gene is knocked out. 40
19. A method for the production of an antibody using the method according to any one of Claims 1 - 15 or the B cells whose CD22 function is inhibited according to Claim 17 or 18. 45
20. A method for the production according to Claim 19, wherein the antibody is specific for a T cell-dependent antigen. 50
21. A method for screening of a substance capable of promoting the immune response based on a change in the CD22 function in the B cells.
22. A method for screening according to Claim 21, comprising the following steps: 55


screened with the B cells;
 (e) a step of detecting the CD22 function in said cells; and
 (f) a step of selecting a substance which inhibits said function.

23. A method for screening according to any one of Claim 19 - 22, wherein the B cells and/or the CD22 or CD22 gene is derived from a mouse.

24. A screening kit used in a method according to any one of Claims 21 - 23.

(d) a step of contacting a substance to be

FIG. 1






FIG. 1




FIG. 2





FIG. 2




FIG. 2




FIG. 3





FIG. 4



INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP2007/074634

A. CLASSIFICATION OF SUBJECT MATTER

A61K45/00(2006.01)i, A61K39/00(2006.01)i, A61K39/395(2006.01)i, A61P37/04(2006.01)i, C12N5/10(2006.01)i, C12N15/09(2006.01)i, C12P21/08(2006.01)i, C12Q1/02(2006.01)i, G01N33/15(2006.01)i, G01N33/50(2006.01)i

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A61K45/00, A61K39/00, A61K39/395, A61P37/04, C12N5/10, C12N15/09, C12P21/08, C12Q1/02, G01N33/15, G01N33/50

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Jitsuyo Shinan Koho 1922-1996 Jitsuyo Shinan Toroku Koho 1996-2008
Kokai Jitsuyo Shinan Koho 1971-2008 Toroku Jitsuyo Shinan Koho 1994-2008

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

BIOSIS (STN), Cplus (STN), EMBASE (STN), MEDLINE (STN)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	O'KEEFE, Theresa L. et al., Hyperresponsive B Cells in CD22-Deficient Mice, Science, 1996, Vol.274, No.5288, P.798-801	16-24
X	KELM Soerge et al., The Ligand-binding Domain of CD22 Is Needed for Inhibition of the B Cell Receptor Signal, as Demonstrated by a Novel Human CD22-specific Inhibitor Compound, Journal of Experimental Medicine, 2002, Vol.195, No.9, p.1207-1213	16-24
A	WAKABAYASHI C. et al., A Distinct Signaling Pathway Used by the IgG-containing B Cell Antigen Receptor, Science, 2002, Vol.298, No.5602, p.2392-2395	16-24

Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier application or patent but published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search
06 March, 2008 (06.03.08)

Date of mailing of the international search report
18 March, 2008 (18.03.08)

Name and mailing address of the ISA/
Japanese Patent Office

Authorized officer

Facsimile No.

Telephone No.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/JP2007/074634

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.: 1 - 15
because they relate to subject matter not required to be searched by this Authority, namely:
Claims 1 to 15 pertain to methods for treatment of the human body by therapy.
2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

- Remark on Protest**
- the The additional search fees were accompanied by the applicant's protest and, where applicable, payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Non-patent literature cited in the description

- **Wakabayashi, C. et al.** *Science*, 2002, vol. 298 (5602), 2392-5 [0004] [0005]
- **Martin, S.W. ; C.C. Goodnow.** *Nat Immunol.*, 2002, vol. 3 (2), 182-8 [0004]
- **Nitschke, L. ; T. Tsubata.** *Trends Immunol*, 2004, vol. 25 (10), 543-50 [0005]
- **Tedder, T.F. et al.** *Annu Rev Immunol*, 1997, vol. 15, 481-504 [0005]
- Molecular cloning. Cold Spring Harbor Lab. Press, 2001 [0034]
- Current protocols in molecular biology. 1987 [0034]
- *Proc. Natl. Acad. Sci. USA*, 1990, vol. 87, 2264-2268 [0040]
- *Proc. Natl. Acad. Sci. USA*, 1993, vol. 90, 5873-5877 [0040]
- **Sambrook ; Maniatis.** *Molecular Cloning - A laboratory Manual.* Cold Spring Harbor Laboratory Press, 1989 [0042]
- **Ausbel, F. M. et al.** *Current Protocols in Molecular Biology.* John Wiley & sons, 1995 [0042]