



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 0 911 314 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
29.12.2004 Bulletin 2004/53

(51) Int Cl.7: **C07C 69/732**, C07C 59/54,
A61K 31/557, C07C 405/00

(21) Application number: **98308570.5**

(22) Date of filing: **20.10.1998**

(54) **Apoptosis inhibitor**
APOPTOSE-INHIBITOR
INHIBITEUR DE L'APOPTOSE

(84) Designated Contracting States:
CH DE FR GB IT LI

(30) Priority: **21.10.1997 JP 28891297**

(43) Date of publication of application:
28.04.1999 Bulletin 1999/17

(73) Proprietors:
• **Japan Science and Technology Agency**
Kawaguchi-shi, Saitama 332-0012 (JP)
• **Hazato, Atsuo**
Hino-shi, Tokyo (JP)

(72) Inventors:
• **Watanabe, Yasuyoshi**
Minoo-shi, Osaka (JP)
• **Suzuki, Masaaki**
Nagoya-shi, Aichi (JP)
• **Watanabe, Yumiko**
Minoo-shi, Osaka (JP)
• **Hazato, Atsuo**
Hino-shi, Tokyo (JP)

• **Sato, Takumi**
Suita-shi, Osaka (JP)

(74) Representative: **Calamita, Roberto**
Frank B. Dehn & Co.,
European Patent Attorneys,
179 Queen Victoria Street
London EC4V 4EL (GB)

(56) References cited:
EP-A- 0 229 844 **CA-A- 2 168 514**
US-A- 5 700 833

• **TRAUTH B C ET AL: "MONOCLONAL**
ANTIBODY-MEDIATED TUMOR REGRESSION
BY INDUCTION OF APOPTOSIS" SCIENCE, vol.
245, no. 4915, 21 July 1989, pages 301-305,
XP000749873
• **PATENT ABSTRACTS OF JAPAN vol. 017, no.**
567 (C-1120), 14 October 1993 & JP 05 163194 A
(TEIJIN LTD), 29 June 1993

EP 0 911 314 B1

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

[0001] The invention of the present application relates to the use of a compound in the manufacture of a composition for use as an apoptosis inhibitor and it relates to a novel apoptosis inhibitor containing isocarbacyclin derivative which has been known as a ligand specific to prostacyclin receptors of the central nervous type as an effective component and having an excellent inhibiting action to apoptosis of nerve cells, etc (CA 2168514 A).

[0002] Apoptosis is a kind of genetically programmed cell death and, morphologically, it takes place according to the following processes. Thus, aggregation of nucleus; cell contraction; vacuole formation; smoothing of cell surface; expansion of intercellular intervals; liberation of cells from the surroundings; fragmentation of cells (apoptosis corpuscles); and phagocytosis by macrophage, etc. It has been also known that, biochemically, nucleosome unit of DNA is fragmented by endonuclease activity into DNA of base length of 180-200 (Immunology Today, 7:115-119, 1986; Science, 245:301-305, 1989).

[0003] At present, it has been clarified that, in addition to physiological functions such as development/differentiation and turnover of normal tissues and cells, this apoptosis participates in certain diseases such as inflammation and reduction of lymphocytes caused by ischemic nerve cell death after cerebral infarction, etc., involution of carcinoma, action of radioactive ray and anticancer agent and infection of virus such as AIDS. Thus, development of drugs for controlling them (i.e., apoptosis inhibitors and apoptosis inducers) has been expected to give birth to drugs based upon new action mechanism in the broad areas including central nervous system as well as cancer, aging, etc.


[0004] With regard to substances and factors for inducing apoptosis, toxicity of neurotransmitters such as glucocorticoid and glutamic acid, irradiation of radioactive ray, NK cells, killer cells, tumor necrosis factor (TNF), cytokines such as lymphotoxin (LT), etc. have been reported. It has been also reported that cycloheximide which is a protein synthesis inhibitor or actinomycin D which is an RNA synthesis inhibitor induces apoptosis to human leukemia cells HL-60. Further, anti-Fas monoclonal antibody to Fas antigen which is a cell membrane molecule participating in apoptosis of immune cells has been prepared recently and various investigations have been conducted for application of anti-Fas antibody to pharmaceuticals.

[0005] On the other hand, substances which inhibit the interleukin 1 converting enzyme, basic fibroblast growth factor (bFGF), etc. have been reported as factors for inhibiting apoptosis. It has been also known that bc1-2 related gene products inhibit apoptosis and have a macrobiotic function for cells. However, all of those apoptosis inhibiting factors are derived from living organisms and, up to now, no apoptosis inhibitor obtained by industrial means such as chemical synthesis has been known yet.

[0006] On the other hand, during the course of investigation of physiological action of brain function in detail, the inventors of the present application have found various isocarbacyclins acting as specific ligands to prostacyclin receptor of the central nerve system and have filed patent applications already [Japanese Patent Application Nos. 051,589/96 (JP-A 245,498/96), 243,122/96 (JP-A 87608/98), 260,957/96 (JP-A 101610/98) and 160,320/97]. As a result of further investigation on the physiological activity of those substances, several of them have been found to show a significant apoptosis inhibiting effect.

[0007] Accordingly, an object of the invention of the present application is to offer a novel apoptosis inhibitor which can be manufactured by means of chemical synthesis in large quantities and also in low cost by a further development of the above-mentioned findings by the present inventors.

[0008] The first invention of this application is the use of a compound of the following formula (1)



EP 0 911 314 B1

wherein R_1 is a hydrocarbon chain having 1-6 carbon atom(s); and R_2 is a hydrogen atom or a protective group in the manufacture of a composition for use as an apoptosis inhibitor.


[0009] In a preferred embodiment the compound of formula (1) is 15R-16-(m-tolyl)-17,18,19,20-tetranorisocarbacyclin represented by the following formula (2) or methyl ester thereof.

5

10

15

20




[0010] The second invention is a 15-deoxy-isocarbacyclin derivative represented by the following formula (3) as an effective component.

25

30

35

40



wherein R_1 is a hydrocarbon chain having 1-6 carbon atom(s); and R_2 is a hydrogen atom or a protective group.


[0011] In the apoptosis inhibitor of the second invention, it is a preferred embodiment that the 15-Deoxy-isocarbacyclin derivative is 15-Deoxy-16-(m-tolyl)-17,18,19,20-tetranorisocarbacyclin represented by the following formula (4) or methyl ester thereof.

50

55

5

10



15

[0012] In the above-mentioned formulae (1) and (3), the protective group which constitutes R_2 is pharmaceutically acceptable salt, ester, etc. and its example is alkyl group which constitutes methyl ester, ethyl ester, etc.

[0013] Embodiments of the present invention will now be further illustrated as hereunder.

20

Figs. 1 and 2 are graphs showing the inhibitory effect of the tested samples on apoptosis of hippocampal neuron. Fig. 3 is a chemical formulae showing the synthetic process of Deoxy-TIC.

Fig. 4(A) and (B) are graphs showing uptake of labeled isocarbacyclin derivatives in whole brain and thalamus, respectively.

25

[0014] An example of the 15R-isocarbacyclin derivative which is an effective component of the first invention is 15R-16-(m-tolyl)-17,18,19,20-tetranorisocarbacyclin (hereinafter, may be referred to as "15R-TIC") or methyl ester thereof, and can be prepared by the method disclosed in JP-A 245,498/96 filed by the present inventors.

[0015] The 15-Deoxy-isocarbacyclin derivative which is an effective component of the second invention is, for example, 15-Deoxy-16-(m-tolyl)-17,18,19,20-tetranorisocarbacyclin (hereinafter, may be referred to as "Deoxy-TIC") or methyl ester thereof, and can be prepared by the method disclosed in the Japanese Patent Application No. 160,320/97 filed by the present inventors.

30

[0016] The compositions resulting from the use of the first invention and the compounds of the second invention can be made into pharmaceutical preparations containing any of those isocarbacyclin derivatives. In the case of manufacturing a pharmaceutical preparation for applying on central nervous system, it is preferred to manufacture the preparation with the methyl ester or the like of those isocarbacyclin derivatives. This is because that, as shown in the following Referential Example 2, such a methyl ester or the like is easily taken up into the brain and is converted to an isocarbacyclin derivative such as 15R-TIC or Deoxy-TIC in the brain thereby achieving an inhibiting effect on neural apoptosis. For the purpose of inhibiting the apoptosis in somatic cells, isocarbacyclin derivative such as 15R-TIC or Deoxy-TIC can be used as an effective component of a preparation. In addition to such an effective component, the apoptosis inhibitor of the present invention may further contain known apoptosis inhibiting substances.

35

40

[0017] The compositions resulting from the use of the first invention and the compounds of the second invention may be administered to human or other animals in a form of common pharmaceutical preparations. For example, it may be administered to human or other animals by means of intravenous injection, subcutaneous injection or oral administration.

45

[0018] The compositions resulting from the use of the first invention and the compounds of the second invention may be made into pharmaceutical preparations containing not less than $5 \mu\text{mol/kg}$ of isocarbacyclin derivative together with other components. Examples of said other components are those which are commonly used in the field of manufacture of pharmaceuticals such as diluting agents and vehicles including fillers, bulking agents, binders, moisturizers, disintegrating agents, surface-active agents and lubricants. With regard to a dosage form, various forms may be selected depending upon the object of therapy and representative examples thereof are tablets, pills, diluted powder, liquid, suspension, emulsion, granules, capsules, suppositories and injections (in liquid, suspension, etc.).

50

[0019] For example, in the case of preparing the injections, it is preferred that liquid, emulsion and suspension are sterilized and are isotonic to blood. Examples of the applicable diluent are water, ethyl alcohol, propylene glycol, ethoxylated isostearyl alcohol, polyoxylated isostearyl alcohol and polyoxyethylene sorbitan fatty acid esters. In that case, it is also possible to add salt, glucose, glycerol, etc. in such an amount that is satisfactory to prepare an isotonic solution. It is further possible to compound auxiliary solubilizers, buffers, analgetizing agent, etc. therewith.

55

[0020] If necessary, the pharmaceutical preparations in each of the above-mentioned forms may be further compounded with commonly used coloring agents, preservatives, perfumes, condiments, sweeteners, etc. Other pharmaceutically effective components may be compounded therewith as well.

[0021] The present invention will be illustrated in detail and in a specific manner by the following examples although

the present invention is not limited to those examples.

Example 1

5 **[0022]** Effect of 15R-TIC and Deoxy-TIC on hippocampal neuron apoptosis induced by a high-oxygen incubation was tested.

(1) Samples

10 **[0023]** Isocarbacyclin was a pure sample supplied from Teijin Co., Ltd. 15S-TIC was prepared in accordance with the method disclosed in Example 3 of JP-A 245,498/96 by using 15S-16-m-tolyl-17,18,19,20-tetra-nor-carbacyclin which is also disclosed in Example 2 of the same Publication JP-A 245,498/96. The 15R-TIC was prepared by the same method as disclosed in Example 3 of the Publication JP-A 245,498/96. bFGF was the commercially available one.

15 (2) Procedures

[0024] Hippocampal region was cut out from the fetal rat brain which was aseptically isolated from a pregnant rat (Wistar strain) of 20 days age and the hippocampal region was dispersed into neurons by shaking culture for 30 minutes in a PBS (Ca- and Mg-free) containing DL-cysteine (0.2 mg/ml), bovine serum albumin (0.2 mg/ml), glucose (5 mg/ml), DNase I (0.01%) and papain (9 units/ml). After that, each neuron was planted at the density of 5×10^5 cells/cm² on a 24-well plate (coated with polyethyleneimine) filled with a DME/F-12 medium (containing 5% equine serum and 5% fetal bovine serum) and incubated for two days in a 5% carbon dioxide gas incubator (9% oxygen). After that, the culture medium was substituted with an DME/F-12 medium (containing 5 μ g/ml of transferrin, 5 μ g/ml of insulin and 5 nM progesterone) wherefrom serum was removed, then each sample to be tested was added to the medium, after 25 30 minutes, transferred into 50% oxygen. After incubating in 50% oxygen for 48 hours, the survival rate of the neurons was measured. The neurons in 9% oxygen were also added each sample and the survival rate was measured.

[0025] With regard to the survival rate, the neurons were immobilized by 4% paraformaldehyde, stained with anti-MAP2 antibody and colored by DAB and the cells positive to MAP2 were counted from the 200-fold enlarged picture whereupon the survival rate was calculated.

30

(3) Results

[0026] Fig. 1 shows the survival rate of hippocampal neurons incubated in 9% or 50% oxygen where bFGF was added at a concentration of 50 nM and other samples were at 5 μ M. The number of survived neurons in 9% oxygen was used as a control. When isocarbacyclin or 15S-TIC was added, the survival rate of hippocampal neuron was greatly reduced by apoptosis induced by a high-oxygen condition as same as under the control condition. On the contrary, when 15R-TIC or Daoxy-TIC was added, an increase in the survival rate of neurons was observed to the same or even better extent as compared with the case of bFGF which was known as an apoptosis inhibitor in this experimental system.

40 **[0027]** Fig. 2 shows the dose-dependent effects of the tasted samples on hippocampal neural apoptosis. In this figure, the survival rate of neurons in 9% oxygen was used as a control, and the survival rate of neurons to which each sample being added was expressed as the mean \pm S.D. (n=4). As being clear from Fig. 2, while isocarbacyclin and 15S-TIC did not have any effect on neuroprotection in the range of concentrations tested, both 15R-TIC and Deoxy-Tic show the dose-dependent inhibitory effect on neural apoptosis, and the survival rate at 3000 nM was equivalent to the control. IC50 values were ca. 30 nM and 300 nM for Deoxy-TIC and 15R-TIC, respectively.

45 **[0028]** From these results, it has been now ascertained that 15R-TIC and Deoxy-TIC, the effective components of the apoptosis inhibitor of the present invention exhibit an excellent inhibiting effect to apoptosis of cells, particularly to apoptosis of neurons.

50 Example 2

[0029] Effect of 15R-TIC on hippocampal CA1 pyramidal neurons apoptosis induced by ischemic damage in mongolian gerbils was tested *in vivo*.

55 **[0030]** To examine whether 15R-TIC can prevent neuronal death *in vivo*, using an osmotic mini-pomp, 15R-TIC solution was continuously infused for 7 days (started 2 days before the transient ischemia and ended at 5 days) into the left lateral ventricle of gerbils that had been subjected to a transient forebrain ischemia for 3 minutes. Survival number of CA1 pyramidal neurons was assessed by counting neurons with cell bodies larger than 10 μ m in the Nissl-staining sections.

[0031] Administration of 15R-TIC blocked the loss of CA1 pyramidal neurons induced by forebrain ischemia. The number of neurons in the 15R-TIC-infused gerbils (984 ± 299 in a $20 \mu\text{m}$ thick section, $n=4$) was comparable to that in ischemia-free contralateral gerbils (1061 ± 220 , $n=4$). However, the number of CA1 pyramidal neurons in vehicle-infused gerbils (142 ± 28 , $n=4$) was significantly less than that of the control (without ischemic treatment) or 15R-TIC-infused gerbils. Pyramidal neurons in the vehicle-infused gerbils have progressively degenerated, with the nuclei having shrunk and the alignment of cells destroyed.

[0032] These results indicate that 15R-TIC acts as an effective neuronal survival-promoting factor *in vivo*, as well as *in vitro* data shown in Examples 1 and 2.

Referential Example 1

[0033] In accordance with the description of Example 1 of the Japanese Patent Application No. 160,320/97, Deoxy-TIC was synthesized as follows, in accordance with Fig 3.

<1> Synthesis of the aldehyde compound (6)

[0034] A solution of (triphenylphosphoranylidene) acetaldehyde (21.8 mg, $71.6 \mu\text{mol}$) in benzene (1.5 ml) was placed in a 10-ml Schlenk tube, then a solution of the above aldehyde compound (5) (23.0 mg, $65.3 \mu\text{mol}$) in benzene (1.5 ml) was added thereto and the mixture was heated to reflux for 20 hours. The reaction mixture was cooled, the solvent was evaporated therefrom and the residue was purified by means of an SiO_2 chromatography using 2:1 and 1:1 mixtures of hexane and ethyl acetate to give an aldehyde compound (6) (14.8 mg, 61%). The R_f value in the TLC was 0.55 (1:1 mixture of hexane and ethyl acetate).

<2> Synthesis of the carbonate compound (7)

[0035] A solution of the above aldehyde (6) (7.6 mg, $20.2 \mu\text{mol}$) in methanol (1.0 ml) was placed in a 10-ml round-bottom flask, then $\text{CeCl}_7 \cdot \text{H}_2\text{O}$ (10 mg, $27 \mu\text{mol}$) and NaBH_4 (2 mg, $53 \mu\text{mol}$) were added thereto and the mixture was stirred for five minutes. After that, ethyl acetate and water were added to the reaction mixture. Extraction was conducted for three times using ethyl acetate as a solvent. The organic phases were combined, dried over MgSO_4 and filtered and the filtrate was concentrated *in vacuo*. The resulting crude product was placed in a 10-ml round-bottom flask and dissolved in CH_2Cl_2 (2.0 ml). To this solution were added DMAP (37.0 mg, 0.303 mmol) and methyl chloroformate (0.015 ml, 0.194 mmol) followed by stirring for four hours. After that, an aqueous solution of NaHCO_3 was added thereto and the mixture was extracted with ethyl acetate. The organic phases were combined, dried over Na_2SO_4 and filtered and the filtrate was concentrated *in vacuo*. This was purified by means of an SiO_2 chromatography using a 4:1 mixture of hexane and ethyl acetate to give the above-mentioned carbonate compound (7) (8.0 mg, 91%). R_f in TLC was 0.42 (2:1 mixture of hexane and ethyl acetate).

<3> Synthesis of the adduct compound (9)

[0036] A solution of tris (dibenzylideneacetone)-dipalladium (O)-chloroform adduct (2.1 mg, $2.0 \mu\text{mol}$) and 1,2-bis(diphenylphosphino)ethane (1.6 mg, $4.0 \mu\text{mol}$) in THF (1.0 ml) was placed in a 20-ml Schlenk tube. To this solution was added a solution of the above carbonate compound (7) (8.0 mg, $18.3 \mu\text{mol}$) and disulfone (7.6 mg, $19.7 \mu\text{mol}$) in THF (1.0 ml) and the mixture was stirred for 15 minutes. The reaction mixture was poured into an aqueous solution of NH_4Cl followed by extracting with ethyl acetate. The organic phases were combined, dried over MgSO_4 and filtered and the filtrate was concentrated *in vacuo*. This was purified by means of an SiO_2 chromatography using 2:1, 1:1 and 1:2 mixtures of hexane and ethyl acetate to give a desired adduct (9) (9.6 mg, 70%). R_f in TLC was 0.27 (1:1 mixture of hexane and ethyl acetate).

<4> Synthesis of the compound (10): 15-Deoxy-16-(m-tolyl)-17,18,19,20-tetranorisocarbacyclin methyl ester

[0037] Mg (10 mg, 0.4 mmol) was placed in a 10-ml round-bottom flask and then a solution of the above adduct (9) (7.5 mg, $10.0 \mu\text{mol}$) in methanol (1.5 ml) was added followed by stirring for three hours. An aqueous solution of HCl (1N) was added to the reaction mixture followed by extracting with ethyl acetate. The organic phases were combined, dried over MgSO_4 and filtered and the filtrate was concentrated *in vacuo*. The resulting crude product was placed in a 10-ml round-bottom flask and dissolved in a 9:1 mixture (2.0 ml) of acetic acid and water. After stirring for 40 hours, ethyl acetate was added to the solution followed by washing with an aqueous solution of NaHCO_3 . The organic phase was dried over Na_2SO_4 and filtered and the filtrate was concentrated *in vacuo*. This was then purified by means of an SiO_2 chromatography using a 3:1 mixture of hexane and ethyl acetate to give a desired compound (10), *i.e.*, 15-Deoxy-

16-(m-tolyl)-17,18,19,20-tetranorisocarbacyclin methyl ester (2.2 mg, 58%). R_f in TLC was 0.6 (1:1 mixture of hexane and ethyl acetate).

<5> Synthesis of the compound (4): 15-Deoxy-16-(m-tolyl)-17,18,19,20-tetranorisocarbacyclin

[0038] A solution of the above compound (10) (1.0 mg, 2.6 μ mol) in methanol (0.5 ml) was placed in a 10-ml test tube, an aqueous solution (3N, 0.2 mol) of NaOH was added thereto and the mixture was stirred at room temperature for 12 hours. NaHCO₃ was added thereto and then ethyl acetate and water were further added. The pH was adjusted to 3. The organic phase was separated and the aqueous phase was extracted with ethyl acetate. The organic phases were combined, dried over MgSO₄ and filtered and the filtrate was concentrated *in vacuo*. This was purified by means of an SiO₂ (0.5 g) chromatography using a 10:1 mixture of CH₂Cl₂ and methanol to give the desired compound (4), the 15-Deoxy-16-(m-tolyl)-17,18,19,20-tetranorisocarbacyclin (0.9 mg, 94%). R_f in TLC was 0.39 (9:1 mixture of CH₂Cl₂ and methanol).

Referential Example 2

[0039] Uptake of the isocarbacyclin derivative into brain was measured by means of positron emission tomography (PET).

(1) Samples

[0040] ¹¹C-Labeled 15R-TIC methyl ester (RTA) was prepared by labeling the methyl group carbon at 16-m-tolyl group in the 15R-TIC methyl ester with ¹¹C. ¹¹C-Labeled methyl of 15R-TIC (RTC) was prepared by labeling the methyl group carbon at methyl group in the 15R-TIC methyl ester with ¹¹C. The 15R-TIC methyl ester was prepared by the same method as in Example 2 of JP-A 245498/96.

(2) Procedures

[0041] A tracer amount (not more than 0.2 μ g/kg) of the above mentioned sample was intravenously injected to an adult rhesus monkey (body weight being about 8 kg) and the behavior of the sample in the brain from the intravenous injection until 60 minutes thereafter was imaged by means of a PET device. ROI (region of interest) was calculated from said image and the uptake amount was calculated by the following formula.

$$\text{Uptake Amount} = \frac{[(\text{Calculated Weight of ROI in Brain}) / (\text{Body Weight})] \div [(\text{ROI Counts Accumulated in ROI}) / (\text{Administered ROI Counts})]}{1}$$

(3) Results

[0042] Fig. 4(A) shows the incorporated amount of each of the labeled compounds in the whole brain and Fig. 4(B) shows the incorporated amount in the thalamus. RTA and RTC showed equivalent pass through blood-brain-barrier (Fig. 4A and B), but RTA retained in the brain further than RTC, indicating the retention of deesterified ¹¹C-Labeled 15R-TIC in the brain. In other words, RTA could serve as a prodrug for 15R-TIC. It is clear from those results that the 15R-TIC methyl ester systemically administered has been confirmed to be efficiently transferred into the brain. Since the labeled compound remains in the brain for not shorter than 60 minutes after the administration, it has been suggested that the 15R-TIC methyl ester (a precursor to 15R-TIC) is converted to 15R-TIC in the brain to bind to prostacyclin receptor and that its apoptosis inhibiting action is achieved in the brain.

[0043] According to the present invention, a novel apoptosis inhibitor, of which ingredient is a compound available easily and in low cost by means of chemical synthesis, is provided. It has been clarified by PET study that the isocarbacyclin derivative, an effective component of the apoptosis inhibitor, passes through the blood-brain-barrier and acts on the central nervous system and, accordingly, the inhibitor is particularly effective for the therapy of various diseases caused by neural apoptosis.


Claims

1. The use of a compound of formula (1)

5

10

15



(1)


20

or a 15-deoxy-isocarbacyclin derivative of formula (3)

25

30

35



(3)

wherein R₁ is a hydrocarbon chain having 1-6 carbon atom(s); and R₂ is a hydrogen atom or a protective group in the manufacture of a composition for use as an apoptosis inhibitor.


40

2. The use as claimed in claim 1 wherein said compound of formula (1) is 15R-16-(m-tolyl)-17,18,19,20-tetranoriso-carbacyclin represented by the following formula (2):

45

50

55



(2)


or a methyl ester thereof.

3. The use as claimed in claim 1 wherein the 15-deoxy-isocarbacyclin derivative is 15-deoxy-16-(m-tolyl)-17,18,19,20-tetranorisocarbacyclin represented by the following formula (4):

5

10

15



(4)

20


or a methyl ester thereof.

4. A 15-deoxy-isocarbacyclin derivative represented by the following formula (3):

25

30

35



(3)

40

wherein R₁ is a hydrocarbon chain having 1-6 carbon atom(s); and R₂ is a hydrogen atom or a protective group.

5. A compound as claimed in claim 4, wherein the 15-deoxy-isocarbacyclin derivative is 15-deoxy-16-(m-tolyl)-17,18,19,20-tetranorisocarbacyclin represented by the following formula (4):

45


50

55

5

10

15



(4)

or a methyl ester thereof.

6. A compound as claimed in claim 4 or claim 5 for use in therapy.

7. A compound as claimed in claim 4 or claim 5 for use as an apoptosis inhibitor.

8. A pharmaceutical preparation comprising a compound of claim 4 or claim 5.


25 **Patentansprüche**

1. Verwendung einer Verbindung der Formel (1)

30

35

40



(1)

oder eines 15-Deoxyisocarbacyclin-Derivats der Formel (3)

45


50

55

5

10

15



(3)

worin R_1 für eine Kohlenwasserstoffkette mit 1 bis 6 Kohlenstoffatomen steht; und R_2 für ein Wasserstoffatom oder eine Schutzgruppe steht, zur Herstellung eines Mittels zur Verwendung als Apoptose-Hemmer.


20

2. Verwendung nach Anspruch 1, wobei es sich bei der Verbindung der Formel (1) um 15R-16-(m-Tolyl)-17,18,19,20-tetranorisocarbacyclin der nachstehenden Formel (2):

25

30

35



(2)

oder einen Methylester davon handelt.


40

3. Verwendung nach Anspruch 1, wobei es sich bei dem 15-Deoxyisocarbacyclin-Derivat um 15-Deoxy-16-(m-tolyl)-17,18,19,20-tetranorisocarbacyclin der nachstehenden Formel (4):

45

50


55



(4)


oder einen Methylester davon handelt.

4. 15-Deoxyisocarbacyclin-Derivat der nachstehenden Formel (3):



20 worin R_1 für eine Kohlenwasserstoffkette mit 1 bis 6 Kohlenstoffatomen steht; und R_2 für ein Wasserstoffatom oder eine Schutzgruppe steht.

- 25 5. Verbindung nach Anspruch 4, wobei es sich bei dem 15-Deoxyisocarbacyclin-Derivat um 15-Deoxy-16-(m-tolyl)-17,18,19,20-tetranorisocarbacyclin der nachstehenden Formel (4):




40 oder einen Methylester davon handelt.

- 45 6. Verbindung nach Anspruch 4 oder 5 zur therapeutischen Verwendung.
7. Verbindung nach Anspruch 4 oder 5 zur Verwendung als Apoptose-Hemmer.
8. Pharmazeutische Zusammensetzung, umfassend eine Verbindung nach Anspruch 4 oder 5.

Revendications

- 50 1. Utilisation d'un composé de formule (1)

5




10

15

ou d'un dérivé de 15-désoxy-isocarbacycline de formule (3)

20



25


30

où R_1 représente une chaîne hydrocarbure contenant de 1 à 6 atome(s) de carbone ; et R_2 représente un atome d'hydrogène ou un groupe protecteur pour la fabrication d'une composition pour une utilisation à titre d'inhibiteur de l'apoptose.

35

2. Utilisation selon la revendication 1, dans laquelle ledit composé de formule (1) est la 15R-16-(m-tolyl)-17,18,19,20-tétranorisocarbacycline représentée par la formule (2) suivante :

40



45


50

ou un ester méthylique de celle-ci.

55

3. Utilisation selon la revendication 1, dans laquelle le dérivé de 15-désoxy-isocarbacycline est la 15-désoxy-16-(m-tolyl)-17,18,19,20-tétranorisocarbacycline représentée par la formule (4) suivante :

5




10

ou un ester méthylique de celle-ci.

15

4. Dérivé de 15-désoxy-isocarbacycline représentée par la formule (3) suivante :

20



25


30

dans laquelle R_1 représente une chaîne d'hydrocarbure contenant de 1 à 6 atome(s) de carbone ; et R_2 représente un atome d'hydrogène ou un groupe protecteur.

35

5. Composé selon la revendication 4, dans lequel le dérivé de 15-désoxy-isocarbacycline est la 15-désoxy-16-(m-tolyl)-17,18,19,20-tétranisocarbacycline représentée par la formule (4) suivante :

40



45

50

ou un ester méthylique de celle-ci.

55

6. Composé selon la revendication 4 ou la revendication 5, pour une utilisation en thérapie.
7. Composé selon la revendication 4 ou la revendication 5, pour une utilisation à titre d'inhibiteur de l'apoptose.
8. Préparation pharmaceutique comprenant un composé selon la revendication 4 ou 5.

Fig. 1




Fig. 2




Fig. 3




Fig. 4

