



US 20030105615A1

(19) **United States**

(12) **Patent Application Publication** (10) **Pub. No.: US 2003/0105615 A1**

**Doi et al.** (43) **Pub. Date: Jun. 5, 2003**

(54) **METHOD AND A SYSTEM FOR PREDICTING PROTEIN FUNCTIONAL SITE, A METHOD FOR IMPROVING PROTEIN FUNCTION, AND A FUNCTION-MODIFIED PROTEIN**

Jan. 31, 1997 (JP) ..... 019249/1997  
Dec. 2, 1997 (JP) ..... 332100/1997  
Jan. 30, 1998 (JP) ..... 018699/1998

**Publication Classification**

(76) Inventors: **Hirofumi Doi**, Chiba (JP); **Hideaki Hiraki**, Tokyo (JP); **Akio Kanai**, Ibaraki (JP)

(51) **Int. Cl.<sup>7</sup>** ..... **G06G 7/48; G06G 7/58**  
(52) **U.S. Cl.** ..... **703/11**

Correspondence Address:  
**WENDEROTH, LIND & PONACK, L.L.P.**  
**2033 K STREET N. W.**  
**SUITE 800**  
**WASHINGTON, DC 20006-1021 (US)**

(57) **ABSTRACT**

(21) Appl. No.: **10/345,205**

(22) Filed: **Jan. 16, 2003**

**Related U.S. Application Data**

(63) Continuation of application No. 09/697,138, filed on Oct. 27, 2000, which is a continuation-in-part of application No. 09/355,486, filed on Sep. 20, 1999, now abandoned, filed as 371 of international application No. PCT/JP98/00430, filed on Feb. 2, 1998.

The present application provides a method for predicting the functional site of a protein using data of the entire proteins of an organism of which genome data or cDNA data is known. More specifically, the present application provides a method for predicting a protein functional site, comprising the steps of calculating the frequency of occurrence of an oligopeptide in the entire proteins, calculating the value of each amino-acid residue contributing to the frequency of occurrence as the representative value of the function, and predicting the protein functional site by using the representative value of function as an indicator. The present also provides a system for predicting a functional site for automatically performing said methods. Additionally, the present application provides a method for preparing a function-modified protein comprising subjecting the amino-acid residues composing the functional site identified by the method described above to artificial mutation, and a novel thermophilic DNA polymerase prepared by the method.

(30) **Foreign Application Priority Data**

Jan. 31, 1997 (JP) ..... 019248/1997

*Fig. 1*







Fig. 3



*Fig. 4*





Fig. 5



*Fig. 6*




Fig. 7







Fig. 8



*Fig. 9*



*Fig. 10*



*Fig. 11*





Fig. 12



*Fig. 13*




Fig. 14






Fig. 15




MJ0558 Position 325..346



MJ0558 Position 550..570

*Fig. 16*



*Fig. 17*

motif C

	550		570
MJ	AEKFGFKVLYIDTDGFYAIWK		
KOD	EEKYGFKVIYSDTDGFFATIP		
Pfu	EEKFGFKVLYIDTDGLYATIP		

Fig. 18




Fig. 19




Fig. 20





Fig. 21



*Fig. 22*







Fig. 24



**METHOD AND A SYSTEM FOR PREDICTING PROTEIN FUNCTIONAL SITE, A METHOD FOR IMPROVING PROTEIN FUNCTION, AND A FUNCTION-MODIFIED PROTEIN**

**TECHNICAL FIELD**

[0001] The present invention relates to a method for predicting a functional site of a protein, a system for predicting the function thereof, and a method for modifying the function of a protein and a function-modified protein. More specifically, the present invention relates to prediction of a functional site of a functionally unknown protein prepared by genome analysis or cDNA analysis, prediction of a novel function and a novel functional site of a protein with a known function, and prediction of a site on a protein to be modified for improving the function of the protein, and a protein with a modified function based on the prediction.

**BACKGROUND ART**

[0002] Following the progress of genome analysis and cDNA analysis of various organisms including pathogenic microorganisms, the number of novel genes whose functions are unknown is rapidly increasing, together with the number of proteins encoded by the genes. So far, the analysis of the nucleotide sequence of the whole genome of a microorganism, for example *Mycoplasma genitalium* (Fraser et al., *Science* 270, 397-403, 1995), *Haemophilus influenzae* (Fleischman et al., *Science* 269, 496-512, 1995), and *Methanococcus jannaschii* (Bult et al., *Science* 273, 1058-1073, 1996), has been completed, so that numerous novel proteins predicted from the genome sequence have been discovered. For humans and mice, the cDNA analysis is under way in combination with the genome analysis, which brings about the discovery of a great number of novel proteins.

[0003] In such circumstance, the prediction of the function of a functionally unknown protein or a functional site thereof has been a significant issue. If not only a novel protein but also a novel function or a novel functional site of a protein with a known function is discovered, whether or not these proteins are worth industrial or clinical application is possibly determined. Furthermore, such prediction of function possibly enables to prepare a modified protein with a further improved function.

[0004] Whether or not a protein encoded by a gene elucidated by genome analysis or cDNA analysis is novel or has a known function has been determined conventionally by searching the homology through protein databases such as Swiss-Prot. So as to predict a functional site, additionally, functionally identical proteins derived from various organisms are extracted from a protein database and are then subjected to alignment, to identify a region conserved in common to them and predict the conserved region as a functional site.

[0005] However, disadvantageously, such alignment method cannot be used if a protein obtained by genome analysis or cDNA analysis is an absolutely novel protein. Even if the protein has homology with known proteins in a protein database, the conserved region occupies most of the amino acid sequence of the protein in case that the protein is homologous to proteins derived from closely related organisms, so that it is impossible to predict the functional site. As to modification of protein, generally, the function of

a protein is potentially deteriorated irrespective of the fact that the function is known or unknown once the conserved region is modified, even if the functional site is predicted by alignment. Accordingly, the amino-acid residues outside the conserved region should be modified to improve the function. In other words, it is required to find a novel functional site in such protein to be modified. Using the conventional alignment method, disadvantageously, a novel functional site cannot be discovered or which amino-acid residue should be modified cannot be predicted.

[0006] Taking account of such circumstance, the present invention has been carried out. It is an object of the present invention to provide a novel method for predicting a functional site of a functionally unknown protein obtained by genome analysis or cDNA analysis.

[0007] In accordance with the present invention, furthermore, it is an object to provide a system for predicting the function.

[0008] In accordance with the present invention, still furthermore, it is an object to provide a method for predicting a novel functional site of a protein with an unknown function or with a known function and subjecting the functional site to mutation to prepare a modified protein.

[0009] Still furthermore, it is an object of the present invention to provide a protein with a function modified by the method described above.

**DISCLOSURE OF INVENTION**

[0010] A first aspect of the present invention proposed by the present application is a method for predicting a functional site of a protein derived from an organism "a" whose genome data, therefore the entire putative proteins, or cDNA data is known, which method comprises the steps of: (1) determining in the amino acid sequences of the entire proteins of the organism "a", the frequency of occurrence of each amino acid and the frequency of occurrence of individual oligopeptides produced by permutations of twenty amino acids, and determining the smallest length (n) of oligopeptides satisfying the following criteria;

[0011] among oligopeptides of length (n), the number of oligopeptides which occur once in the entire proteins is smaller than the number of oligopeptides which occur twice in the entire proteins; among oligopeptides of length (n+1), the number of oligopeptides which occur once in the entire proteins is larger than the number of oligopeptides which occur twice in the entire proteins,

[0012] (2) determining in the entire proteins of the organism "a", the frequency of occurrence of the following Aji-oligopeptide of length (n+1), which is a part of the amino acid sequence of the protein as a subject for predicting a functional site and contains the j-th amino-acid residue Aj from the amino terminal end (N-terminus) of the amino acid sequence (length of L) of the protein, on condition that the j-th amino-acid residue Aj is defined as the i-th residue Aji from the N-terminus of the Aji-oligopeptide;

[0013] Aji-oligopeptide:  $aj_1aj_2 \dots Aji \dots aj_naj_{n+1}$  (wherein,  $1 \leq i \leq n+1$ ;  $n+1 \leq j \leq L-n$ ;  $A_j = A_{ji}$ ; and Aji is the i-th residue of the oligopeptide; and moreover,  $aj_1 = A_{j-i+1}, \dots, aj_{n+1} = A_{j-i+(n+1)}$ ),

[0014] and determining in the entire proteins of the organism "a", the frequency of occurrence of the following Xji-oligopeptide of length (n+1);

[0015] Xji-oligopeptide: aj1aj2 . . . Xji . . . ajnaj(n+1) (wherein,  $1 \leq i \leq n+1$ ;  $n+1 \leq j \leq L-n$ ; and the i-th amino-acid residue Xji is any amino acid; and moreover,  $aj1=Aj-i+1, \dots, aj(n+1)=Aj-i+(n+1)$ )

[0016] (3) calculating ratio Yji of the frequency of occurrence of the Aji-oligopeptide to that of the Xji-oligopeptide,

[0017] (4) determining mean Yj of the Yji;

$$Y_j = \sum_{i=1}^{n+1} Y_{ji}/(n+1),$$

[0018] (5) determining functional value Zj of Yj;  $Z_j=f(Y_j)$  (Function f is a monotonously decreasing function or a monotonously increasing function),

[0019] and defining the Zj value as the representative value of the function of the j-th amino-acid residue Aj of the amino acid sequence (length of L), and

[0020] (6) repeating the steps (2) to (5) sequentially and determining the Zj value of each Aj of all the amino-acid residues at positions between  $n+1 \leq j \leq L-n$  in the amino acid sequence (length of L) of the protein, thereby predicting the degree of the involvement of each amino-acid residue in the function of the protein by using the dimension of the Zj value as an indicator.

[0021] A second aspect of the present invention is a method for predicting a functional site of a protein derived from the entire putative proteins of an organism "a" of which genome data or cDNA data is known, which method comprises the steps of:

[0022] (1) determining the frequency of occurrence of each amino acid and the frequency of occurrence of individual oligopeptides produced by permutations of twenty amino acids, in the amino acid sequences of the entire proteins of the organism "a",

[0023] (2) as to an appropriate protein of the organism "a",

[0024] (2') determining in the entire proteins of the organism "a", the frequency of occurrence of the following Aji-oligopeptide of given length of (n) ( $1 \leq n \leq M$ , provided that M is the smallest length of oligopeptides satisfying the criterion that all the oligopeptides of length M are at frequency 1 of the occurrence), which the Aji-oligopeptide is a part of the amino acid sequence of the protein and contains the j-th amino-acid residue Aj, on condition that the j-th amino-acid residue Aj is defined as the i-th residue Aji from the N-terminus of the Aji-oligopeptide ( $n \leq j \leq L-n+1$ );

[0025] Aji-oligopeptide: aj1aj2 . . . Aji . . . ajn (wherein,  $1 \leq n$ ;  $A_j=A_{ji}$ , and Aji is the i-th residue of the oligopeptide; and moreover,  $aj1=A_{j-i+1}, \dots, ajn=A_{j-i+n}$ ),

[0026] and determining in the entire proteins of the organism "a", the frequency of occurrence of the following Xji-oligopeptide of length (n) corresponding to the length of Aji-oligopeptide;

[0027] Xji-oligopeptide: aj1aj2 . . . Xji . . . ajn (wherein,  $1 \leq i \leq n$ ; and the i-th residue Xji is any amino acid; and moreover,  $aj1=A_{j-i+1}, \dots, ajn=A_{j-i+n}$ ),

[0028] (3) calculating ratio Yji of the frequency of occurrence of the Aji-oligopeptide to that of the Xji-oligopeptide,

[0029] (4) determining mean Y(j,n) of the Yji;

$$Y(j, n) = \sum_{i=1}^n Y_{ji}/n,$$

[0030] (5) determining functional value Z(j,n) of Y(j,n);  $Z(j,n)=-\log(Y(j,n))$ ,

[0031] (6) repeating the steps (2') to (5) sequentially and determining the Z(j,n) value of each amino-acid residue Aj at position j ( $n \leq j \leq L-n+1$ ) in the amino acid sequence (length of L),

[0032] (7) sequentially repeating the steps (2) to (6) for the entire proteins of the organism "a", thereby determining the distribution of the Z(j,n) value of each amino-acid residue in the entire proteins, and the Z(j,n) values are classified into twenty according to the twenty amino acids, and then determining mean Av(Aa) of the Z(j,n) values for each amino acid Aa and the standard deviation Sd(Aa) of the distribution thereof, on the basis of the distribution, to determine function g to the j-th amino-acid residue Aj of a protein for normalizing the difference in distribution due to the species of amino-acid residues;

[0033]  $g=g(Z(j,n), A_j)=[Z(j,n)-Av(Aa)]/Sd(Aa)$  (provided that  $A_j=A_a$ ; and  $1 \leq n \leq M$ ),

[0034] (8) determining value D(j,n) of the function g of each Aj of all the amino-acid residues at position j ( $n \leq j \leq L-n+1$  and  $1 \leq n \leq M$ ) of a protein in the entire proteins as recovered in the step (7);

[0035]  $D(j,n)=g(Z(j,n), A_j)$ , and

[0036] (9) defining the representative value of the function of the j-th amino-acid residue in the amino acid sequence (length of L) as functional value Wj of the Z(j,n) and D(j,n);

[0037]  $W_j=h(Z(j,1), Z(j,2), \dots, Z(j,M), D(j,1), D(j,2), \dots, D(j,M))$  thereby predicting the degree of the involvement of each amino-acid residue in the function of the protein by using the dimension of the Wj value as an indicator.

[0038] A third aspect of the present invention is a system for automatically conducting the method according to claim 1, at least comprising the following units (a) to (g);

[0039] (a) an outer memory unit memorizing the amino acid sequence data of the entire putative proteins derived from organism "a" of which genome data or cDNA data is known, as well as an existing protein data base,

[0040] (b) a calculation/memory unit, composed of CPU calculating the frequency of occurrence of each amino acid and the frequency of occurrence of individual oligopeptides produced by permutations of twenty amino acids, in the

amino acid sequences of the entire proteins from the organism "a", and a memory unit having the memory of the calculation results,

[0041] (c) a calculation/memory unit, composed of CPU calculating the smallest length (n) of oligopeptide satisfying the following criteria among the individual oligopeptides of which the frequencies of the occurrences being memorized in the unit (b);

[0042] among oligopeptides of length (n), the number of oligopeptides which occur once in the entire proteins is smaller than the number of oligopeptides which occur twice in the entire proteins; among oligopeptides of length (n+1), the number of oligopeptides which occur once in the entire proteins is larger than the number of oligopeptides which occur twice in the entire proteins, and a memory unit having the memory of the length (n),

[0043] (d) a calculation/memory unit, composed of CPU determining in the entire proteins of the organism "a", the frequency of occurrence of the following Aji-oligopeptide of length (n+1), which is a part of the amino acid sequence of the protein as a subject for predicting a functional site and contains the j-th amino-acid residue Aj from the amino terminal end (N-terminus) of the amino acid sequence (length of L) of the protein, on condition that the j-th amino-acid residue Aj is defined as the i-th residue Aji from the N-terminus of the Aji-oligopeptide;

[0044] Aji-oligopeptide: aj1aj2 . . . Aji . . . ajnaj(n+1) (wherein,  $1 \leq i \leq n+1$ ;  $n+1 \leq j \leq L-n$ ;  $A_j = A_{ji}$ ; and Aji is the i-th residue of the oligopeptide; and moreover,  $aj1 = A_{j-i+1}, \dots, aj(n+1) = A_{j-i+(n+1)}$ ),

[0045] and determining in the entire proteins of the organism "a", the frequency of occurrence of the following Xji-oligopeptide of length (n+1);

[0046] Xji-oligopeptide: aj1aj2 . . . Xji . . . ajnaj(n+1) (wherein,  $1 \leq n+1$ ;  $n+1 \leq j \leq L-n$ ; and the i-th amino-acid residue Xji is any amino acid; and moreover,  $aj1 = A_{j-i+1}, aj(n+1) = A_{j-i+(n+1)}$ ), and a memory unit having the memory of the calculation results,

[0047] (e) a calculation/memory unit, composed of CPU determining ratio Yji of the frequency of occurrence of the Aji-oligopeptide to that of the Xji-oligopeptide, and a memory unit having the memory of Yji,

[0048] (f) a calculation/memory unit, composed of CPU determining mean Yj of the Yji;

$$Y_j = \sum_{i=1}^{n+1} Y_{ji} / (n+1),$$

[0049] and a memory unit having the memory of Yj, and

[0050] (g) a calculation/memory unit, composed of CPU determining functional value Zj of Yj;

[0051]  $Z_j = f(Y_j)$

[0052] (Function f is a monotonously decreasing function or a monotonously increasing function),

[0053] and a memory unit having the memory of Zj.

[0054] A fourth aspect of the present invention is a system for automatically conducting the method according to claim 3, at least comprising the following units (a) to (i);

[0055] (a) an outer memory unit memorizing the amino acid sequence data of the entire putative proteins of the organism "a" of which genome data or cDNA data is known, as well as an existing protein data base,

[0056] (b) a calculation/memory unit, composed of CPU calculating in the amino acid sequences of the entire proteins of the organism "a", the frequency of occurrence of each amino acid and the frequency of occurrence of individual oligopeptides produced by permutations of twenty amino acids, and a memory unit having the memory of the calculation results,

[0057] (c) a calculation/memory unit, composed of CPU determining in the entire proteins of the organism "a", the frequency of occurrence of the following Aji-oligopeptide of given length of (n) ( $1 \leq n \leq M$ , provided that M is the smallest length of oligopeptides satisfying the criterion that all the oligopeptides of length M are at frequency 1 of the occurrence), which the Aji-oligopeptide is a part of the amino acid sequence of the protein and contains the j-th amino-acid residue Aj, on condition that the j-th amino-acid residue Aj is defined as (or identical to?) the i-th residue Aji from the N-terminus of the Aji-oligopeptide ( $n \leq j \leq L-n+1$ );

[0058] Aji-oligopeptide: aj1aj2 . . . Aji . . . ajn (wherein,  $1 \leq i \leq n$ ;  $A_j = A_{ji}$ , and Aji is the i-th residue of the oligopeptide; and moreover,  $aj1 = A_{j-i+1}, \dots, ajn = A_{j-i+n}$ ), and determining in the entire proteins of the organism "a", the frequency of occurrence of the following Xji-oligopeptide of length (n) corresponding to the length of Aji-oligopeptide;

[0059] Xji-oligopeptide: aj1aj2 . . . Xji . . . ajn (wherein,  $1 \leq i \leq n$ ; and the i-th residue Xji is any amino acid; and moreover,  $aj1 = A_{j-i+1}, \dots, ajn = A_{j-i+n}$ ),

[0060] and a memory unit memorizing the calculation results,

[0061] (d) a calculation/memory unit, composed of CPU calculating ratio Yji of the frequency of occurrence of the Aji-oligopeptide to that of the Xji-oligopeptide, and a memory unit having the memory of the Yji,

[0062] (e) a calculation/memory unit, composed of CPU determining mean Y(j,n) of the Yji;

$$Y(j, n) = \sum_{i=1}^n Y_{ji} / n,$$

[0063] and a memory unit having the memory of Y(j,n),

[0064] (f) a calculation/memory unit, composed of CPU determining functional value Z(j,n) of Y(j,n);

[0065]  $Z(j,n) = -\log(Y(j,n))$ ,

[0066] and a memory unit having the memory of Z(j,n),

[0067] (g) a calculation/memory unit, composed of CPU determining the distribution of the Z(j,n) value of each

amino-acid residue in the entire proteins, and the  $Z(j,n)$  values are classified into twenty according to the twenty amino acids, and then determining mean  $Av(Aa)$  of the  $Z(j,n)$  values for each amino acid  $Aa$  and the standard deviation  $Sd(Aa)$  of the distribution thereof, on the basis of the distribution, to determine function  $g$  to the  $j$ -th amino-acid residue  $A_j$  of a protein for normalizing the difference in distribution due to the species of amino-acid residues;

$$[0068] \quad g = g(Z(j,n), A_j) = [Z(j,n) - Av(Aa)] / Sd(Aa)$$

[0069] (wherein,  $A_j = Aa$ ; and  $1 \leq n \leq M$ )

[0070] and a memory unit having the memory of  $g$ ,

[0071] (h) a calculation/memory unit, composed of CPU determining value  $D(j,n)$  of function  $g$  memorized in the unit ( $g$ ) concerning each of all the amino-acid residues  $A_j$  at position  $j$  ( $n \leq j \leq L-n+1$ ) in the amino acid sequence (length of  $L$ );

$$[0072] \quad D(j,n) = g(Z(j,n), A_j)$$

[0073] and a memory unit having the memory of the  $D(j,n)$  value, and (i) a calculation/memory unit, composed of a calculation unit determining appropriate functional value  $W_j$  of the  $Z(j,n)$  and  $D(j,n)$  of each amino-acid residue in the amino acid sequence;

$$[0074] \quad W_j = h(Z(j,1), Z(j,2), \dots, Z(j,M), D(j,1), D(j,2), \dots, D(j,M))$$

[0075] and a memory unit having the memory of the  $W_j$  value.

[0076] A fifth aspect of the present invention is a method for modifying the known function of protein "A" derived from the entire proteins of organism "a" of which genome data or cDNA data has been known, which method comprises the steps of:

[0077] (1) extracting a protein closely related to the protein "A" from an existing protein data base and subjecting the proteins to alignment,

[0078] (2) determining in the amino acid sequences of the entire proteins of the organism "a", the frequency of occurrence of each amino acid and the frequency of occurrence of individual oligopeptides produced by permutations of twenty amino acids, and determining the smallest length ( $n$ ) of oligopeptides satisfying the following criteria;

[0079] among oligopeptides of length ( $n$ ), the number of oligopeptides which occur once in the entire proteins is smaller than the number of oligopeptides which occur twice in the entire proteins; among oligopeptides of length ( $n+1$ ), the number of oligopeptides which occur once in the entire proteins is larger than the number of oligopeptides which occur twice in the entire proteins,

[0080] (3) determining in the entire proteins of the organism "a", the frequency of occurrence of the following Aji-oligopeptide of length ( $n+1$ ), which is a part of the amino acid sequence of the protein as a subject for predicting a functional site and contains the  $j$ -th amino-acid residue  $A_j$  from the amino terminal end (N-terminus) of the amino acid sequence (length of  $L$ ) of the protein, on condition that the  $j$ -th amino-acid residue  $A_j$  is defined as the  $i$ -th residue  $A_{ji}$  from the N-terminus of the Aji-oligopeptide;

[0081] Aji-oligopeptide:  $a_{j1}a_{j2} \dots a_{ji} \dots a_{jn}a_{j(n+1)}$  (wherein,  $1 \leq i \leq n+1$ ;  $n+1 \leq j \leq L-n$ ;  $A_j = A_{ji}$ ; and  $A_{ji}$  is the  $i$ -th residue of the oligopeptide; and moreover,  $a_{j1} = A_{j-i+1}, \dots, a_{j(n+1)} = A_{j-i+(n+1)}$ ), and determining in the entire proteins of the organism "a", the frequency of occurrence of the following Xji-oligopeptide of length ( $n+1$ );

[0082] Xji-oligopeptide:  $a_{j1}a_{j2} \dots X_{ji} \dots a_{jn}a_{j(n+1)}$  (wherein,  $1 \leq i \leq n+1$ ;  $n+1 \leq j \leq L-n$ ; and the  $i$ -th amino-acid residue  $X_{ji}$  is any amino acid; and moreover,  $a_{j1} = A_{j-i+1}, \dots, a_{j(n+1)} = A_{j-i+(n+1)}$ ),

[0083] (4) calculating ratio  $Y_j$  of the frequency of occurrence of the Aji-oligopeptide to that of the Xji-oligopeptide,

[0084] (5) determining mean  $Y_j$  of the  $Y_j$ ;

$$Y_j = \sum_{i=1}^{n+1} Y_{ji} / (n+1),$$

[0085] (6) determining functional value  $Z_j$  of  $Y_j$ ;  $Z_j = f(Y_j)$  (Function  $f$  is a monotonously decreasing function or a monotonously increasing function), and defining the  $Z_j$  value as the representative value of the function of the  $j$ -th amino-acid residue  $A_j$  of the amino acid sequence (length of  $L$ ) of the protein "A",

[0086] (7) sequentially repeating the steps (3) to (6) and determining the  $Z_j$  value of each of all the amino-acid residues at positions between  $n+1 \leq j \leq L-n$  in the amino acid sequence (length of  $L$ ) of the protein "A",

[0087] (8) selecting at least one amino-acid residue to be subjected to mutation from the amino acid sequence (length of  $L$ ) of the protein "A" on the basis of the alignment data carried out in the step (1), sequentially repeating the steps (3) to (6) for variant amino-acid residues in various mutated (or mutant?) amino acid sequences where the selected amino-acid residue has been mutated into another amino-acid residue, to determine the  $Z_j$  value of the variant amino-acid residues,

[0088] (9) selecting a mutated amino acid sequence wherein the  $Z_j$  value of the variant amino-acid residue as determined in the step (8) is larger or smaller than the  $Z_j$  value of the wild type amino-acid residue as determined in the step (7), and

[0089] (10) preparing a modified gene encoding the modified amino acid sequence from the protein "A" gene, and producing the modified protein as the expression product of the gene.

[0090] A sixth aspect of the present invention is a method for modifying the function of protein "B" derived from an organism "b" of which genome data or cDNA data has been unknown, which method comprises the steps of:

[0091] (1) extracting protein "A" most closely related to protein "B" from the entire proteins of organism "a" of which genome data or cDNA data being known and subjecting the protein to alignment, or extracting a protein closely related to protein "B" from an existing protein data base to subject the protein to alignment,

**[0092]** (2) determining in the amino acid sequences of the entire proteins of the organism "a", the frequency of occurrence of each amino acid and the frequency of occurrence of individual oligopeptides produced by permutations of twenty amino acids, and determining the smallest length (n) of oligopeptides satisfying the following criteria;

**[0093]** among oligopeptides of length (n), the number of oligopeptides which occur once in the entire proteins is smaller than the number of oligopeptides which occur twice in the entire proteins; among oligopeptides of length (n+1), the number of oligopeptides which occur once in the entire proteins is larger than the number of oligopeptides which occur twice in the entire proteins,

**[0094]** (3) determining in the entire proteins of the organism "a", the frequency of occurrence of the following Aji-oligopeptide of length (n+1), which is a part of the amino acid sequence of the protein as a subject for predicting a functional site and contains the j-th amino-acid residue Aj from the amino terminal end (N-terminus) of the amino acid sequence (length of L) of the protein, on condition that the j-th amino-acid residue Aj is defined as the i-th residue Aji from the N-terminus of the Aji-oligopeptide;

**[0095]** Aji-oligopeptide: aj1aj2 . . . Aji . . . ajnaj(n+1) (wherein,  $1 \leq i \leq n+1$ ;  $n+1 \leq j \leq L-n$ ;  $A_j = A_{ji}$ ; and Aji is the i-th residue of the oligopeptide; and moreover,  $aj1 = A_{j-i+1}, \dots, aj(n+1) = A_{j-i+(n+1)}$ ), and determining in the entire proteins of the organism "a", the frequency of occurrence of the following Xji-oligopeptide of length (n+1);

**[0096]** Xji-oligopeptide: aj1aj2 . . . Xji . . . ajnaj(n+1) (wherein,  $1 \leq i \leq n+1$ ;  $n+1 \leq j \leq L-n$ ; and the i-th amino-acid residue Xji is any amino acid; and moreover,  $aj1 = A_{j-i+1}, \dots, aj(n+1) = A_{j-i+(n+1)}$ ),

**[0097]** (4) calculating ratio Yji of the frequency of occurrence of the Aji-oligopeptide to that of the Xji-oligopeptide,

**[0098]** (6) determining mean Yj of the Yji;

$$Y_j = \sum_{i=1}^{n+1} Y_{ji} / (n+1),$$

**[0099]** (6) determining functional value Zj of Yj;

**[0100]**  $Z_j = f(Y_j)$

**[0101]** (Function f is a monotonously decreasing function or a monotonously increasing function),

**[0102]** and defining the Zj value as the representative value of the function of the j-th amino-acid residue Aj of the amino acid sequence (length of L) of the protein "A",

**[0103]** (7) sequentially repeating the steps (3) to (6) and determining the Zj value of each of all the amino-acid residues at positions between  $n+1 \leq j \leq L-n$  in the amino acid sequence (length of L),

**[0104]** (8) selecting at least one amino-acid residue to be subjected to mutation from the amino acid sequence (length of L) of the protein "A" on the basis of the alignment data carried out in the step (1), sequentially repeating the steps (3) to (6) for variant amino-acid residues in various mutated amino acid sequences where the selected amino-acid residue has been mutated into another amino-acid residues, to determine the Zj value of the variant amino-acid residues,

**[0105]** (9) selecting the mutation position and the mutated amino-acid residue wherein the Zj value of the variant amino-acid residue as determined in the step (8) is larger or smaller than the Zj value of the wild type amino-acid residue as determined in the step (7), and

**[0106]** (10) preparing a modified gene encoding the modified amino acid sequence having the mutated amino-acid residue at the position from the protein "B" gene, and producing the modified protein as the expression product of the gene.

**[0107]** A seventh aspect of the present invention is a method for modifying the known function of protein "A" derived from the entire proteins of organism "a" of which genome data or cDNA data has been known, which method comprises the steps of:

**[0108]** (1) extracting proteins closely related to the protein "A" from an existing protein data base and subjecting the proteins to alignment,

**[0109]** (2) determining the frequency of occurrence of each amino acid and the frequency of occurrence of individual oligopeptides produced by permutations of twenty amino acids, in the amino acid sequences of the entire proteins of the organism "a"

**[0110]** (3) as to an appropriate protein of the organism "a", (3') determining in the entire proteins of the organism "a", the frequency of occurrence of the following Aji-oligopeptide of given length of (n) ( $1 \leq n \leq M$ , provided that M is the smallest length (n) of oligopeptides satisfying the criterion that all the oligopeptides of length M are at frequency 1 of the occurrence), which the Aji-oligopeptide is a part of the amino acid sequence of the protein and contains the j-th amino-acid residue Aj, on condition that the j-th amino-acid residue Aj is defined as the i-th residue Aji from the N-terminus of the Aji-oligopeptide ( $n \leq j \leq L-n+1$ );

**[0111]** Aji-oligopeptide: aj1aj2 . . . Aji . . . ajn (wherein,  $1 \leq i \leq n$ ;  $A_j = A_{ji}$ , and Aji is the i-th residue of the oligopeptide; and moreover,  $aj1 = A_{j-i+1}, \dots, ajn = A_{j-i+n}$ ),

**[0112]** and determining in the entire proteins of the organism "a", the frequency of occurrence of the following Xji-oligopeptide of length (n) corresponding to the length of Aji-oligopeptide;

**[0113]** Xji-oligopeptide: aj1aj2 . . . Xji . . . ajn (wherein,  $1 \leq i \leq n$ ; and the i-th residue Xji is any amino acid; and moreover,  $aj1 = A_{j-i+1}, \dots, ajn = A_{j-i+n}$ ),

[0114] (4) calculating ratio  $Y_{ji}$  of the frequency of occurrence of the  $A_{ji}$ -oligopeptide to that of the  $X_{ji}$ -oligopeptide,

[0115] (5) determining mean  $Y(j,n)$  of the  $Y_{ji}$ ;

$$Y(j,n) = \sum_{i=1}^n Y_{ji}/n,$$

[0116] (6) determining functional value  $Z(j,n)$  of  $Y(j,n)$ ;  $Z(j,n) = -\log(Y(j,n))$ ,

[0117] (7) repeating the steps (3') to (6) sequentially and determining the  $Z(j,n)$  value of each amino-acid residue  $A_j$  at position  $j$  ( $n \leq j \leq L-n+1$ ) in the amino acid sequence (length of  $L$ ),

[0118] (8) sequentially repeating the steps (3) to (7) for the entire proteins of the organism "a", thereby determining the distribution of the  $Z(j,n)$  value of each amino-acid residue in the entire proteins, and the  $Z(j,n)$  values are classified into twenty according to the twenty amino acids, and then determining mean  $Av(Aa)$  of the  $Z(j,n)$  values for each amino acid  $Aa$  and the standard deviation  $Sd(Aa)$  of the distribution thereof, on the basis of the distribution, to determine function  $g$  to the  $j$ -th amino-acid residue  $A_j$  of a protein for normalizing the difference in distribution due to the species of amino-acid residues;

[0119]  $g = g(Z(j,n), A_j) = [Z(j,n) - Av(Aa)] / Sd(Aa)$  (provided that  $A_j = Aa$ ; and  $1 \leq n \leq M$ ),

[0120] (9) determining value  $D(j,n)$  of the function  $g$  of each  $A_j$  of all the amino-acid residues at position  $j$  ( $n \leq j \leq L-n+1$  and  $1 \leq n \leq M$ ) of a protein in the entire proteins as recovered in the step (8);

[0121]  $D(j,n) = g(Z(j,n), A_j)$ , and

[0122] (10) defining the representative value of the function of the  $j$ -th amino-acid residue in the amino acid sequence (length of  $L$ ) as functional value  $W_j$  of the  $Z(j,n)$  and  $D(j,n)$ ;

[0123]  $W_j = h(Z(j,1), Z(j,2), \dots, Z(j,M), D(j,1), D(j,2), D(j,M))$

[0124] (11) sequentially repeating the steps (3) to (10), to determine the individual  $W_j$  values of all the amino-acid residues at position  $n \leq j \leq L-n+1$  in the amino acid sequence (length of  $L$ ),

[0125] (12) selecting at least one amino-acid residue to be subjected to mutation on the basis of the alignment data carried out in the step (1) from the amino acid sequence (length of  $L$ ) of the protein "A", and sequentially repeating the steps (3) to (10) for variant amino-acid residues in various mutated amino acid sequences where the selected amino-acid residue has been mutated into another amino-acid residue, to determine the  $W_j$  value of the variant amino-acid residue,

[0126] (13) selecting a mutated amino acid sequence wherein the  $W_j$  value of the variant amino-acid residue as determined in the step (12) is larger or smaller than the  $W_j$  value of the wild type amino-acid residue as determined in the step (10), and

[0127] (14) preparing a modified gene encoding the modified amino acid sequence from the protein "A" gene, and producing the modified protein as the expression product of the gene.

[0128] An eighth aspect of the present invention is a protein of which function is artificially modified by the methods for modifying the function of protein set forth above.

[0129] One embodiment of the function-modified protein is a thermophilic DNA polymerase, prepared by artificially modifying the amino acid sequence of Pfu DNA polymerase so that the elongation of synthesized DNA chain might not be terminated intermediately during the catalysis for the synthesis of a DNA chain complementary to a single-stranded DNA, and more specifically, the thermophilic DNA polymerase is one comprising the amino acid sequence of SQ ID No.1. In association with this thermophilic DNA polymerase, the present application provides a DNA sequence encoding the amino acid sequence of SQ ID No.1 and a recombinant vector carrying the DNA sequence. Such recombinant vector includes recombinant plasmid pDP320 carried on *Escherichia coli* HMS174 (DE3)/pDP320 (FERM P-16052). Still furthermore, in accordance with the present invention, it is provided a method for preparing this thermophilic DNA polymerase, comprising culturing a cell transformed with an expression vector carrying the DNA sequence and isolating and purifying the objective enzyme generated in a culture medium.

[0130] Another embodiment of the function-modified protein is a thermophilic DNA polymerase, prepared by artificially modifying the amino acid sequence of Pfu DNA polymerase so that the synthesized DNA chain might be more elongated during the catalysis for the synthesis of a DNA chain complementary to a single-stranded DNA, and more specifically, the thermophilic DNA polymerase is one comprising the amino acid sequence of SQ ID No.6 or a DNA polymerase comprising the amino acid sequence of SQ ID No.7. In association with this thermophilic DNA polymerase, the present application provides a DNA sequence encoding the amino acid sequence of SQ ID No.6 or 7, and a recombinant vectors carrying such DNA sequences, respectively. As such vectors, there are provided recombinant plasmid pDP5b17 carried on *Escherichia coli* HMS174 (DE3)/pDP5b17 (FERM BP-6189) (vector carrying the DNA sequence encoding the amino acid sequence of SQ ID No.1), and recombinant plasmid pDP5C4 carried on *Escherichia coli* HMS174 (DE3)/pDP5C4 (FERM BP-6190) (vector carrying the DNA sequence encoding the amino acid sequence of SQ ID No.1). Still furthermore, it is provided a method for producing the DNA polymerase, comprising culturing a cell transformed with an expression vector carrying the DNA sequence and isolating and purifying the objective enzyme produced in a culture medium.

[0131] The method for predicting a protein functional site in accordance with the first aspect of the present invention has been established on what will be described below. More specifically, protein is composed of a sequence of twenty amino acids, but the sequence is not random. Hence, the frequency of occurrence of a specific oligopeptide as a partial amino acid sequence in the entire proteins encoded by genome derived from an appropriate organism species is not constant, but some oligopeptides occur at high frequen-

cies in various proteins while other oligopeptides rarely occur therein. It is recognized that among them, oligopeptides highly frequently occurring in common to various proteins do not have any potency to determine the uniqueness (specificity) of individual proteins, namely any potency to determine the functions, while oligopeptides occurring at low frequencies adversely determine the uniqueness and functions of individual proteins.

[0132] It is suggested that the functional site of protein is composed of oligopeptides occurring at low frequencies. Additionally, longer oligopeptides, more rarely occurring, increase in number. In other words, oligopeptide of length (n+1) as shown in the step (3) according to the method of the first aspect of the present invention is mostly the shortest oligopeptide occurring at a low frequency, and the calculated functional value  $Z_j$  of amino-acid residue  $A_j$  at an appropriate position  $j$  in the oligopeptide is the coefficient of the occurrence (namely, the representative value of the function) of the amino-acid residue  $A_j$  at the position.

[0133] According to the method for predicting the protein functional site in accordance with the second aspect of the present invention, the contribution degree of the amino-acid residue  $A_j$  to the frequency of the occurrence of  $A_j$ -oligopeptide can be evaluated on the basis of the ratio  $Y_{ji}$  of the frequency of the occurrence of the  $A_j$ -oligopeptide to that of the  $X_{ji}$ -oligopeptide as shown in the step (3), and thus, the calculated functional value  $Z(j, n)$  of the amino-acid residue  $A_j$  at an appropriate position of a protein serves as the coefficient of the occurrence of the amino-acid residue  $A_j$  at the position (namely, the representative value of the function).

[0134] Furthermore, the value  $Z(j, n)$  varies, depending on the species of amino-acid residue  $A_j$ . In the step (7) according to the inventive method, the distribution of the  $Z(j, n)$  value of each of twenty amino acids is determined in the entire proteins of organism "a", to determine  $D(j, n)$  value by normalizing the  $Z(j, n)$  value on the basis of the mean and standard deviation of  $Z(j, n)$  value of each amino acid, as determined on the basis of the distribution, which serves as the representative value of the function, after correction of the bias due to each amino-acid residue species.

[0135] Furthermore, longer oligopeptides, more rarely occurring, increase in number. Because the  $Z(j, n)$  and  $D(j, n)$  values generally vary, depending on the length (n), accordingly, the functional value  $W_j$  of the  $Z(j, n)$  and  $D(j, n)$  as determined on a variety of length (n) is defined as the representative value of the function.

[0136] The systems for predicting a protein functional site in accordance with the third and fourth aspects of the present invention are individually systems for automatically carrying out the methods in accordance with the first and second aspects of the present invention; the methods for modifying protein in accordance with the fifth and sixth aspects of the present invention are methods for preparing mutant proteins by substituting the amino-acid residue at the functional site predicted by the method of the first aspect of the present invention with another amino-acid residue. Still further, the method for modifying a protein in accordance with the seventh aspect of the present invention is a method for preparing a mutant protein by substituting the amino-acid residue at the functional site predicted by the method in accordance with the second aspect of the present invention

with another amino-acid residue. In accordance with the eighth aspect of the present invention, a function-modified protein is provided. Furthermore, thermophilic DNA polymerase is provided as an embodiment of such protein.

[0137] The term "DNA polymerase" is the generic name of enzymes catalyzing the synthesis of a DNA chain complementary to a single-stranded DNA. DNA polymerase is an essential enzyme for DNA sequencing and in vitro DNA amplification, and "thermophilic DNA polymerase" is inevitable for PCR (polymerase chain reaction) in terms of the automation of a series of the reaction cycles.

[0138] Such thermophilic DNA polymerase includes known ones, for example Taq, Pfu, KOD, which are separately used, depending on the characteristic performance. Pfu DNA polymerase in particular has been known as an enzyme with an extremely low frequency of erroneous reading during the synthesis of DNA strands (at a high fidelity). However, the Pfu DNA polymerase is inappropriate for the amplification of polymeric DNAs such as genome DNA, because the synthetic DNA yielded by the Pfu DNA polymerase is low and the activity thereof to elongate a synthetic chain is insufficient. Thus, the present application provides a novel Pfu DNA polymerase prepared according to the method of the fifth aspect of the present invention.

#### BRIEF DESCRIPTION OF DRAWINGS

[0139] FIG. 1 depicts graphically the individual frequencies of the occurrences of oligopeptides of lengths 3, 4 and 5, and the distributions of the individual frequencies thereof according to the method of the first aspect of the present invention;

[0140] FIG. 2 depicts an example of an amino acid sequence of a length of 20, and examples of  $A_j$ -oligopeptide of a length of 4, containing the amino-acid residue Met at position 5 in the sequence and examples of  $X_{ji}$ -oligopeptides;

[0141] FIG. 3 depicts graphically the individual frequencies of the occurrences of oligopeptides of lengths of 2, 3, 4 and 5, and the distributions of the individual frequencies thereof according to the method of the second aspect of the present invention;

[0142] FIG. 4 depicts an example of the flow chart for conducting the step (1) of the method of the second aspect of the present invention;

[0143] FIG. 5 depicts an example of the flow chart for conducting the steps (2') to (3) of the method of the second aspect of the present invention;

[0144] FIG. 6 depicts an example of the flow chart for conducting the steps (4) to (5) of the method of the second aspect of the present invention;

[0145] FIG. 7 depicts a distribution of the frequency of  $Z(j, 3)$  of each of three amino acids, see below, according to the method of the second aspect of the present invention, wherein the solid line expresses the distribution of that of isoleucine (Ile); the dotted line expresses the distribution of that of alanine (Ala); and the alternate long and short dash line expresses the distribution of that of methionine (Met);

[0146] FIG. 8 depicts an example of the flow chart for conducting the step (7) of the method of the second aspect of the present invention;

[0147] FIG. 9 depicts an example of the flow chart for conducting the step (8) of the method of the second aspect of the present invention;

[0148] FIG. 10 depicts an example of the flow chart for conducting the step (9) of the method of the second aspect of the present invention;

[0149] FIG. 11 depicts a block diagram illustrating the system of the third aspect of the present invention;

[0150] FIG. 12 depicts a block diagram illustrating the system of the fourth aspect of the present invention;

[0151] FIG. 13 depicts the electrophoresis results of conventional Pfu DNA polymerase and KOD DNA polymerase, indicating the primer elongating activities thereof;

[0152] FIG. 14 depicts a distribution chart of the plotted  $Z_j = -\log Y_j$  value for the whole amino acid sequence of the  $\alpha$ -type DNA polymerase MJ encoded by MJ0885 (gene name from the genome of *Methanococcus jannaschii*), which value is calculated by the first aspect of the present invention;

[0153] FIG. 15 depicts a distribution chart of the plotted  $Z_j = -\log Y_j$  value for partial sequences (motif A and motif C) of the amino acid sequence of which the distribution chart is shown in FIG. 14;

[0154] FIG. 16 depicts a frequency distribution chart of the  $Z_j = -\log Y_j$  value calculated on the basis of the amino acid sequence of the  $\alpha$ -type DNA polymerase MJ encoded by MJ0885;

[0155] FIG. 17 depicts alignment charts of the amino acid sequences of the individual motif Cs from the  $\alpha$ -type DNA polymerases Pfu, KOD and MJ derived from *Pyrococcus furiosus*, *Pyrococcus* sp. KOD, and *Methanococcus jannaschii*, respectively;

[0156] FIG. 18 depicts distribution charts of the plotted  $Z_j = -\log Y_j$  values for the individual motif Cs of the  $\alpha$ -type DNA polymerases Pfu, KOD and MJ;

[0157] FIG. 19 depicts distribution charts of the plotted values of  $W_j = Z(j, 3) - Z(j, 1)$  (in solid line),  $W_j = Z(j, 4) - Z(j, 3)$  (in dotted line) and  $W_j = Z(j, 5) - Z(j, 3)$  (in alternate long and short dash line) of the 100 residues from the N-terminus of the whole amino acid sequence of the  $\alpha$ -type DNA polymerase MJ encoded by MJ0885, and these values are calculated by the method of the second aspect of the present invention;

[0158] FIG. 20 depicts distribution charts of the plotted value  $W_j = Z(j, 5) - Z(j, 3)$  for partial sequences (regions comprising exoi, exoII, motif A, motif B and motif C) of the amino acid sequence of the  $\alpha$ -type DNA polymerase MJ encoded by MJ0885;

[0159] FIG. 21 depicts distribution charts of the plotted values of  $W_j = D(j, 3)$  (in dark color) and  $W_j = D(j, 5)$  (in pale color) for partial sequences (regions comprising exoi, exoII, motif A, motif B and motif C) of the amino acid sequence of the  $\alpha$ -type DNA polymerase MJ encoded by MJ0885;

[0160] FIG. 22 depicts distribution charts in dark color of the positions of amino-acid residues with  $W_j = D(j, 3)$  of 2 or more or of 2 or less in the three-dimensional structure of the amino acid sequence of enolase encoded by MJ0232

(gene name from the genome of *Methanococcus jannaschii*) on a three-dimensional structure model;

[0161] FIG. 23 depicts the results of electrophoresis, indicating the primer elongating activities of the conventional Pfu DNA polymerase (wild type) and the modified Pfu DNA polymerase I of the present invention; and

[0162] FIG. 24 depicts the results of electrophoresis, indicating the primer elongating activities of the conventional Pfu DNA polymerase (wild type) and the modified Pfu DNA polymerases II and III of the present invention.

#### BEST MODE FOR CARRYING OUT THE INVENTION

[0163] The method for predicting a functional site of a protein in accordance with the first aspect of the present invention is a method for predicting the functional site of an appropriate protein of organism "a" with a known genome data or cDNA analysis data, in the entire putative proteins of the organism "a", essentially comprising the following steps (1) to (6).

[0164] Step (1)

[0165] By determining the frequency of the occurrence of each amino acid and the frequencies of the occurrences of individual oligopeptides produced by permutations of twenty amino acids in the amino acid sequences of the entire proteins of the organism "a", the oligopeptide length (n) is determined.

[0166] The length n is determined, then, as the smallest integer satisfying the following criteria.

[0167] "Among oligopeptides of length n, the number of oligopeptides that occur once in the entire proteins is smaller than the number of oligopeptides that occur twice in the entire proteins; among oligopeptides of length (n+1), the number of oligopeptides that occur once in the entire proteins is larger than the number of oligopeptides that occur twice in the entire proteins."

[0168] For example, FIG. 1 depicts distribution charts of the frequencies of the occurrences of oligopeptides of a length of 3, oligopeptides of a length of 4 and oligopeptides of a length of 5, in the entire proteins encoded by the genome of a microorganism *Methanococcus jannaschii* (Bult et al., Science 273, 1058-1073, 1996). In the case of the three types of length of the oligopeptides shown in FIG. 1, the smallest n in the step (1) is 3.

[0169] Step (2)

[0170] Given that the j-th amino-acid residue from the N-terminus of the amino acid sequence (length of L) of the protein as a subject for predicting a functional site is described here as  $A_j$  ( $n+1 \leq j \leq L-n$ ), the frequency of occurrence of a partial sequence of the amino acid sequence of the protein, which sequence corresponds to the following  $A_{ji}$ -oligopeptide of length (n+1), containing the j-th amino-acid residue  $A_j$ ;

[0171]  $A_{ji}$ -oligopeptide:  $a_j 1 a_j 2 \dots A_{ji} \dots a_j n a_j (n+1)$  ( $1 \leq i \leq n+1$ ;  $A_j = A_{ji}$ ; and  $A_j$  is the i-th residue of the oligopeptide; and moreover,  $a_j 1 = A_{j-i+1}$ ,  $a_j (n+1) = A_{j-i+(n+1)}$ ),

[0172] and the frequency of the occurrence of the following X<sub>ji</sub>-oligopeptide of length (n+1);

[0173] X<sub>ji</sub>-oligopeptide: aj<sub>1</sub>aj<sub>2</sub> . . . X<sub>ji</sub> . . . aj<sub>n</sub>aj<sub>(n+1)</sub> (the i-th residue X<sub>ji</sub> is any amino acid; and moreover, aj<sub>1</sub>=A<sub>j-i+1</sub>, . . . , aj<sub>(n+1)</sub>=A<sub>j-i+(n+1)</sub>) should be determined in the entire proteins of the organism "a".

[0174] Such A<sub>ji</sub>-oligopeptide and X<sub>ji</sub>-oligopeptide can be illustrated for example in FIG. 2. The upper row (1) in FIG. 2 expresses in single letter code the partial sequence from the N-terminus to the 20-th amino-acid residue of the putative amino acid sequence on the basis of the gene MJ0885, which is believed to encode the α-type DNA polymerase of *Methanococcus jannaschii* (Bult et al., Science 273, 1058-1073, 1996); the middle row (2) expresses examples of A<sub>ji</sub>-oligopeptide of a length of 4, containing the 5-th amino-acid residue Met(M) in the amino acid sequence; and the rows (3) to (6) further below express examples of X<sub>ji</sub>-oligopeptide containing the 5-th amino-acid residue M.

[0175] Step (3)

[0176] Calculating ratio Y<sub>ji</sub> of the frequency of the occurrence of the A<sub>ji</sub>-oligopeptide to that of the X<sub>ji</sub>-oligopeptide.

[0177] Step (4)

[0178] The mean Y<sub>j</sub> of the Y<sub>ji</sub> is determined as follows.

$$Y_j = \sum_{i=1}^{n+1} Y_{ji} / (n+1),$$

[0179] Step (5)

[0180] Monotonously decreasing functional value or monotonously increasing functional value Z<sub>j</sub> of Y<sub>j</sub> is determined as follows;

$$[0181] Z_j = f(Y_j)$$

[0182] The Z<sub>j</sub> value is defined as the representative value of the function of the j-th amino-acid residue of the amino acid sequence (length of L).

[0183] Step (6)

[0184] By subsequently repeating the steps (2) to (5) sequentially and determining the Z<sub>j</sub> value of each of all the amino-acid residues at position n+1 ≤ j ≤ L-n, the degree of the involvement of each amino-acid residue in the function of the protein is predicted by using the dimension of the Z<sub>j</sub> value as an indicator. More specifically, because the manner of occurring of each amino-acid residue in the context is expressed as the functional value Z<sub>j</sub> of Y<sub>j</sub>, a larger Z<sub>j</sub> value indicates a lower frequency of occurrence of the amino-acid residue if Z<sub>j</sub> is a monotonously decreasing functional value, which suggests that the amino-acid residue has higher responsibility over the fulfillment of the function. If Z<sub>j</sub> is a monotonously increasing function, additionally, it is suggested that an amino-acid residue with a smaller Z<sub>j</sub> value has greater responsibility over the function.

[0185] By expressing the Z<sub>j</sub> value of each amino-acid residue for example in a distribution chart wherein the Z<sub>j</sub> value is plotted on the vertical axis while the amino acid sequence is shown on the horizontal axis, furthermore, the

functional site can be confirmed at a glance, which is preferable as an embodiment for carrying out the present invention.

[0186] The method for predicting a protein functional site in accordance with the second aspect of the present invention is a method for predicting a functional site of an appropriate protein in the entire putative proteins of the organism "a" with a known genome data or cDNA analysis data, essentially comprising the following steps (1) to (9).

[0187] Step (1)

[0188] The frequency of the occurrence of each amino acid and the frequencies of the occurrences of individual oligopeptides produced by permutations of twenty amino acids, in the amino acid sequences of the entire proteins of the organism "a", are determined.

[0189] For example, FIG. 3 shows a distribution chart of the frequencies of the occurrences of oligopeptides of a length of 3, oligopeptides of a length of 4 and oligopeptides of a length of 5, which are determined in the entire proteins encoded by the genome of a microorganism *Methanococcus jannaschii* (Bult et al., Science 273, 1058-1073, 1996) on the basis of the genome data of the microorganism.

[0190] FIG. 4 depicts an example of the flow chart for carrying out the step (1).

[0191] Step (2)

[0192] As to an appropriate protein of organism "a",

[0193] Step (2')

[0194] given that the j -th amino-acid residue from the N-terminus of the amino acid sequence (length of L) of the protein is described here as A<sub>j</sub>, the frequency of the occurrence of a partial sequence of the amino acid sequence of the protein, which sequence corresponds to the following A<sub>ji</sub>-oligopeptide of an appropriate length n (1 ≤ n ≤ M, provided that "M" is the smallest length of oligopeptides satisfying the following criterion; all the oligopeptides of length M are at frequency 1 of the occurrence), containing the j-th amino-acid residue A<sub>j</sub> (n ≤ j ≤ L-n+1);

[0195] A<sub>ji</sub>-oligopeptide: aj<sub>1</sub>aj<sub>2</sub> . . . A<sub>ji</sub> . . . aj<sub>n</sub> (1 ≤ i ≤ n; A<sub>j</sub>=A<sub>ji</sub> and A<sub>ji</sub> is the i-th residue of the oligopeptide; and moreover, aj<sub>1</sub>=A<sub>j-i+1</sub>, aj<sub>(n+1)</sub>=A<sub>j-i+(n+1)</sub>),

[0196] and the frequency of the occurrence of the following X<sub>ji</sub>-oligopeptide of the length n corresponding to the length of the A<sub>ji</sub>-oligopeptide;

[0197] X<sub>ji</sub>-oligopeptide: aj<sub>1</sub>aj<sub>2</sub> . . . X<sub>ji</sub> . . . aj<sub>n</sub> (the i-th residue X<sub>ji</sub> is any amino acid; and moreover, aj<sub>1</sub>=A<sub>j-i+1</sub>, . . . , aj<sub>(n+1)</sub>=A<sub>j-i+(n+1)</sub>), are determined in the entire proteins of the organism "a".

[0198] In the same manner as by the method of the first aspect of the present invention, such A<sub>ji</sub>-oligopeptide and X<sub>ji</sub>-oligopeptide are for example illustrated as in FIG. 2.

[0199] Step (3)

[0200] The ratio Y<sub>ji</sub> of the frequency of the occurrence of the A<sub>ji</sub>-oligopeptide to that of the X<sub>ji</sub>-oligopeptide is determined.

[0201] FIG. 5 depicts an example of the flow chart for carrying out the aforementioned steps (2') to (3).

[0202] Step (4)

[0203] The mean  $Y(j, n)$  of the  $Y_{ji}$  is determined as described below.

$$Y(j, n) = \sum_{i=1}^n Y_{ji}/n.$$

[0204] Step (5)

[0205] The logarithmic value  $Z(j, n)$  of  $Y(j, n)$  is determined as follows.

$$[0206] \quad Z(j, n) = -\log(Y(j, n))$$

[0207] FIG. 6 depicts an example of the flow chart for carrying out the aforementioned steps (4) to (5).

[0208] Step (6)

[0209] By subsequently repeating the steps (2') to (5) sequentially, the  $Z(j, n)$  value of each of all the amino-acid residues at position  $n \leq j \leq L-n+1$  in the amino acid sequence (length of  $L$ ) is determined.

[0210] Step (7)

[0211] By sequentially repeating the steps (2) to (6) over the entire proteins of the organism "a", thereby determining the distribution of the  $Z(j, n)$  value of each amino-acid residue in the entire proteins, and the  $Z(j, n)$  values are classified into twenty according to the twenty amino acids, and then determining mean  $Av(Aa)$  of the  $Z(j, n)$  values for each amino acid  $Aa$  and the standard deviation  $Sd(Aa)$  of the distribution thereof, on the basis of the distribution, to determine function  $g$  to the  $j$ -th amino-acid residue  $A_j$  of a protein for normalizing the difference in distribution due to (or among?) the species of amino-acid residues is determined;

$$[0212] \quad g = g(Z(j, n), A_j) = [Z(j, n) - Ad(Aa)] / Sd(Aa)$$

(provided that  $A_j = Aa$ ; and  $1 \leq n \leq M$ ).

[0213] For example, FIG. 7 depicts a distribution of the frequency of  $Z(j, n)$  for three species of amino acids, namely isoleucine (Ile), alanine (Ala) and methionine (Met), in the entire proteins encoded by the genome of *Methanococcus jannaschii* (Bult et al., Science 273, 1058-1073, 1996). Based on the distribution, the mean and standard deviation of the  $Z(j, n)$  values for an amino acid isoleucine (Ile), namely  $Ad(Ile)$  and  $Sd(Ile)$ , respectively, are determined as  $Ad(Ile) = 3.16$  and  $Sd(Ile) = 0.17$ , and the function  $g$  for  $A_j = Ile$  is determined as follows.

$$[0214] \quad g = g(Z(j, n), A_j) = (Z(j, n) - 3.16) / 0.17$$

[0215] FIG. 8 depicts an example of the flow chart for carrying out the step (7).

[0216] Step (8)

[0217] The value  $D(j, n)$  of the function  $g$  of each of all the amino-acid residues  $A_j$  at position  $n \leq j \leq L-n+1$  in the amino acid sequence (length of  $L$ ) as recovered in the step (7) is determined;

$$[0218] \quad D(j, n) = g(Z(j, n), A_j).$$

[0219] FIG. 9 depicts an example of the flow chart for carrying out the step (8).

[0220] Step (9)

[0221] The functional value  $W_j$  of the  $Z(j, n)$  value and the  $D(j, n)$  value is determined as follows.

$$[0222] \quad W_j = h(Z(j, 1), Z(j, 2), \dots, Z(j, M), D(j, 1), D(j, 2), \dots, D(j, M))$$

[0223] By defining the value of the  $W_j$  as the representative value of the function of the  $j$ -th amino-acid residue in the amino acid sequence (length of  $L$ ), the degree of the responsibility of each amino-acid residue over the function of the protein is estimated by using the dimension of the  $W_j$  value as an indicator.

[0224] FIG. 10 depicts an example of the flow chart for carrying out the step (9).

[0225] By expressing the  $W_j$  value of each amino-acid residue for example in a distribution chart wherein the  $W_j$  value is plotted on the vertical axis while the amino acid sequence is shown on the horizontal axis, furthermore, the functional site can be confirmed at a glance, which is preferable as an embodiment for carrying out the present invention.

[0226] If the three-dimensional structure of the protein as a subject for predicting the functional site is known or if a three-dimensional structure model thereof can be prepared by known methods (for example, homology modeling method, Peitsch, Proceedings of the Fifth International Conference on Intelligent Systems for Molecular Biology, 1997, 5, 234-236), the distribution is expressed on the three-dimensional structure, whereby a spatial arrangement of an amino-acid residue as a candidate of a novel functional site can be confirmed, which is preferable as an embodiment for carrying out the invention.

[0227] The system for predicting a functional site of a protein in accordance with the third aspect of the present invention is a system for automatically carrying out the method for predicting a functional site in accordance with the first aspect of the present invention, at least comprising the following units (a) to (g) for conducting the steps (1) to (6) according to the method of the first aspect of the present invention, as shown for example in the composition example in FIG. 11.

[0228] Outer Memory Unit (a):

[0229] Unit memorizing the amino acid sequence data of a protein or an existing protein data base for use in the step (1).

[0230] Calculation/Memory Unit (b):

[0231] Unit, composed of CPU calculating the frequencies of the occurrences of individual oligopeptides as determined in the step (1), and a memory unit having the memory of the calculation results.

[0232] Calculation/Memory Unit (c):

[0233] Unit, composed of CPU calculating the smallest length ( $n$ ) of oligopeptides as determined in the step (1) and a memory unit having the memory of the length  $n$ .

[0234] Calculation/Memory Unit (d):

[0235] Unit, composed of CPU calculating the the frequencies of occurrence of each amino acid and the frequencies of occurrence of A<sub>ji</sub>-oligopeptide and X<sub>ji</sub>-oligopeptide in the entire proteins as determined in the step (2) and a memory unit having the memory of the calculation results.

[0236] Calculation/Memory Unit (e):

[0237] Unit, composed of CPU calculating the Y<sub>ji</sub> value as determined in the step (3) and a memory unit having the memory of the Y<sub>ji</sub> value.

[0238] Calculation/Memory Unit (f):

[0239] Unit, composed of CPU calculating the Y<sub>j</sub> value as determined in the step (4) and a memory unit having the memory of the Y<sub>j</sub> value.

[0240] Calculation/Memory Unit (g):

[0241] Unit, composed of CPU calculating the Z<sub>j</sub> value as determined in the step (5) and a memory unit having the memory of Z<sub>j</sub>.

[0242] Additionally, the system for predicting a functional site is provided with the following display unit (h) in a preferable embodiment.

[0243] Display Unit (h):

[0244] Unit displaying the Z<sub>j</sub> value. of each amino-acid residue recovered in the calculation/memory unit (g) in a distribution chart.

[0245] The system of the present invention may be equipped with keyboard (i) and control unit (j) and the like as illustrated in **FIG. 11**, in addition to these units (a) to (h).

[0246] According to the fourth aspect of the present invention, the system for predicting a protein functional site is a system for automatically conducting the method of the second aspect of the present invention, at least comprising the following units (a) to (i) for carrying out the steps (1) to (9) according to the method of the second aspect of the present invention, as shown in the composition example in **FIG. 12**.

[0247] Outer Memory Unit (a):

[0248] Unit memorizing the amino acid sequence data and an existing protein data base for use in the step (1).

[0249] Calculation/Memory Unit (b):

[0250] Unit, composed of CPU calculating the frequencies of the occurrences of individual oligopeptides as determined in the step (1) and a memory unit having the memory of the calculation results.

[0251] Calculation/Memory Unit (c):

[0252] Unit, composed of CPU calculating the frequencies of occurrence of each amino acid and the individual frequencies of the occurrences of A<sub>ji</sub>-oligopeptide and X<sub>ji</sub>-oligopeptide in the entire proteins as determined in the step (2) and a memory unit having the memory of the calculation results.

[0253] Calculation/Memory Unit (d):

[0254] Unit, composed of CPU calculating Y<sub>ji</sub> as determined in the step (3) and a memory unit having the memory of the Y<sub>ji</sub> value.

[0255] Calculation/Memory Unit (e):

[0256] Unit, composed of CPU calculating the Y<sub>(j, n)</sub> value as determined in the step (4) and a memory unit having the memory of the Y<sub>(j, n)</sub> value.

[0257] Calculation/Memory Unit (f):

[0258] Unit, composed of CPU calculating the Z<sub>(j, n)</sub> value as determined in the steps (5) and (6) and a memory unit having the memory of the Z<sub>(j, n)</sub> value.

[0259] Calculation/Memory Unit (g):

[0260] Unit, composed of CPU calculating the g value as determined in the step (7) and a memory unit having the memory of the g value.

[0261] Calculation/Memory Unit (h):

[0262] Unit, composed of CPU calculating the D<sub>(j, n)</sub> value as determined in the step (8) and a memory unit having the memory of the D<sub>(j, n)</sub> value.

[0263] Calculation/Memory Unit (i):

[0264] Unit, composed of CPU calculating the W<sub>j</sub> value as determined in the step (9) and a memory unit having the memory of the W<sub>j</sub> value.

[0265] Additionally, the system for predicting a functional site in accordance with the fourth aspect of the present invention may be equipped with an appropriate combination of the following units (j) to (l).

[0266] Display Unit (j):

[0267] Unit displaying the W<sub>j</sub> value of each amino-acid residue as recovered with the unit (i) in a distribution chart.

[0268] Calculation/Memory Unit (k):

[0269] Unit memorizing an existing database of protein three-dimensional structures or unit preparing a three-dimensional structure model based on an amino acid sequence according to a known method and memorizing the three-dimensional structure.

[0270] Display Unit (l):

[0271] Unit displaying the W<sub>j</sub> value of each amino-acid residue in a distribution chart on the three-dimensional structure stored in the database or three-dimensional structure model recorded on the unit (k).

[0272] The system of the present invention may satisfactorily be equipped with keyboard (m) and control unit (n) and the like as illustrated in **FIG. 12**, in addition to these units (a) to (l).

[0273] The method for modifying the function of a protein in accordance with the fifth aspect of the present invention will be described below. The method is a method for modifying the function of protein "A", which function has been known, derived from the entire putative proteins of organism "a" with a known genome data or cDNA analysis data, essentially comprising the following steps (1) to (10).

[0274] Step (1):

[0275] Extracting proteins closely related to the protein "A" from an existing protein data base and subjecting the proteins to alignment.

[0276] Steps (2) to (7):

[0277] Subjecting the amino acid sequences of the entire proteins of the organism "a" to the steps (1) to (6) according to the method of the first aspect of the present invention.

[0278] Step (8):

[0279] Selecting at least one amino-acid residue to be subjected to mutation from the amino acid sequence (length of L) of the protein "A" on the basis of the alignment data recovered in the step (1), sequentially repeating the steps (3) to (6) for a variant amino-acid residue of various mutated amino acid sequences where the selected amino-acid residue has been mutated into another amino-acid residue, to determine the Z<sub>j</sub> value of the variant amino-acid residue.

[0280] Step (9):

[0281] Selecting a mutated amino acid sequence wherein the Z<sub>j</sub> value of the variant amino-acid residue as determined in the step (8) is larger or smaller than the Z<sub>j</sub> value of the intact amino-acid residue as determined in the step (7).

[0282] Step (10):

[0283] Preparing a modified gene of the protein "A", which gene encodes the mutant amino acid sequence selected in the step (9), and expressing the modified gene in an appropriate host-vector system to prepare modified protein "A".

[0284] In accordance with the sixth aspect of the present invention, the method for modifying a protein function is a method for modifying the function of protein "B" derived from organism "b" with an unknown genome data or cDNA analysis data, essentially comprising the following steps (1) to (10).

[0285] Step (1):

[0286] Extracting protein "A" most closely related to protein "B" from the entire putative proteins of organism "a" with a known genome data or cDNA analysis data and subjecting the protein "A" to alignment, or extracting proteins closely related to protein "B" from an existing protein data base to subject the proteins to alignment.

[0287] Steps (2) to (8):

[0288] Conducting the steps (2) to (8) of the method of the third aspect of the present invention over the amino acid sequences of the entire proteins of the organism "a".

[0289] Step (9):

[0290] Selecting a position that should be mutated and an amino acid residue for which should be substituted wherein the Z<sub>j</sub> value of the substituted amino-acid residue as determined in the step (8) is larger or smaller than the Z<sub>j</sub> value of the intact amino-acid residue as determined in the step (7).

[0291] Step (10):

[0292] Preparing a modified gene of the protein B according to a known method, which gene encodes the amino acid

sequence mutated at the position to another amino acid residue as selected in the step (9), and expressing the modified gene in an appropriate host-vector system to prepare modified protein B.

[0293] As has been described above, the methods for modifying the protein function in the fifth and sixth aspects of the present invention, comprising the method for predicting a functional site of the first aspect of the present invention, are characterized in that an unknown functional site of protein is newly found and is subjected to mutation.

[0294] Furthermore, the method for modifying a protein function according to the seventh aspect of the present invention also utilizes the method for predicting the function in accordance with the second aspect of the present invention, whereby the method can be carried out in the same manner as in the case of the fifth aspect of the present invention.

[0295] The eighth aspect of the present invention is a protein of which function is artificially modified by the method of the fifth to seventh aspect of the present invention. An embodiment of such protein is thermophilic DNA polymerase, which is more specifically an enzyme prepared by modifying, in accordance with the sixth aspect of the present invention, a thermophilic Pfu DNA polymerase derived from *Pyrococcus furiosus* in a genetic engineering manner by the known method for preparing mutant gene (Strategies, Vol.9, p.3-4, 1996) (the thermophilic DNA polymerase of the present invention is sometimes referred to as "modified Pfu DNA polymerase"). The enzyme can be prepared as follows. More specifically, because the nucleotide sequence of the gene of Pfu DNA polymerase is known (Nucleic Acids Research, Vol.21, p.259-265, 1993), the gene of Pfu DNA polymerase is prepared by PCR comprising synthetically preparing an oligopeptide complementary to both the ends by using the genome DNA of the archaeobacterium as template and using the oligopeptide as primer. The DNA fragment of the gene is cloned into a vector, and the gene is subsequently subjected to mutation by the method described in the reference mentioned above. In accordance with the present invention, in particular, the mutation of the gene was executed by nucleotide substitution, so that a part of the amino acid sequence of Pfu DNA polymerase might be substituted with the amino-acid residues from KOD DNA polymerase. In terms of amino acid sequence, Pfu DNA polymerase has about 80% homology with KOD DNA polymerase, and therefore, similar synthetic termination occurs during PCR (FIG. 13), but the elongation rate with KOD DNA polymerase is about 6-fold the rate with Pfu DNA polymerase. By substituting some amino-acid residues of Pfu DNA polymerase with some amino-acid residues of KOD DNA polymerase, the synthetic termination of the chain elongation might be improved or the elongation rate turns rapid, which possibly enables the recovery of an enzyme capable of elongating a DNA chain under way of synthesis more longer. By expressing in *Escherichia coli* the mutant gene that was mutated in such a manner and recovering and purifying the expression product, the modified Pfu DNA polymerase of the present invention was recovered.

[0296] The thermophilic DNA polymerase (modified Pfu DNA polymerase I) of the present invention is more specifically an enzyme of the amino acid sequence of SQ ID No.1. The amino acid sequence is a novel sequence prepared

by identifying potentially function-modifiable amino-acid residues of the amino acid sequence of the conventionally known Pfu DNA polymerase, according to the inventive method for predicting a functional site, and substituting the amino-acid residues as shown in Table 1. By using the novel enzyme then for DNA synthesis by PCR, for example, the synthetic termination occurring when using the conventional DNA polymerases is almost totally overcome, as shown in the following examples. It is needless to say that template DNA chains to be highly efficiently amplified with the conventional polymerase can be amplified at high efficiency in the same manner.

[0297] Furthermore, the thermophilic DNA polymerases (modified Pfu DNA polymerases II and III) of the present invention are more specifically enzymes of amino acid sequences of SQ ID Nos 6 and 7, which are novel sequences prepared by identifying potentially function-modifiable amino-acid residues of the amino acid sequence of the Pfu DNA polymerase, according to the inventive method for predicting a functional site, and substituting the amino-acid residues as shown in Table 1. By using the novel enzymes then for DNA synthesis by PCR, for example, synthetic polymeric products can be recovered at large scales, as shown in the following examples.

Table 1

Modified DNA polymerases	Positions	Wild-type amino acid	Modified amino acid
I	2	Ile	Val
	533	Phe	Tyr
	538	Leu	Ile
	540	Ile	Ser
	545	Leu	Phe
	546	Tyr	Phe
II	2	Ile	Val
	710	Pro	Arg
	712	Ser	Arg
	713	Asn	Asp
	717	Leu	Pro
	717	Ile	Val
III	2	Ile	Val
	717	Leu	Pro

[0298] The DNA sequences encoding these modified Pfu DNA polymerases include for example the mutant genes of the Pfu DNA polymerase gene, as recovered during the process of enzyme preparation. As to these mutant genes, the DNA sequences encoding the amino acid sequences of SQ ID Nos. 1, 6 and 7 for example have been cloned in recombinant plasmids p320, pDP5b17 and pDP5C4, respectively, and these recombinant plasmids have been integrated in *Escherichia coli* HMS174 (DE3) and deposited at the Life Engineering and Industrial Technology Research Institute, the Agency of Industrial Science and Technology, Japan (Accession Nos. FERM P-16052, FERM BP-6189 and FERM BP-6190, respectively).

[0299] Additionally, the DNA sequences of the present invention may appropriately be designed as DNA sequences with conjugated nucleotide codons corresponding to the individual amino-acid residues of SQ ID No. 1, 6 or 7.

[0300] The thermophilic DNA polymerases of the present invention may be expressed in microorganisms such as *Escherichia coli*, which may thereafter be recovered. By inserting and integrating the DNA sequence into an expres-

sion vector with an origin of replication in a microorganism, a promoter, a ribosome-binding site, a cDNA cloning site, and a terminator and the like to prepare an expression vector, transforming a host cell with the expression vector and thereafter culturing the resulting transformant, an enzyme encoded by the DNA sequence can be generated in the microorganism at a large scale.

## EXAMPLES

[0301] The present invention will now be described more specifically in more detail with reference to examples, but the invention is not limited to the following examples.

### Example 1

[0302] According to the method of the first aspect of the present invention and based on the genome data of *Methanococcus jannaschii* (Bult et al., Science 273, 1058-1073, 1996),  $Z_j = -\log Y_j$  ( $f = -\log$ ) was calculated, concerning each amino-acid residue in the amino acid sequence (from the N-terminus to the C terminus) of a DNA polymerase speculated from the microbial gene MJ0885 believed to encode the  $\alpha$ -type DNA polymerase. The results are plotted in a distribution chart in FIG. 14.

[0303] Among the motifs known as the functional sites of the  $\alpha$ -type DNA polymerase, furthermore, motif A and motif C were extracted, and the  $Z_j$  values of the individual amino-acid residues were plotted in FIG. 15. FIG. 15 and FIG. 16 below suggest that the  $Z_j = -\log Y_j$  values of amino-acid residues responsible for the function are larger than those of the remaining amino-acid residues.

[0304] FIG. 16 depicts a distribution chart of the frequency of the value  $Z_j = -\log Y_j$  for the amino acid sequence of the  $\alpha$ -type DNA polymerase encoded by MJ0885. It is confirmed in the figure that amino-acid residues with value  $Z_j = -\log Y_j$  of 4.8 or more are highly possibly amino-acid residues responsible for the protein function.

### Example 2

[0305] Following the chart on FIG. 15 in Example 1, the characteristic properties of  $\alpha$ -type DNA polymerase Pfu (DDBJ Accession No. D12983) derived from *Pyrococcus furiosus* were modified, on the basis of the amino acid sequence of an  $\alpha$ -type DNA polymerase KOD (DDBJ Accession No. D29671) derived from *Pyrococcus* sp., and the genome data of *Methanococcus jannaschii* (Bult et al., Science 273, 1058-1073, 1996), and the amino acid sequence of the  $\alpha$ -type DNA polymerase (named here as MJ) encoded by MJ0885.

[0306] FIG. 17 depicts alignment charts of the amino acid sequences of the motif Cs from Pfu, KOD and MJ, with no difference between the region 531 to 544 from Pfu and the region 551 to 564 from MJ (the numbers 550 and 570 indicate the N-terminus and C-terminus amino acid residues of the motif C shown here from MJ).

[0307] FIG. 18 depicts the results of the prediction of functional sites in the amino acid sequences of the motif Cs of Pfu, KOD and MJ according to the inventive method, on the basis of the genome data of *Methanococcus jannaschii*. The results indicate that mutations Ile540Ser, Leu545Phe, Tyr546Phe, and Ile548Thr increase the  $Z_j = -\log Y_j$  values of the amino-acid residues. Furthermore, the values  $Z_j = -\log Y_j$

of Asp541 and Ala547 are increased. By subjecting the  $\alpha$ -type DNA polymerase MJ of *Methanococcus jannaschii* to such mutation, the resulting sequence turns more unique (specific) which possibly brings about the improvement of some function.

#### Example 3

[0308] Based on the genome data of *Methanococcus jannaschii* (Bult et al., Science 273, 1058-1073, 1996), the values  $Z(j, 1) = -\log Y(j, 1)$ ,  $Z(j, 3) = -\log Y(j, 3)$ ,  $Z(j, 4) = -\log Y(j, 4)$ , and  $Z(j, 5) = -\log Y(j, 5)$  were calculated, concerning the individual amino-acid residues of the amino acid sequence (from the N to C termini) of a DNA polymerase speculated on the basis of the microbial gene MJO885 believed to encode the  $\alpha$ -type DNA polymerase, according to the method of the second aspect of the present invention, so that  $W_j = Z(j, 3) - Z(j, 1)$  ( $h = Z(j, 3) - Z(j, 1)$ ) was calculated. Similarly,  $W_j = Z(j, 4) - Z(j, 3)$  ( $h = Z(j, 4) - Z(j, 3)$ ) and  $W_j = Z(j, 5) - Z(j, 3)$  ( $h = Z(j, 5) - Z(j, 3)$ ) were also calculated.

[0309] FIG. 19 depicts the results of the 100 residues from the N-terminus in a plotted distribution chart. Given  $h = Z(j, 5) - Z(j, 3)$ , regions with significantly different distributions from those of the remaining two cases are present in the region from the 35-th to 60-th residues and the like. The distributions indicate that smaller  $W_j = Z(j, 5) - Z(j, 3)$  more specifically characterizes the amino acid sequence.

[0310] Among the motifs known as the functional sites of the  $\alpha$ -type DNA polymerases, furthermore, regions containing exoI, exoII, motif A, motif B and motif C were extracted, and subsequently, the  $W_j$  values of the individual amino-acid residues are plotted in FIG. 20. As shown in FIG. 20, the regions with the characteristic reduction of  $W_j$  are consistent with the functional sites.

#### Example 4

[0311] FIG. 21 depicts the values  $W_j = D(j, 3)$  and  $W_j = D(j, 5)$  of individual amino-acid residues in the regions containing the exoI, exoII, motif A, motif B and motif C extracted among the motifs known as the functional sites in the  $\alpha$ -type DNA polymerases ( $h = D(j, 3)$  and  $h = D(j, 5)$ ). Amino-acid residues with  $W_j = D(j, n)$  of 2 or more or of 2 or less are present outside the motifs, and these amino-acid residues are candidates of new functional sites.

#### Example 5

[0312] FIG. 22 depicts in a dark color the positions of amino-acid residues having  $W_j = D(j, 3)$  of 2 or more or of 2 or less in the amino acid sequence of MJ0232, which is speculated as enolase of *Methanococcus jannaschii*, on a three-dimensional structure model prepared on the basis of the enolase of budding yeast. It is indicated that residues positioned apart on the amino acid sequence are closely positioned on the three-dimensional structure.

#### Example 6

[0313] Modified Pfu DNA polymerase I was prepared, by applying the mutation of putative amino-acid residues for improving the function of the DNA polymerase MJ in Example 2 to the Pfu DNA polymerase.

[0314] (1) Preparation of Modified Pfu DNA Polymerase Gene Cloning of Pfu DNA Polymerase Gene:

[0315] Following the nucleotide sequence of Pfu DNA polymerase gene (Nucleic Acids Research, Vol.21, p.259-265, 1993), a PCR primer was prepared, for amplifying the objective gene by PCR by using the genome DNA of *P. furiosus* as template, which was then cloned in an expression vector for *Escherichia coli*. The detail is described below.

[0316] *P. furiosus* DSM3638 was cultured according to the method described in the reference described above. First, the culture medium described in the reference was prepared, followed by sterilization at a high temperature under pressure, and subsequently, nitrogen gas was purged into the resulting culture medium. The bacterium was inoculated into the culture medium, for stationary culturing at 95° C. for 15 hours. From 200 ml of the culture broth were recovered the bacteria of about 0.5 mg by centrifugation. The collected bacteria were suspended in buffer A (10 mM Tris/HCl (pH 8.0), 1 mM EDTA, 100 mM NaCl), followed by addition of 1 ml of 10% SDS and subsequent agitation, and to the resulting suspension was added 0.5 mg of proteinase K for reaction at 55° C. for 60 minutes. The reaction solution was extracted sequentially in phenol, phenol/chloroform, and chloroform, and to the extract was added ethanol to make the DNA insoluble, which was then recovered. The resulting DNA was dissolved in 1 ml of TE buffer (10 mM Tris/HCl (pH 8.0), 1 mM EDTA), followed by addition of 0.5 mg RNase A for reaction at 37° C. for 60 minutes and re-extraction sequentially in phenol, phenol/chloroform, and chloroform, and subsequent ethanol precipitation, to recover the DNA, which was then dissolved in the TE buffer, to recover the DNA at about 0.3 mg.

[0317] For PCR amplification of the objective DNA polymerase gene, then, two primer DNAs of SQ ID Nos.2 and 3 were synthesized on the basis of the known sequence data. More specifically, it was designed that the initiation codon ATG of the objective gene and a restriction nuclease NcoI sequence (5'-CCATGG-3') might be, introduced in the forward primer sequence, while the reverse primer might be conjugated at an appropriate position downstream the termination codon. PCR was conducted in a reaction system of 50  $\mu$ l, by using 2  $\mu$ g of *P. furiosus* DNA and 10 pmol each of the primers under conditions for LA Taq (manufactured by TaKaRa Brewery) and attached buffers. The cycle conditions were as follows; 93° C./3 minutes prior to the addition of the enzyme, and 30 cycles of each cycle composed of 94° C./0.5 minute, 55° C./0.5 minute and 72° C./1.0 minute. The amplified DNA fragment was purified, followed by treatment with NcoI, and the resulting DNA fragment was similarly cleaved with NcoI and subsequently blunt ended, and the resulting fragment was then integrated downstream the T7 promoter of an NcoI-treated expression vector pET-15b. The expression vector was defined as pDPWT100, to confirm the nucleotide sequence of the inserted gene.

[0318] Modification of Pfu DNA Polymerase Gene:

[0319] According to the known method (Strategies, Vol.9, p.3-4, 1996) and for the expression vector pDPWT100 with

the cloned Pfu DNA polymerase gene integrated therein, a modified Pfu DNA polymerase gene was prepared on the expression vector pDPWT100 by using oligopeptides containing desired mutations (SQ ID Nos.4 and 5) and the mutation induction kit manufactured by Promega Corporation, whereby an expression vector pDP320 was constructed. By determining the nucleotide sequence of the modified gene, furthermore, the amino acid sequence of the modified Pfu DNA polymerase (SQ ID No.1) was verified.

**[0320]** (2) Expression and Purification of the Modified Pfu DNA Polymerase in *Escherichia coli*

**[0321]** The gene of the modified Pfu DNA polymerase I was expressed in *Escherichia coli* as follows, which was then purified.

**[0322]** The expression vector pDP320 with the modified Pfu DNA polymerase gene was inserted in a strain *Escherichia coli* HMS174 (DE3) and cultured in an LB culture medium supplemented with IPTG to a final concentration of 0.1 mM for 14 hours, to induce the expression of the enzyme in the bacteria of the strain *Escherichia coli*. After harvesting the bacteria by centrifugation, a modified Pfu DNA polymerase was extracted under ultrasonic treatment in a buffer containing 150 mM Tris/HCl (pH 7.5), 2 mM EDTA, 0.24 mM APMSF and 0.2% Tween 20. The crude extract solution was thermally treated at 80° C. for 15 minutes, to inactivate the DNA polymerase derived from *Escherichia coli* and partially purify the DNA polymerase of the present invention. The partially purified fraction was dialyzed against a buffer composed of 50 mM Tris/HCl (pH 7.5), 1 mM EDTA, 0.2% Tween 20, 7 mM 2-mercaptoethanol and 10% glycerol. At the stage was detected a DNA polymerizing activity specific to the modified Pfu DNA polymerase I.

#### Example 7

**[0323]** By using the modified Pfu DNA polymerase I partially purified in Example 6, the primer elongation reaction of a DNA chain complementary to the template DNA was tested.

**[0324]** One  $\mu\text{g}$  of the partially purified enzyme fraction described above was placed in 20  $\mu\text{l}$  of a reaction solution containing 20 mM Tris/HCl (pH 8.0), 2 mM  $\text{MgCl}_2$ , 50  $\mu\text{g/ml}$  BSA, 0.1% Triton X-100, 1 mM each of cold dNTPs (0.1 mM for dCTP), 0.63  $\mu\text{g}$  of pBLUESCRIPT plasmid prepared by annealing together 10  $\mu\text{Ci}$  of  $[\alpha\text{-}^{32}\text{P}]\text{dCTP}$  and a primer of M13(-21), for reaction at 75° C. for one minute and 3 minutes. The elongated DNA chain was separated by electrophoresis on a polyacrylamide gel containing 8M urea, and the resulting pattern was analyzed with an image analyzer. As a control, additionally, the conventional wild-type Pfu DNA polymerase was used for the same DNA synthesis.

**[0325]** The results are shown in FIG. 23. When the conventional wild-type Pfu DNA polymerase was used, at least 10 bands indicating the presence of incomplete DNA chains due to synthetic termination were observed. However, these bands disappeared during the DNA synthesis with the modified Pfu DNA polymerase I of the present invention. Alternatively, no difference in the accumulation of highly elongated DNA chains around 1000 bases was observed.

#### Example 8

**[0326]** Pfu DNA polymerases II and III were prepared.

**[0327]** (1) Preparation of Modified Pfu DNA Polymerase Gene

**[0328]** In the same manner as in Example 6(1), the Pfu DNA polymerase gene was cloned, to prepare modified genes II and III as follows.

**[0329]** Preparation of Modified Pfu DNA Polymerase II:

**[0330]** According to the known method (Strategies, Vol.9, p.3-4, 1996) and for the expression vector pDPWT100 with the cloned Pfu DNA polymerase gene integrated therein, the gene of modified Pfu DNA polymerase II was prepared on the expression vector pDPWT100 by using oligopeptides containing desired mutations (SQ ID Nos.8 and 9) and the mutation induction kit manufactured by Promega Corporation, whereby an expression vector pDP5b17 was constructed. By determining the nucleotide sequence of the modified gene, furthermore, the amino acid sequence of the modified Pfu DNA polymerase II (SQ ID No.6) was confirmed.

**[0331]** Preparation of the Gene of Modified Pfu DNA Polymerase III:

**[0332]** By the same method as described above except for the use of the oligonucleotides of SQ ID Nos.10 and 11, the gene of modified Pfu DNA polymerase III was prepared, to construct an expression vector pDP5C4. By determining the nucleotide sequence of the modified gene, the amino acid sequence (SQ ID No.7) of the modified Pfu DNA polymerase III was confirmed.

**[0333]** (2) Expression in *Escherichia coli* and purification of the modified Pfu DNA polymerases II and III

**[0334]** The genes of the modified Pfu DNA polymerases II and III, thus prepared, were expressed in *Escherichia coli* as follows, which were then purified.

**[0335]** The expression vectors pDP5b17 and pDP5C4 were independently inserted in a strain *Escherichia coli* HMS174 (DE3) and cultured in an LB culture medium supplemented with IPTG to a final concentration of 0.1 mM for 14 hours, to induce the expression of the enzymes in the bacteria of the strain *Escherichia coli*. After harvesting the bacteria by centrifugation, modified Pfu DNA polymerases II and III were extracted, with ultrasonic treatment, in a buffer containing 150 mM Tris/HCl (pH 7.5), 2 mM EDTA, 0.24 mM APMSF and 0.2% Tween 20. The crude extract solution was thermally treated at 80° C. for 15 minutes, to inactivate the DNA polymerases derived from *Escherichia coli* and partially purify the modified DNA polymerases II and III. The partially purified fractions were dialyzed against a buffer composed of 50 mM Tris/HCl (pH 7.5), 1 mM EDTA, 0.2% Tween 20, 7 mM 2-mercaptoethanol and 10% glycerol. At the stage were detected DNA polymerizing activities specific to the modified Pfu DNA polymerases II and III.

#### Example 9

**[0336]** By using the modified Pfu DNA polymerases II and III partially purified in Example 8, the primer elongation reaction of a DNA chain complementary to the template DNA was tested.

[0337] One  $\mu\text{g}$  of each of the partially purified enzyme fractions described above was placed in 20  $\mu\text{l}$  of a reaction solution containing 20 mM Tris/HCl (pH 8.0), 2 mM  $\text{MgCl}_2$ , 50  $\mu\text{g/ml}$  BSA, 0.1% Triton X-100, 1 mM each of cold dNTPs (0.1 mM for dCTP), 0.63  $\mu\text{g}$  of pBLUESCRIPT plasmid prepared by annealing together 10  $\mu\text{Ci}$  of [ $\alpha$ - $^{32}\text{P}$ ] dCTP and a primer M13(-21), for reaction at 75° C. for one minute and 3 minutes. The elongated DNA chain was separated by electrophoresis on a polyacrylamide gel containing 8M urea, and the resulting pattern was analyzed with an image analyzer. As a control, additionally, the conventional wild-type Pfu DNA polymerase was used for the same DNA synthesis.

[0338] The results are shown in FIG. 24. When the conventional wild-type Pfu DNA polymerase was used, bands indicating the presence of incomplete DNA chains were present under observation, because of the presence of a large region at about 1000 bases where synthetic termination occurred. However, the yield of synthesized products including those of bands of about 1000 bases was elevated during the DNA synthesis with the modified Pfu DNApoly-

merases II and III of the present invention, together with bands indicating the presence of more polymeric (more elongated) PCR products under observation.

[0339] The results described above indicate that the DNA polymerases of the present invention can more markedly elongate DNA chains in the course of synthesis during the DNA synthesis by PCR.

#### INDUSTRIAL APPLICABILITY

[0340] In accordance with the present invention, the functional site of a functionally unknown protein recovered by genome analysis or cDNA analysis can be predicted. A novel functional site of a protein with a known function can also be predicted.

[0341] The thermophilic DNA polymerases provided by the present invention can highly efficiently synthesize and amplify the whole length of a polymeric DNA by PCR, whereby the in vitro synthesis and amplification of a DNA chain and the nucleotide sequencing thereof can be attained at a high precision in a simple manner.

---

#### SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 11

<210> SEQ ID NO 1

<211> LENGTH: 775

<212> TYPE: PRT

<213> ORGANISM: Unknown

<220> FEATURE:

<223> OTHER INFORMATION: Description of Unknown Organism: DNA polymerase

<400> SEQUENCE: 1

```

Met Val Leu Asp Val Asp Tyr Ile Thr Glu Glu Gly Lys Pro Val Ile
 1                               10                    15
Arg Leu Phe Lys Lys Glu Asn Gly Lys Phe Lys Ile Glu His Asp Arg
 20                    25                    30
Thr Phe Arg Pro Tyr Ile Tyr Ala Leu Leu Arg Asp Ser Lys Ile
 35                    40                    45
Glu Glu Val Lys Lys Ile Thr Gly Glu Arg His Gly Lys Ile Val Arg
 50                    55                    60
Ile Val Asp Val Glu Lys Val Glu Lys Lys Phe Leu Gly Lys Pro Ile
 65                    70                    75
Thr Val Trp Lys Leu Tyr Leu Glu His Pro Gln Asp Val Pro Thr Ile
 85                    90                    95
Arg Glu Lys Val Arg Glu His Pro Ala Val Val Asp Ile Phe Glu Tyr
100                    105                    110
Asp Ile Pro Phe Ala Lys Arg Tyr Leu Ile Asp Lys Gly Leu Ile Pro
115                    120                    125
Met Glu Gly Glu Glu Glu Leu Lys Ile Leu Ala Phe Asp Ile Glu Thr
130                    135                    140
Leu Tyr His Glu Gly Glu Glu Phe Gly Lys Gly Pro Ile Ile Met Ile
145                    150                    155
Ser Tyr Ala Asp Glu Asn Glu Ala Lys Val Ile Thr Trp Lys Asn Ile
165                    170                    175
Asp Leu Pro Tyr Val Glu Val Val Ser Ser Glu Arg Glu Met Ile Lys

```

-continued

180				185				190							
Arg	Phe	Leu	Arg	Ile	Ile	Arg	Glu	Lys	Asp	Pro	Asp	Ile	Ile	Val	Thr
		195					200					205			
Tyr	Asn	Gly	Asp	Ser	Phe	Asp	Phe	Pro	Tyr	Leu	Ala	Lys	Arg	Ala	Glu
	210					215					220				
Lys	Leu	Gly	Ile	Lys	Leu	Thr	Ile	Gly	Arg	Asp	Gly	Ser	Glu	Pro	Lys
	225				230					235					240
Met	Gln	Arg	Ile	Gly	Asp	Met	Thr	Ala	Val	Glu	Val	Lys	Gly	Arg	Ile
			245							250				255	
His	Phe	Asp	Leu	Tyr	His	Val	Ile	Thr	Arg	Thr	Ile	Asn	Leu	Pro	Thr
			260						265				270		
Tyr	Thr	Leu	Glu	Ala	Val	Tyr	Glu	Ala	Ile	Phe	Gly	Lys	Pro	Lys	Glu
		275					280					285			
Lys	Val	Tyr	Ala	Asp	Glu	Ile	Ala	Lys	Ala	Trp	Glu	Ser	Gly	Glu	Asn
	290					295					300				
Leu	Glu	Arg	Val	Ala	Lys	Tyr	Ser	Met	Glu	Asp	Ala	Lys	Ala	Thr	Tyr
	305				310					315					320
Glu	Leu	Gly	Lys	Glu	Phe	Leu	Pro	Met	Glu	Ile	Gln	Leu	Ser	Arg	Leu
			325						330					335	
Val	Gly	Gln	Pro	Leu	Trp	Asp	Val	Ser	Arg	Ser	Ser	Thr	Gly	Asn	Leu
			340						345				350		
Val	Glu	Trp	Phe	Leu	Leu	Arg	Lys	Ala	Tyr	Glu	Arg	Asn	Glu	Val	Ala
		355					360					365			
Pro	Asn	Lys	Pro	Ser	Glu	Glu	Glu	Tyr	Gln	Arg	Arg	Leu	Arg	Glu	Ser
	370					375					380				
Tyr	Thr	Gly	Gly	Phe	Val	Lys	Glu	Pro	Glu	Lys	Gly	Leu	Trp	Glu	Asn
	385				390					395					400
Ile	Val	Tyr	Leu	Asp	Phe	Arg	Ala	Leu	Tyr	Pro	Ser	Ile	Ile	Ile	Thr
			405						410					415	
His	Asn	Val	Ser	Pro	Asp	Thr	Leu	Asn	Leu	Glu	Gly	Cys	Lys	Asn	Tyr
			420						425				430		
Asp	Ile	Ala	Pro	Gln	Val	Gly	His	Lys	Phe	Cys	Lys	Asp	Ile	Pro	Gly
		435					440					445			
Phe	Ile	Pro	Ser	Leu	Leu	Gly	His	Leu	Leu	Glu	Glu	Arg	Gln	Lys	Ile
	450					455					460				
Lys	Thr	Lys	Met	Lys	Glu	Thr	Gln	Asp	Pro	Ile	Glu	Lys	Ile	Leu	Leu
	465				470					475					480
Asp	Tyr	Arg	Gln	Lys	Ala	Ile	Lys	Leu	Leu	Ala	Asn	Ser	Phe	Tyr	Gly
			485						490					495	
Tyr	Tyr	Gly	Tyr	Ala	Lys	Ala	Arg	Trp	Tyr	Cys	Lys	Glu	Cys	Ala	Glu
			500						505				510		
Ser	Val	Thr	Ala	Trp	Gly	Arg	Lys	Tyr	Ile	Glu	Leu	Val	Trp	Lys	Glu
		515					520					525			
Leu	Glu	Glu	Lys	Tyr	Gly	Phe	Lys	Val	Ile	Tyr	Ser	Asp	Thr	Asp	Gly
	530					535					540				
Phe	Phe	Ala	Thr	Ile	Pro	Gly	Gly	Glu	Ser	Glu	Glu	Ile	Lys	Lys	Lys
	545				550					555					560
Ala	Leu	Glu	Phe	Val	Lys	Tyr	Ile	Asn	Ser	Lys	Leu	Pro	Gly	Leu	Leu
			565						570					575	
Glu	Leu	Glu	Tyr	Glu	Gly	Phe	Tyr	Lys	Arg	Gly	Phe	Phe	Val	Thr	Lys
			580						585				590		

-continued

Lys Arg Tyr Ala Val Ile Asp Glu Glu Gly Lys Val Ile Thr Arg Gly  
595 600 605

Leu Glu Ile Val Arg Arg Asp Trp Ser Glu Ile Ala Lys Glu Thr Gln  
610 615 620

Ala Arg Val Leu Glu Thr Ile Leu Lys His Gly Asp Val Glu Glu Ala  
625 630 635 640

Val Arg Ile Val Lys Glu Val Ile Gln Lys Leu Ala Asn Tyr Glu Ile  
645 650 655

Pro Pro Glu Lys Leu Ala Ile Tyr Glu Gln Ile Thr Arg Pro Leu His  
660 665 670

Glu Tyr Lys Ala Ile Gly Pro His Val Ala Val Ala Lys Lys Leu Ala  
675 680 685

Ala Lys Gly Val Lys Ile Lys Pro Gly Met Val Ile Gly Tyr Ile Val  
690 695 700

Leu Arg Gly Asp Gly Pro Ile Ser Asn Arg Ala Ile Leu Ala Glu Glu  
705 710 715 720

Tyr Asp Pro Lys Lys His Lys Tyr Asp Ala Glu Tyr Tyr Ile Glu Asn  
725 730 735

Gln Val Leu Pro Ala Val Leu Arg Ile Leu Glu Gly Phe Gly Tyr Arg  
740 745 750

Lys Glu Asp Leu Arg Tyr Gln Lys Thr Arg Gln Val Gly Leu Thr Ser  
755 760 765

Trp Leu Asn Ile Lys Lys Ser  
770 775

<210> SEQ ID NO 2

<211> LENGTH: 35

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic DNA

<400> SEQUENCE: 2

gtggggagca ccatggtttt agatgtggat tacat 35

<210> SEQ ID NO 3

<211> LENGTH: 35

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic DNA

<400> SEQUENCE: 3

gcatgcagat agaccatttc taacgaagc gtttg 35

<210> SEQ ID NO 4

<211> LENGTH: 66

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic DNA

<400> SEQUENCE: 4

ctcgaagaaa agtatggatt taaagtcac tacagtgaca ctgatggtt ctttgcaact 60

-continued

---

atccca 66

<210> SEQ ID NO 5  
 <211> LENGTH: 66  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 DNA

<400> SEQUENCE: 5

tgggatagtt gcaaagaac catcagtgtc actgtagatg actttaaatc catacttttc 60  
 ttcgag 66

<210> SEQ ID NO 6  
 <211> LENGTH: 775  
 <212> TYPE: PRT  
 <213> ORGANISM: Unknown  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Unknown Organism: DNA polymerase

<400> SEQUENCE: 6

Met Val Leu Asp Val Asp Tyr Ile Thr Glu Glu Gly Lys Pro Val Ile  
 1 5 10 15  
 Arg Leu Phe Lys Lys Glu Asn Gly Lys Phe Lys Ile Glu His Asp Arg  
 20 25 30  
 Thr Phe Arg Pro Tyr Ile Tyr Ala Leu Leu Arg Asp Asp Ser Lys Ile  
 35 40 45  
 Glu Glu Val Lys Lys Ile Thr Gly Glu Arg His Gly Lys Ile Val Arg  
 50 55 60  
 Ile Val Asp Val Glu Lys Val Glu Lys Lys Phe Leu Gly Lys Pro Ile  
 65 70 75 80  
 Thr Val Trp Lys Leu Tyr Leu Glu His Pro Gln Asp Val Pro Thr Ile  
 85 90 95  
 Arg Glu Lys Val Arg Glu His Pro Ala Val Val Asp Ile Phe Glu Tyr  
 100 105 110  
 Asp Ile Pro Phe Ala Lys Arg Tyr Leu Ile Asp Lys Gly Leu Ile Pro  
 115 120 125  
 Met Glu Gly Glu Glu Glu Leu Lys Ile Leu Ala Phe Asp Ile Glu Thr  
 130 135 140  
 Leu Tyr His Glu Gly Glu Glu Phe Gly Lys Gly Pro Ile Ile Met Ile  
 145 150 155 160  
 Ser Tyr Ala Asp Glu Asn Glu Ala Lys Val Ile Thr Trp Lys Asn Ile  
 165 170 175  
 Asp Leu Pro Tyr Val Glu Val Val Ser Ser Glu Arg Glu Met Ile Lys  
 180 185 190  
 Arg Phe Leu Arg Ile Ile Arg Glu Lys Asp Pro Asp Ile Ile Val Thr  
 195 200 205  
 Tyr Asn Gly Asp Ser Phe Asp Phe Pro Tyr Leu Ala Lys Arg Ala Glu  
 210 215 220  
 Lys Leu Gly Ile Lys Leu Thr Ile Gly Arg Asp Gly Ser Glu Pro Lys  
 225 230 235 240  
 Met Gln Arg Ile Gly Asp Met Thr Ala Val Glu Val Lys Gly Arg Ile  
 245 250 255  
 His Phe Asp Leu Tyr His Val Ile Thr Arg Thr Ile Asn Leu Pro Thr

-continued

260					265					270					
Tyr	Thr	Leu	Glu	Ala	Val	Tyr	Glu	Ala	Ile	Phe	Gly	Lys	Pro	Lys	Glu
		275					280					285			
Lys	Val	Tyr	Ala	Asp	Glu	Ile	Ala	Lys	Ala	Trp	Glu	Ser	Gly	Glu	Asn
	290					295					300				
Leu	Glu	Arg	Val	Ala	Lys	Tyr	Ser	Met	Glu	Asp	Ala	Lys	Ala	Thr	Tyr
	305					310					315				320
Glu	Leu	Gly	Lys	Glu	Phe	Leu	Pro	Met	Glu	Ile	Gln	Leu	Ser	Arg	Leu
				325					330					335	
Val	Gly	Gln	Pro	Leu	Trp	Asp	Val	Ser	Arg	Ser	Ser	Thr	Gly	Asn	Leu
			340					345					350		
Val	Glu	Trp	Phe	Leu	Leu	Arg	Lys	Ala	Tyr	Glu	Arg	Asn	Glu	Val	Ala
		355					360					365			
Pro	Asn	Lys	Pro	Ser	Glu	Glu	Glu	Tyr	Gln	Arg	Arg	Leu	Arg	Glu	Ser
	370					375					380				
Tyr	Thr	Gly	Gly	Phe	Val	Lys	Glu	Pro	Glu	Lys	Gly	Leu	Trp	Glu	Asn
	385					390					395				400
Ile	Val	Tyr	Leu	Asp	Phe	Arg	Ala	Leu	Tyr	Pro	Ser	Ile	Ile	Ile	Thr
				405					410					415	
His	Asn	Val	Ser	Pro	Asp	Thr	Leu	Asn	Leu	Glu	Gly	Cys	Lys	Asn	Tyr
			420					425					430		
Asp	Ile	Ala	Pro	Gln	Val	Gly	His	Lys	Phe	Cys	Lys	Asp	Ile	Pro	Gly
		435					440					445			
Phe	Ile	Pro	Ser	Leu	Leu	Gly	His	Leu	Leu	Glu	Glu	Arg	Gln	Lys	Ile
	450					455					460				
Lys	Thr	Lys	Met	Lys	Glu	Thr	Gln	Asp	Pro	Ile	Glu	Lys	Ile	Leu	Leu
	465					470					475				480
Asp	Tyr	Arg	Gln	Lys	Ala	Ile	Lys	Leu	Leu	Ala	Asn	Ser	Phe	Tyr	Gly
				485					490					495	
Tyr	Tyr	Gly	Tyr	Ala	Lys	Ala	Arg	Trp	Tyr	Cys	Lys	Glu	Cys	Ala	Glu
			500					505					510		
Ser	Val	Thr	Ala	Trp	Gly	Arg	Lys	Tyr	Ile	Glu	Leu	Val	Trp	Lys	Glu
		515					520					525			
Leu	Glu	Glu	Lys	Phe	Gly	Phe	Lys	Val	Leu	Tyr	Ile	Asp	Thr	Asp	Gly
	530					535					540				
Leu	Tyr	Ala	Thr	Ile	Pro	Gly	Gly	Glu	Ser	Glu	Glu	Ile	Lys	Lys	Lys
	545					550					555				560
Ala	Leu	Glu	Phe	Val	Lys	Tyr	Ile	Asn	Ser	Lys	Leu	Pro	Gly	Leu	Leu
				565					570					575	
Glu	Leu	Glu	Tyr	Glu	Gly	Phe	Tyr	Lys	Arg	Gly	Phe	Phe	Val	Thr	Lys
			580					585					590		
Lys	Arg	Tyr	Ala	Val	Ile	Asp	Glu	Glu	Gly	Lys	Val	Ile	Thr	Arg	Gly
		595					600					605			
Leu	Glu	Ile	Val	Arg	Arg	Asp	Trp	Ser	Glu	Ile	Ala	Lys	Glu	Thr	Gln
	610					615					620				
Ala	Arg	Val	Leu	Glu	Thr	Ile	Leu	Lys	His	Gly	Asp	Val	Glu	Glu	Ala
	625					630					635				640
Val	Arg	Ile	Val	Lys	Glu	Val	Ile	Gln	Lys	Leu	Ala	Asn	Tyr	Glu	Ile
				645					650					655	
Pro	Pro	Glu	Lys	Leu	Ala	Ile	Tyr	Glu	Gln	Ile	Thr	Arg	Pro	Leu	His
			660					665					670		

-continued

---

Glu Tyr Lys Ala Ile Gly Pro His Val Ala Val Ala Lys Lys Leu Ala  
 675 680 685  
 Ala Lys Gly Val Lys Ile Lys Pro Gly Met Val Ile Gly Tyr Ile Val  
 690 695 700  
 Leu Arg Gly Asp Gly Arg Ile Arg Asp Arg Ala Ile Pro Ala Glu Glu  
 705 710 715 720  
 Tyr Asp Pro Lys Lys His Lys Tyr Asp Ala Glu Tyr Tyr Ile Glu Asn  
 725 730 735  
 Gln Val Leu Pro Ala Val Leu Arg Ile Leu Glu Gly Phe Gly Tyr Arg  
 740 745 750  
 Lys Glu Asp Leu Arg Tyr Gln Lys Thr Arg Gln Val Gly Leu Thr Ser  
 755 760 765  
 Trp Leu Asn Ile Lys Lys Ser  
 770 775

&lt;210&gt; SEQ ID NO 7

&lt;211&gt; LENGTH: 775

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Unknown

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Unknown Organism: DNA polymerase

&lt;400&gt; SEQUENCE: 7

Met Val Leu Asp Val Asp Tyr Ile Thr Glu Glu Gly Lys Pro Val Ile  
 1 5 10 15  
 Arg Leu Phe Lys Lys Glu Asn Gly Lys Phe Lys Ile Glu His Asp Arg  
 20 25 30  
 Thr Phe Arg Pro Tyr Ile Tyr Ala Leu Leu Arg Asp Asp Ser Lys Ile  
 35 40 45  
 Glu Glu Val Lys Lys Ile Thr Gly Glu Arg His Gly Lys Ile Val Arg  
 50 55 60  
 Ile Val Asp Val Glu Lys Val Glu Lys Lys Phe Leu Gly Lys Pro Ile  
 65 70 75 80  
 Thr Val Trp Lys Leu Tyr Leu Glu His Pro Gln Asp Val Pro Thr Ile  
 85 90 95  
 Arg Glu Lys Val Arg Glu His Pro Ala Val Val Asp Ile Phe Glu Tyr  
 100 105 110  
 Asp Ile Pro Phe Ala Lys Arg Tyr Leu Ile Asp Lys Gly Leu Ile Pro  
 115 120 125  
 Met Glu Gly Glu Glu Glu Leu Lys Ile Leu Ala Phe Asp Ile Glu Thr  
 130 135 140  
 Leu Tyr His Glu Gly Glu Glu Phe Gly Lys Gly Pro Ile Ile Met Ile  
 145 150 155 160  
 Ser Tyr Ala Asp Glu Asn Glu Ala Lys Val Ile Thr Trp Lys Asn Ile  
 165 170 175  
 Asp Leu Pro Tyr Val Glu Val Val Ser Ser Glu Arg Glu Met Ile Lys  
 180 185 190  
 Arg Phe Leu Arg Ile Ile Arg Glu Lys Asp Pro Asp Ile Ile Val Thr  
 195 200 205  
 Tyr Asn Gly Asp Ser Phe Asp Phe Pro Tyr Leu Ala Lys Arg Ala Glu  
 210 215 220  
 Lys Leu Gly Ile Lys Leu Thr Ile Gly Arg Asp Gly Ser Glu Pro Lys  
 225 230 235 240

-continued

---

Met Gln Arg Ile Gly Asp Met Thr Ala Val Glu Val Lys Gly Arg Ile  
 245 250 255  
 His Phe Asp Leu Tyr His Val Ile Thr Arg Thr Ile Asn Leu Pro Thr  
 260 265 270  
 Tyr Thr Leu Glu Ala Val Tyr Glu Ala Ile Phe Gly Lys Pro Lys Glu  
 275 280 285  
 Lys Val Tyr Ala Asp Glu Ile Ala Lys Ala Trp Glu Ser Gly Glu Asn  
 290 295 300  
 Leu Glu Arg Val Ala Lys Tyr Ser Met Glu Asp Ala Lys Ala Thr Tyr  
 305 310 315 320  
 Glu Leu Gly Lys Glu Phe Leu Pro Met Glu Ile Gln Leu Ser Arg Leu  
 325 330 335  
 Val Gly Gln Pro Leu Trp Asp Val Ser Arg Ser Ser Thr Gly Asn Leu  
 340 345 350  
 Val Glu Trp Phe Leu Leu Arg Lys Ala Tyr Glu Arg Asn Glu Val Ala  
 355 360 365  
 Pro Asn Lys Pro Ser Glu Glu Tyr Gln Arg Arg Leu Arg Glu Ser  
 370 375 380  
 Tyr Thr Gly Gly Phe Val Lys Glu Pro Glu Lys Gly Leu Trp Glu Asn  
 385 390 395 400  
 Ile Val Tyr Leu Asp Phe Arg Ala Leu Tyr Pro Ser Ile Ile Ile Thr  
 405 410 415  
 His Asn Val Ser Pro Asp Thr Leu Asn Leu Glu Gly Cys Lys Asn Tyr  
 420 425 430  
 Asp Ile Ala Pro Gln Val Gly His Lys Phe Cys Lys Asp Ile Pro Gly  
 435 440 445  
 Phe Ile Pro Ser Leu Leu Gly His Leu Leu Glu Glu Arg Gln Lys Ile  
 450 455 460  
 Lys Thr Lys Met Lys Glu Thr Gln Asp Pro Ile Glu Lys Ile Leu Leu  
 465 470 475 480  
 Asp Tyr Arg Gln Lys Ala Ile Lys Leu Leu Ala Asn Ser Phe Tyr Gly  
 485 490 495  
 Tyr Tyr Gly Tyr Ala Lys Ala Arg Trp Tyr Cys Lys Glu Cys Ala Glu  
 500 505 510  
 Ser Val Thr Ala Trp Gly Arg Lys Tyr Ile Glu Leu Val Trp Lys Glu  
 515 520 525  
 Leu Glu Glu Lys Phe Gly Phe Lys Val Leu Tyr Ile Asp Thr Asp Gly  
 530 535 540  
 Leu Tyr Ala Thr Ile Pro Gly Gly Glu Ser Glu Glu Ile Lys Lys Lys  
 545 550 555 560  
 Ala Leu Glu Phe Val Lys Tyr Ile Asn Ser Lys Leu Pro Gly Leu Leu  
 565 570 575  
 Glu Leu Glu Tyr Glu Gly Phe Tyr Lys Arg Gly Phe Phe Val Thr Lys  
 580 585 590  
 Lys Arg Tyr Ala Val Ile Asp Glu Glu Gly Lys Val Ile Thr Arg Gly  
 595 600 605  
 Leu Glu Ile Val Arg Arg Asp Trp Ser Glu Ile Ala Lys Glu Thr Gln  
 610 615 620  
 Ala Arg Val Leu Glu Thr Ile Leu Lys His Gly Asp Val Glu Glu Ala  
 625 630 635 640

-continued

---

Val	Arg	Ile	Val	Lys	Glu	Val	Ile	Gln	Lys	Leu	Ala	Asn	Tyr	Glu	Ile
			645						650					655	
Pro	Pro	Glu	Lys	Leu	Ala	Ile	Tyr	Glu	Gln	Ile	Thr	Arg	Pro	Leu	His
		660						665					670		
Glu	Tyr	Lys	Ala	Ile	Gly	Pro	His	Val	Ala	Val	Ala	Lys	Lys	Leu	Ala
		675					680						685		
Ala	Lys	Gly	Val	Lys	Ile	Lys	Pro	Gly	Met	Val	Ile	Gly	Tyr	Ile	Val
		690				695					700				
Leu	Arg	Gly	Asp	Gly	Pro	Ile	Ser	Asn	Arg	Ala	Ile	Pro	Ala	Glu	Glu
705					710					715				720	
Tyr	Asp	Pro	Lys	Lys	His	Lys	Tyr	Asp	Ala	Glu	Tyr	Tyr	Ile	Glu	Asn
				725					730					735	
Gln	Val	Leu	Pro	Ala	Val	Leu	Arg	Ile	Leu	Glu	Gly	Phe	Gly	Tyr	Arg
			740					745						750	
Lys	Glu	Asp	Leu	Arg	Tyr	Gln	Lys	Thr	Arg	Gln	Val	Gly	Leu	Thr	Ser
		755					760						765		
Trp	Leu	Asn	Ile	Lys	Lys	Ser									
	770					775									

<210> SEQ ID NO 8  
 <211> LENGTH: 49  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic DNA

<400> SEQUENCE: 8

agaggcgatg gtcgaattcg cgataggcca attccagctg aggaatacy 49

<210> SEQ ID NO 9  
 <211> LENGTH: 49  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic DNA

<400> SEQUENCE: 9

cgtattcctc agctggaatt gccctatcgc gaattcgacc atcgcctct 49

<210> SEQ ID NO 10  
 <211> LENGTH: 40  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic DNA

<400> SEQUENCE: 10

ccaattagca atagggcaat tccagctgag gaatcagatc 40

<210> SEQ ID NO 11  
 <211> LENGTH: 40  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic DNA

-continued

<400> SEQUENCE: 11

gatcgtattc ctcagctgga attgcctat tgctaattgg

40

1. A method for predicting a functional site of a protein derived from the entire putative proteins of an organism "a" of which genome data or cDNA data is known, which method comprises the steps of:

- (1) determining in the amino acid sequences of the entire proteins of the organism "a", the frequency of occurrence of each amino acid and the frequency of occurrence of individual oligopeptides produced by permutations of twenty amino acids, and determining the smallest length (n) of oligopeptides satisfying the following criteria;

among oligopeptides of length (n), the number of oligopeptides which occur once in the entire proteins is smaller than the number of oligopeptides which occur twice in the entire proteins; among oligopeptides of length (n+1), the number of oligopeptides which occur once in the entire proteins is larger than the number of oligopeptides which occur twice in the entire proteins,

- (2) determining in the entire proteins of the organism "a", the frequency of occurrence of the following Aji-oligopeptide of length (n+1), which is a part of the amino acid sequence of the protein as a subject for predicting a functional site and contains the j-th amino-acid residue Aj ( $n+1 \leq j \leq L-n$ ) from the amino terminal end (N-terminus) of the amino acid sequence (length of L) of the protein, on condition that the j-th amino-acid residue Aj is defined as the i-th residue Aji from the N-terminus of the Aji-oligopeptide;

Aji-oligopeptide:  $aj_1aj_2 \dots Aji \dots aj_naj_{n+1}$   
(wherein,  $1 \leq i \leq n+1$ ;  $A_j = A_{ji}$ ; and Aj is the i-th residue of the oligopeptide; and moreover,  $aj_1 = A_{j-i+1}, \dots, aj_{n+1} = A_{j-i+(n+1)}$ ), and determining in the entire proteins of the organism "a", the frequency of occurrence of the following Xji-oligopeptide of length (n+1);

Xji-oligopeptide:  $aj_1aj_2 \dots Xji \dots aj_naj_{n+1}$   
(wherein,  $1 \leq i \leq n+1$ ;  $n+1 \leq j \leq L-n$ ; and the i-th residue Xji is any amino acid; and moreover,  $aj_1 = A_{j-i+1}, \dots, aj_{n+1} = A_{j-i+(n+1)}$ ),

- (3) calculating ratio Yji of the frequency of occurrence of the Aji-oligopeptide to that of the Xji-oligopeptide,  
(4) determining mean Yj of the Yji;

$$Y_j = \sum_{i=1}^{n+1} Y_{ji} / (n+1),$$

- (5) determining functional value Zj of Yj;

$Z_j = f(Y_j)$  (Function f is a monotonously decreasing function or a monotonously increasing function),

and defining the Zj value as the representative value of the function of the j-th amino-acid residue Aj of the amino acid sequence (length of L), and

- (6) repeating the steps (2) to (5) sequentially and determining the Zj value of each Aj of all the amino-acid residues at positions between  $n+1 \leq j \leq L-n$  in the amino acid sequence (length of L), thereby predicting the degree of the involvement of each amino-acid residue in the function of the protein by using the dimension of the Zj value as an indicator.

2. The method according to claim 1, wherein the Zj value ( $n+1 \leq j \leq L-n$ ) of each amino-acid residue in the amino acid sequence (length of L) is expressed in a distribution chart.

3. A method for predicting a functional site of a protein derived from the entire putative proteins of an organism "a" of which genome data or cDNA data is known, which method comprises the steps of:

- (1) determining the frequency of occurrence of each amino acid and the frequency of occurrence of individual oligopeptides produced by permutations of twenty amino acids, in the amino acid sequences of the entire proteins of the organism "a",

- (2) as to an appropriate protein of the organism "a",

- (2') determining in the entire proteins of the organism "a", the frequency of occurrence of the following Aji-oligopeptide of given length of (n) ( $1 \leq n \leq M$ , provided that M is the the smallest length of oligopeptides satisfying the criterion that all the oligopeptides of length M are at frequency 1 of the occurrence), which Aji-oligopeptide is a part of the amino acid sequence of the protein and contains the j-th amino-acid residue Aj ( $n \leq j \leq L-n+1$ ), on condition that the j-th amino-acid residue Aj is defined as the i-th residue Aji from the N-terminus of the Aji-oligopeptide;

Aji-oligopeptide:  $aj_1aj_2 \dots aji \dots aj_n$  (wherein,  $1 \leq i \leq n+1$ ;  $A_j = a_{ji}$ , and Aj is the i-th residue of the oligopeptide; and moreover,  $aj_1 = A_{j-i+1}, \dots, aj_n = A_{j-i+n}$ ), and determining in the entire proteins of the organism "a", the frequency of occurrence of the following Xji-oligopeptide of length (n) corresponding to the length of Aji-oligopeptide;

Xji-oligopeptide:  $aj_1aj_2 \dots Xji \dots aj_n$  (the i-th residue Xji is any amino-acid residue; and moreover,  $aj_1 = A_{j-i+1}, \dots, aj_n = A_{j-i+n}$ ),

- (3) calculating ratio Yji of the frequency of occurrence of the Aji-oligopeptide to that of the Xji-oligopeptide,

(4) determining mean  $Y(j,n)$  of the  $Y_{ji}$ ;

$$Y(j, n) = \sum_{i=1}^n Y_{ji}/n,$$

(5) determining functional value  $Z(j,n)$  of  $Y(j,n)$ ;

$$Z(j, n) = -\log(Y(j,n)),$$

(6) repeating the steps (2') to (5) sequentially and determining the  $Z(j,n)$  value of each amino-acid residue  $A_j$  at position  $j$  ( $n \leq j \leq L-n+1$ ) in the amino acid sequence (length of  $L$ ),

(7) sequentially repeating the steps (2) to (6) for the entire proteins of the organism "a", thereby determining the distribution of the  $Z(j,n)$  value of each amino-acid residue in the entire proteins, and the  $Z(j,n)$  values are classified into twenty according to the twenty amino acids, and then determining mean  $Av(Aa)$  of the  $Z(j,n)$  values for each amino acid  $Aa$  and the standard deviation  $Sd(Aa)$  of the distribution thereof, on the basis of the distribution, to determine function  $g$  to the  $j$ -th amino-acid residue  $A_j$  of a protein for normalizing the difference in distribution due to the species of amino-acid residues;

$$g = g(Z(j,n), A_j) = [Z(j,n) - Av(Aa)] / Sd(Aa) \text{ (provided that } A_j \text{ Aa; and } 1 \leq n \leq M),$$

(8) determining value  $D(j,n)$  of the function  $g$  of each  $A_j$  of all the amino-acid residues at position  $j$  ( $n \leq j \leq L-n+1$  and  $1 \leq n \leq M$ ) of a protein in the entire proteins as recovered in the step (7);

$$D(j,n) = g(Z(j,n), A_j),$$

and

(9) defining the representative value of the function of the  $j$ -th amino-acid residue in the amino acid sequence (length of  $L$ ) as functional value  $W_j$  of the  $Z(j,n)$  and  $D(j,n)$ ;

$$W_j = h(Z(j,1), Z(j,2), \dots, Z(j,M), D(j,1), D(j,2), \dots, D(j,M))$$

thereby predicting the degree of the involvement of each amino-acid residue in the function of the protein by using the dimension of the  $W_j$  value as an indicator.

4. The method according to claim 3, wherein the  $W_j$  value of each amino-acid residue is expressed in a two-dimensional distribution chart.

5. The method according to claim 3, wherein the  $W_j$  value of each amino-acid residue is expressed on a three-dimensional structure model of the protein.

6. A system for automatically conducting the method according to claim 1, at least comprising the following units (a) to (g);

(a) an outer memory unit memorizing the amino acid sequence data of the entire putative proteins of organism "a" of which genome data or cDNA data is known, as well as an existing protein data base,

(b) a calculation/memory unit, composed of CPU calculating the frequency of occurrence of each amino acid

and the frequency of occurrence of individual oligopeptides produced by permutations of twenty amino acids, in the amino acid sequences of the entire proteins of the organism "a", and a memory unit having the memory of the calculation results,

(c) a calculation/memory unit, composed of CPU calculating the smallest length ( $n$ ) of oligopeptides satisfying the following criteria among the individual oligopeptides of which the frequencies of the occurrences being memorized in the unit (b);

among oligopeptides of length ( $n$ ), the number of oligopeptides which occur once in the entire proteins is smaller than the number of oligopeptides which occur twice in the entire proteins; among oligopeptides of length ( $n+1$ ), the number of oligopeptides which occur once in the entire proteins is larger than the number of oligopeptides which occur twice in the entire proteins,

and a memory unit having the memory of the ( $n$ ),

(d) a calculation/memory unit, composed of CPU determining in the entire proteins of the organism "a", the frequency of occurrence of the following  $A_{ji}$ -oligopeptide of length ( $n+1$ ), which is a part of the amino acid sequence of the protein as a subject for predicting a functional site and contains the  $j$ -th amino-acid residue  $A_j$  ( $n+1 \leq j \leq L-n$ ) from the amino terminal end (N-terminus) of the amino acid sequence (length of  $L$ ) of the protein, on condition that the  $j$ -th amino-acid residue  $A_j$  is defined as (or identical to?) the  $i$ -th residue  $A_{ji}$  from the N-terminus of the  $A_{ji}$ -oligopeptide;

$$A_{ji}\text{-oligopeptide: } a_1 a_2 \dots a_{j-1} A_j \dots a_{n+1} \text{ (} 1 \leq n+1; A_j = A_{ji}; \text{ and } A_j \text{ is the } i\text{-th residue of the oligopeptide; and moreover, } a_{j-1} = A_{j-1}, \dots, a_{n+1} = A_{n+1})$$

and determining in the entire proteins of the organism "a", the frequency of occurrence of the following  $X_{ji}$ -oligopeptide of length ( $n+1$ );

$$X_{ji}\text{-oligopeptide: } a_1 a_2 \dots X_{ji} \dots a_{n+1} \text{ (} 1 \leq i \leq n+1; n+1 \leq j \leq L-n; \text{ and the } i\text{-th residue } X_{ji} \text{ is any amino acid)}$$

and a memory unit having the memory of the calculation results,

(e) a calculation/memory unit, composed of CPU determining ratio  $Y_{ji}$  of the frequency of occurrence of the  $A_{ji}$ -oligopeptide to that of the  $X_{ji}$ -oligopeptide, and a memory unit having the memory of  $Y_{ji}$ ,

(f) a calculation/memory unit, composed of CPU determining mean  $Y_j$  of the  $Y_{ji}$ ;

$$Y_j = \sum_{i=1}^{n+1} Y_{ji} / (n+1),$$

and a memory unit having the memory of  $Y_j$ , and

(g) a calculation/memory unit, composed of CPU determining functional value  $Z_j$  of  $Y_j$ ;

$Z_j=f(Y_j)$  (Function  $f$  is a monotonously decreasing function or a monotonously increasing function),

and a memory unit having the memory of  $Z_j$ .

7. The system according to claim 6, the system being equipped with a display unit displaying the  $Z_j$  value ( $n+1 \leq j \leq L-n$ ) of each amino-acid residue in the amino acid sequence (length of  $L$ ) in a distribution chart.

8. A system for automatically conducting the method according to claim 3, at least comprising the following units (a) to (i);

- (a) an outer memory unit memorizing the amino acid sequence data of the entire putative proteins of organism "a" of which genome data or cDNA data is known, as well as an existing protein data base,
- (b) a calculation/memory unit, composed of CPU calculating in the amino acid sequences of the entire proteins of the organism "a", the frequency of occurrence of each amino acid and the frequency of occurrence of individual oligopeptides produced by permutations of twenty amino acids, and a memory unit having the memory of the calculation results,
- (c) a calculation/memory unit, composed of CPU determining in the entire proteins of the organism "a", the frequency of occurrence of the following  $A_{ji}$ -oligopeptide of given length of ( $n$ ) ( $1 \leq n \leq M$ , provided that  $M$  is the smallest length of oligopeptides satisfying the criterion that all the oligopeptides of length  $M$  are at frequency 1 of the occurrence), which  $A_{ji}$ -oligopeptide is a part of the amino acid sequence of a given protein of the organism "a" and contains the  $j$ -th amino-acid residue  $A_j$ , on condition that the  $j$ -th amino-acid residue  $A_j$  is defined as the  $i$ -th residue  $A_{ji}$  from the N-terminus of the  $A_{ji}$ -oligopeptide ( $n \leq j \leq L-n+1$ );

$A_{ji}$ -oligopeptide:  $a_j1 a_j2 \dots a_{ji} \dots a_{jn}$  ( $1 \leq n+1$ ;  $A_j=a_{ji}$ , and  $A_j$  is the  $i$ -th residue of the oligopeptide; and moreover,  $a_{j1}=A_j-i+1, \dots, a_{jn}=A_j-i+n$ ),

and determining in the entire proteins of the organism "a", the frequency of occurrence of the following  $X_{ji}$ -oligopeptide of length ( $n$ ) corresponding to the length of  $A_{ji}$ -oligopeptide;

$X_{ji}$ -oligopeptide:  $a_j1 a_j2 \dots X_{ji} \dots a_{jn}$  (the  $i$ -th residue  $X_{ji}$  is any amino acid; and moreover,  $a_{j1}=A_j-i+1, \dots, a_{jn}=A_j-i+n$ ), and a memory unit memorizing the calculation results,

- (d) a calculation/memory unit, composed of CPU calculating ratio  $Y_{ji}$  of the frequency of occurrence of the  $A_{ji}$ -oligopeptide to that of the  $X_{ji}$ -oligopeptide, and a memory unit having the memory of the  $Y_{ji}$ ,
- (e) a calculation/memory unit, composed of CPU determining mean  $Y(j,n)$  of the  $Y_{ji}$ ;

$$Y(j, n) = \sum_{i=1}^n Y_{ji} / n,$$

and a memory unit having the memory of  $Y(j,n)$ ,

(f) a calculation/memory unit, composed of CPU determining functional value  $Z(j,n)$  of  $Y(j,n)$ ;

$$Z(j,n)=-\log(Y(j,n)),$$

and a memory unit having the memory of  $Z(j,n)$ ,

(g) a calculation/memory unit, composed of CPU determining the  $Z(j,n)$  value of each amino-acid residue in the amino acid sequences of the entire proteins of the organism "a", determining the distribution of the  $Z(j,n)$  value of each amino-acid residue in the entire proteins, and the  $Z(j,n)$  values are classified into twenty according to the twenty amino acids, and then determining mean  $Av(Aa)$  of the  $Z(j,n)$  values for each amino acid  $Aa$  and the standard deviation  $Sd(Aa)$  of the distribution thereof, on the basis of the distribution, to determine function  $g$  for normalizing the difference in distribution due to the species of amino-acid residues;

$$g(Z(j,n), A_j)=[Z(j,n)-Av(Aa)]/Sd(Aa) \text{ (wherein, } A_j=Aa; \text{ and } 1 \leq n \leq M)$$

and a memory unit having the memory of  $g$ ,

(h) a calculation/memory unit, composed of CPU determining value  $D(j,n)$  of function  $g$  memorized in the unit (g) concerning each of all the amino-acid residues  $A_j$  at position  $j$  ( $n \leq j \leq L-n+1$ ) in the amino acid sequence (length of  $L$ );

$$D(j,n)=g(Z(j,n), A_j)$$

and a memory unit having the memory of the  $D(j, n)$  value, and

(i) a calculation/memory unit, composed of a calculation unit determining appropriate functional value  $W_j$  of the  $Z(j,n)$  and  $D(j,n)$  of each amino-acid residue in the amino acid sequence;

$$W_j=h(Z(j,1),Z(j,2), \dots, Z(j,M),D(j,1),D(j,2), \dots, D(j, M))$$

and a memory unit having the memory of the  $W_j$  value.

9. The system according to claim 8, the system being equipped with a display unit displaying the  $W_j$  value of each amino-acid residue in a two-dimensional distribution chart.

10. The system according to claim 8, the system being equipped with a calculation/memory unit memorizing an existing database of three-dimensional structure of proteins or preparing a three-dimensional structure model based on an amino acid sequence according to a known method and memorizing the three-dimensional structure model, and a display unit displaying the  $W_j$  value of each amino-acid residue in the amino acid sequence in a distribution chart on the three-dimensional structure stored in the data base or three-dimensional structure model memorized in the calculation/memory unit.

11. A method for modifying the known function of protein "A" derived from the entire proteins of organism "a" of which genome data or cDNA data has been known, which method comprises the steps of:

- (1) extracting a protein closely related to the protein "A" from an existing protein data base and subjecting the proteins to alignment,
- (2) determining in the amino acid sequences of the entire proteins of the organism "a", the frequency of occurrence of each amino acid and the frequency of occur-

rence of individual oligopeptides produced by permutations of twenty amino acids, and determining the smallest length (n) of oligopeptides satisfying the following criteria;

among oligopeptides of length (n), the number of oligopeptides which occurrence in the entire proteins is smaller than the number of oligopeptides which occur twice in the entire proteins; among oligopeptides of length (n+1), the number of oligopeptides which occur once in the entire proteins is larger than the number of oligopeptides which occur twice in the entire proteins,

- (3) determining in the entire proteins of the organism "a", the frequency of occurrence of the following Aji-oligopeptide of length (n+1), which is a part of the amino acid sequence of the protein as a subject for predicting a functional site and contains the j-th amino-acid residue Aj from the amino terminal end (N-terminus) of the amino acid sequence (length of L) of the protein, on condition that the j-th amino-acid residue Aj is defined as (or identical to?) the i-th residue Aji from the N-terminus of the Aji-oligopeptide ( $n+1 \leq j \leq L-n$ );

Aji-oligopeptide:  $aj_1aj_2 \dots Aji \dots aj_naj_{(n+1)}$   
(wherein,  $1 \leq i \leq n+1$ ;  $A_j = A_{ji}$ ; and  $A_j$  is the i-th residue of the oligopeptide; and moreover,  $aj_1 = A_{j-i+1}$ ,  $\dots$ ,  $aj_{(n+1)} = A_{j-i+(n+1)}$ ),

and determining in the entire proteins of the organism "a", the frequency of occurrence of the following Xji-oligopeptide of length (n+1);

Xji-oligopeptide:  $aj_1aj_2 \dots Xji \dots aj_naj_{(n+1)}$   
(wherein,  $1 \leq i \leq n+1$ ;  $n+1 \leq j \leq L-n$ ; and the i-th residue Xji is any amino acid; and moreover,  $aj_1 = A_{j-i+1}$ ,  $\dots$ ,  $aj_{(n+1)} = A_{j-i+(n+1)}$ ),

- (4) calculating ratio Yji of the frequency of occurrence of the Aji-oligopeptide to that of the Xji-oligopeptide,  
(5) determining mean Yj of the Yji;

$$Y_j = \sum_{i=1}^{n+1} Y_{ji} / (n+1),$$

- (6) determining functional value Zj of Yj;

$Z_j = f(Y_j)$  (Function f is a monotonously decreasing function or a monotonously increasing function),

and defining the Zj value as the representative value of the function of the j-th amino-acid residue of the amino acid sequence (length of L) of the protein "A",

- (7) sequentially repeating the steps (3) to (6) and determining the Zj value of each of all the amino-acid residues at position between  $n+1 \leq j \leq L-n$  in the amino acid sequence (length of L),  
(8) selecting at least one amino-acid residue to be subjected to mutation on the basis of the alignment data carried out in the step (1) from the amino acid sequence (length of L) of the protein "A", sequentially repeating the steps (3) to (6) for variant amino-acid residues in various mutated amino acid sequences where the

selected amino-acid residue has been mutated into another amino-acid residue, to determine the Zj value of the variant amino-acid residues,

- (9) selecting a mutated amino acid sequence wherein the Zj value of the variant amino-acid residue as determined in the step (8) is larger or smaller than the Zj value of the wild type amino-acid residue as determined in the step (7), and (10) preparing a modified gene encoding the modified amino acid sequence from the protein "A" gene, and producing the modified protein as the expression product of the gene.

12. A method for modifying the function of protein "B" derived from an organism "b" of which genome data or cDNA data has been unknown, which method comprises the steps of:

- (1) extracting protein "A" most closely related to protein "B" from the entire proteins of organism "a" of which genome data or cDNA data being known and subjecting the protein to alignment, or extracting a protein closely related to protein "B" from an existing protein data base to subject the protein to alignment,

- (2) determining in the amino acid sequences of the entire proteins of the organism "a", the frequency of occurrence of each amino acid and the frequency of occurrence of individual oligopeptides produced by permutations of twenty amino acids, and determining the smallest length (n) of oligopeptides satisfying the following criteria;

among oligopeptides of length (n), the number of oligopeptides which occur once in the entire proteins is smaller than the number of oligopeptides which occur twice in the entire proteins; among oligopeptides of length (n+1), the number of oligopeptides which occur once in the entire proteins is larger than the number of oligopeptides which occur twice in the entire proteins,

- (3) determining in the entire proteins of the organism "a", the frequency of occurrence of the following Aji-oligopeptide of length (n+1), which is a part of the amino acid sequence of the protein as a subject for predicting a functional site and contains the j-th amino-acid residue Aj from the amino terminal end (N-terminus) of the amino acid sequence (length of L) of the protein, on condition that the j-th amino-acid residue Aj is defined as the i-th residue Aji from the N-terminus of the Aji-oligopeptide ( $n+1 \leq j \leq L-n$ );

Aji-oligopeptide:  $aj_1aj_2 \dots Aji \dots aj_naj_{(n+1)}$   
(wherein,  $1 \leq i \leq n+1$ ;  $A_j = A_{ji}$ ; and  $A_j$  is the i-th residue of the oligopeptide; and moreover,  $aj_1 = A_{j-i+1}$ ,  $\dots$ ,  $aj_{(n+1)} = A_{j-i+(n+1)}$ ),

and determining in the entire proteins of the organism "a", the frequency of occurrence of the following Xji-oligopeptide of length (n+1);

Xji-oligopeptide:  $aj_1aj_2 \dots Xji \dots aj_naj_{(n+1)}$   
(wherein,  $1 \leq i \leq n+1$ ;  $n+1 \leq j \leq L-n$ ; and the i-th residue Xji is any amino acid; and moreover,  $aj_1 = A_{j-i+1}$ ,  $\dots$ ,  $aj_{(n+1)} = A_{j-i+(n+1)}$ )

- (4) calculating ratio  $Y_{ji}$  of the frequency of occurrence of the  $A_{ji}$ -oligopeptide to that of the  $X_{ji}$ -oligopeptide,  
 (6) determining mean  $Y_j$  of the  $Y_{ji}$ ;

$$Y_j = \sum_{i=1}^{n+1} Y_{ji}/(n+1),$$

- (6) determining functional value  $Z_j$  of  $Y_j$ ;

$Z_j=f(Y_j)$  (Function  $f$  is a monotonously decreasing function or a monotonously increasing function),

and defining the  $Z_j$  value as the representative value of the function of the  $j$ -th amino-acid residue  $A_j$  of the amino acid sequence (length of  $L$ ) of the protein "A",

- (7) sequentially repeating the steps (3) to (6) and determining the  $Z_j$  value of each of all the amino-acid residues at position between  $n+1 \leq j \leq L-n$  in the amino acid sequence (length of  $L$ ),  
 (8) selecting at least one amino-acid residue to be subjected to mutation on the basis of the alignment data carried out in the step (1) from the amino acid sequence (length of  $L$ ) of the protein "A", sequentially repeating the steps (3) and (6) for variant amino-acid residues in various mutated amino acid sequences where the selected amino-acid residue has been mutated into another amino-acid residues, to determine the  $Z_j$  value of the variant amino-acid residues,  
 (9) selecting the mutation position and the mutated amino-acid residue wherein the  $Z_j$  value of the variant amino-acid residue as determined in the step (8) is larger or smaller than the  $Z_j$  value of the wild type amino-acid residue as determined in the step (7), and  
 (10) preparing a modified gene encoding the modified amino acid sequence having the mutated amino-acid residue at the position from the protein "B" gene, and producing the modified protein as the expression product of the gene.

**13.** A method for modifying the known function of protein "A" derived from the entire proteins of organism "a" of which genome data or cDNA data has been known, which method comprises the steps of:

- (1) extracting proteins closely related to the protein "A" from an existing protein data base and subjecting the proteins to alignment,  
 (2) determining the frequency of occurrence of each amino acid and the frequency of occurrence of individual oligopeptides produced by permutations of twenty amino acids, in the amino acid sequences of the entire proteins of the organism "a",  
 (3) as to an appropriate protein of the organism "a",  
 (3') determining in the entire proteins of the organism "a", the frequency of occurrence of the following  $A_{ji}$ -oligopeptide of given length of ( $n$ ) ( $1 \leq n \leq M$ , provided that  $M$  is the smallest length of oligopeptides satisfying the criterion that all the oligopeptides of length  $M$  are at frequency 1 of the occurrence), which  $A_{ji}$ -oligopeptide is a part of the amino acid sequence of the protein

and contains the  $j$ -th amino-acid residue  $A_j$ , on condition that the  $j$ -th amino-acid residue  $A_j$  is defined as the  $i$ -th residue  $A_{ji}$  from the N-terminus of the  $A_{ji}$ -oligopeptide ( $n \leq j \leq L-n+1$ );

$A_{ji}$ -oligopeptide:  $a_{j1}a_{j2} \dots a_{ji} \dots a_{jn}$  ( $1 \leq i \leq n$ ;  $A_j=a_{ji}$ , and  $A_j$  is the  $i$ -th residue of the oligopeptide; and moreover,  $a_{j1}=A_{j-i+1}, \dots, a_{jn}=A_{j-i+n}$ ),

and determining in the entire proteins of the organism "a", the frequency of occurrence of the following  $X_{ji}$ -oligopeptide of length ( $n$ ) corresponding to the length of  $A_{ji}$ -oligopeptide;

$X_{ji}$ -oligopeptide:  $a_{j1}a_{j2} \dots X_{ji} \dots a_{jn}$  ( $1 \leq i \leq n$ ; and the  $i$ -th residue  $X_{ji}$  is any amino acid; and moreover,  $a_{j1}=A_{j-i+1}, \dots, a_{jn}=A_{j-i+n}$ ),

- (4) calculating ratio  $Y_{ji}$  of the frequency of occurrence of the  $A_{ji}$ -oligopeptide to that of the  $X_{ji}$ -oligopeptide,  
 (5) determining mean  $Y(j,n)$  of the  $Y_{ji}$ ;

$$Y(j,n) = \sum_{i=1}^n Y_{ji}/n,$$

- (6) determining functional value  $Z(j,n)$  of  $Y(j,n)$ ;

$Z(j,n)=-\log(Y(j,n))$ ,

- (7) repeating the steps (3') to (6) sequentially and determining the  $Z(j,n)$  value of each amino-acid residue  $A_j$  at position  $j$  ( $n \leq j \leq L-n+1$ ) in the amino acid sequence (length of  $L$ ),  
 (8) sequentially repeating the steps (3) to (7) for the entire proteins of the organism "a", thereby determining the distribution of the  $Z(j,n)$  value of each amino-acid residue in the entire proteins, and the  $Z(j,n)$  values are classified into twenty according to the twenty amino acids, and then determining mean  $Av(Aa)$  of the  $Z(j,n)$  values for each amino acid  $Aa$  and the standard deviation  $Sd(Aa)$  of the distribution thereof, on the basis of the distribution, to determine function  $g$  to the  $j$ -th amino-acid residue  $A_j$  of a protein for normalizing the difference in distribution due to the species of amino-acid residues;  
 $g=g(Z(j,n), A_j)=[Z(j,n)-Av(Aa)]/Sd(Aa)$  (provided that  $A_j=Aa$ ; and  $1 \leq n \leq M$ ),  
 (9) determining value  $D(j,n)$  of the function  $g$  of each  $A_j$  of all the amino-acid residues at position  $j$  ( $n \leq j \leq L-n+1$  and  $1 \leq n \leq M$ ) of a protein in the entire proteins as recovered in the step (8);  
 $D(j,n)=g(Z(j,n), A_j)$ , and  
 (10) defining the representative value of the function of the  $j$ -th amino-acid residue in the amino acid sequence (length of  $L$ ) as functional value  $W_j$  of the  $Z(j,n)$  and  $D(j,n)$ ;  
 $W_j=h(Z(j,1), Z(j,2), \dots, Z(j,M), D(j,1), D(j,2), \dots, D(j,M))$   
 (11) sequentially repeating the steps (3) to (10), to determine the individual  $W_j$  values of all the amino-acid residues at position  $n \leq j \leq L-n+1$  in the amino acid sequence (length of  $L$ ),

- (12) selecting at least one amino-acid residue to be subjected to mutation on the basis of the alignment data carried out in the step (1) from the amino acid sequence (length of L) of the protein "A", and sequentially repeating the steps (3) and (10) for variant amino-acid residues in various mutated amino acid sequences where the selected amino-acid residue has been mutated into another amino-acid residue, to determine the  $W_j$  value of the variant amino-acid residue,
- (13) selecting a mutated amino acid sequence wherein the  $W_j$  value of the variant amino-acid residue as determined in the step (12) is larger or smaller than the  $W_j$  value of the wild type amino-acid residue as determined in the step (10), and
- (14) preparing a modified gene encoding the modified amino acid sequence from the protein "A" gene, and producing the modified protein as the expression product of the gene.
- 14.** A protein of which function is artificially modified by the method according to claim **11**, **12** or **13**.
- 15.** The protein according to claim 14, which is a thermophilic DNA polymerase prepared by artificially modifying the amino acid sequence of Pfu DNA polymerase so that the elongation of synthesized DNA chain might not be terminated intermediately during the catalysis for the synthesis of a DNA chain complementary to a single-stranded DNA.
- 16.** The thermophilic DNA polymerase according to claim 14, comprising the amino acid sequence of SQ ID No.1.
- 17.** The protein according to claim 14, which is a thermophilic DNA polymerase prepared by artificially modifying the amino acid sequence of Pfu DNA polymerase so that the synthesized DNA chain might be more elongated during the catalysis for the synthesis of a DNA chain complementary to a single-stranded DNA.
- 18.** The DNA polymerase according to claim 17, comprising the amino acid sequence of SQ ID No.6.
- 19.** The DNA polymerase according to claim 17, comprising the amino acid sequence of SQ ID. No.7.
- 20.** A DNA sequence encoding the amino acid sequence of SQ ID No.1.
- 21.** A DNA sequence encoding the amino acid sequence of SQ ID No. 6.
- 22.** A DNA sequence encoding the amino acid sequence of SQ ID No.7.
- 23.** A recombinant vector carrying the DNA sequence according to claim 20.
- 24.** The recombinant vector according to claim 24, which is a recombinant plasmid pDP320 carried on *Escherichia coli* HMS 174(DE3)/pDP 320 (FERM P-16052).
- 25.** A recombinant vector carrying the DNA sequence according to claim 21.
- 26.** The recombinant vector according to claim 25, which is a recombinant plasmid pDP 5b17 carried on *Escherichia coli* HMS 174(DE3)/pDP 5b17 (FERM BP-6189).
- 27.** A recombinant vector carrying the DNA sequence according to claim 22.
- 28.** The recombinant vector according to claim 27, which is a recombinant plasmid pDP 5c4 carried on *Escherichia coli* HMS 174(DE3)/pDP 5c4(FERM BP-6190).
- 29.** A method for preparing a thermophilic DNA polymerase, comprising culturing a cell transformed with an expression vector carrying the DNA sequence according to claim 20 and isolating and purifying the objective enzyme produced in the culture medium.
- 30.** A method for preparing a DNA polymerase, comprising culturing a cell transformed with an expression vector carrying the DNA sequence according to claim 21 or **22** and isolating and purifying the objective enzyme produced in the culture medium.

\* \* \* \* \*