



(11)

EP 2 505 660 B1

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention of the grant of the patent:

21.05.2014 Bulletin 2014/21

(21) Application number: **10833235.4**

(22) Date of filing: **25.11.2010**

(51) Int Cl.:

C12Q 1/48 ^(2006.01)	A61K 39/395 ^(2006.01)
A61K 45/00 ^(2006.01)	A61P 3/06 ^(2006.01)
A61P 9/10 ^(2006.01)	A61P 43/00 ^(2006.01)
C12N 15/09 ^(2006.01)	C12Q 1/02 ^(2006.01)
G01N 33/15 ^(2006.01)	G01N 33/50 ^(2006.01)
G01N 33/92 ^(2006.01)	C12N 5/10 ^(2006.01)
C12Q 1/60 ^(2006.01)	C12Q 1/61 ^(2006.01)

(86) International application number:

PCT/JP2010/070959

(87) International publication number:

WO 2011/065389 (03.06.2011 Gazette 2011/22)

(54) **METHOD FOR SCREENING OF THERAPEUTIC AGENT FOR HYPERLIPEMIA**

SCREENING-VERFAHREN FÜR EIN MITTEL ZUR BEHANDLUNG VON HYPERLIPÄMIE

MÉTHODE DE CRIBLAGE D'UN AGENT DE TRAITEMENT DE L'HYPERLIPÉMIE

(84) Designated Contracting States:

AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR

(30) Priority: **27.11.2009 JP 2009269593**

(43) Date of publication of application:

03.10.2012 Bulletin 2012/40

(73) Proprietor: **Japan Science and Technology Agency**

**Kawaguchi-shi
Saitama 332-0012 (JP)**

(72) Inventors:

- **INOKUCHI Jinichi**
Sendai-shi
Miyagi 981-3203 (JP)
- **NAGAFUKU Masakazu**
Sendai-shi, Miyagi
989-3124 Japan (JP)
- **HAYAMIZU Hirota**
Minamiuonuma-shi, Niigata
949-7312 Japan (JP)

(74) Representative: **Cabinet Plasseraud**

**52, rue de la Victoire
75440 Paris Cedex 09 (FR)**

(56) References cited:

EP-A1- 1 726 313 WO-A2-02/084301
JP-A- 2003 238 410

- **JIN'ICHI INOKUCHI ET AL.:** 'Insulin resistance in type 2 diabetes as a microdomain syndrome, Ganglioside GM3 no Kan'yo o Chushin to shite' **PROTEIN, NUCLEIC ACID AND ENZYME** vol. 48, no. 8, 2003, pages 1179 - 1183
- **GARNER, B. ET AL.:** 'Increased glycosphingolipid levels in serum and aortae of apolipoprotein E gene knockout mice' **J. LIPID RES.** vol. 43, 2002, pages 205 - 214
- **YAMASHITA, T. ET AL.:** 'Enhanced insulin sensitivity in mice lacking ganglioside GM3' **PROC. NATL. ACAD. SCI. U.S.A.** vol. 100, no. 6, 2003, pages 3445 - 3449
- **JIN'ICHI INOKUCHI ET AL.:** 'Microdomain Kino Ijo ni Motozuku 2-Gata Tonyobyono no Byotai Kaimei' **CORE RESEARCH FOR EVOLUTIONAL SCIENCE AND TECHNOLOGY CREST RESEARCH AREAS** vol. 64, no. 1, 2009, pages 65 - 105

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 505 660 B1

Description

TECHNICAL FIELD

5 [0001] The present invention provides a method for screening a novel therapeutic agent for hyperlipidemia. More specifically, the present invention provides a method for screening a substance that provides a blood lipid level lowering activity through inhibiting the production or function of gangliosides, particularly GM3, or inhibiting the activity or expression of GM3 synthase. The present disclosure also provides a pharmaceutical composition effective for treating hyperlipidemia that specifically inhibits the synthesis of gangliosides, particularly GM3, and others.

BACKGROUND ART

15 [0002] Circulatory disorders associated with arteriosclerosis including myocardial infarction, cerebral infarction, etc. are increasing yearly and are one of the leading causes of death in adults. There are various causes to induce arteriosclerosis and hyperlipidemia (including hypercholesteremia, hypertriglyceridemia, etc.) is recognized to be one of the most important causes. For the purpose of treating hypercholesteremia, drugs such as HMG-CoA reductase inhibitors (especially, statin type medicaments), anionic exchange resin drugs, etc. have been used. However, these drugs not only inhibit cholesterol biosynthesis but also inhibit the biosynthesis of some other components such as ubiquinone, dolichol and heme A, which are essential for the living body. It is thus concerned that adverse effects caused thereby might take place.

20 [0003] In recent years, the importance of glycosphingolipids (GSLs) in various metabolic disorders (including insulin resistance, type 2 diabetes mellitus, hyperlipidemia, arteriosclerosis, fatty liver, etc.) caused by obesity has been revealed. GSL falls in a molecular group formed by the addition of various sugar chains to the backbone of ceramide and is present in blood or on cell membranes of all cells. GSL is biosynthesized *in vivo* through a series of enzyme reactions starting from ceramide (FIG 1). GSL containing sialic acid is called a ganglioside family and starts from GM3 which is synthesized from lactocylceramide (LacCer) by GM3 synthase (sialic acid transferase I: SAT-I) (FIG 1). Inhibitors to the enzymes involved in biosynthesis of glucosylceramide (GlcCer), which is the early stage of GSL biosynthesis pathway, include: D-threo-1-phenyl-2-decanoylamino-3-morpholino-1-propanol (D-threo-PDMP) (Non-Patent Document 1), (1R,2R)-nonanoic acid [-2-(2,3-dihydro-benzo[1,4] dioxin-6-yl)-2-hydroxy-1-pyrrolidin-1-ylmethyl-ethyl] -amide-1-tartaric acid salt (Genz-123346), which is a D-threo-PDMP analog, N-(5-adamantane-1-yl-methoxy)-pentyl-1-deoxyojirimycin (AMP-DNM), etc. These inhibitors are reported improving insulin resistance *in vitro* and in obese animal models, having an effect of improving fatty liver and even promoting excretion of free fatty acids (and reducing cholesterol to promote excretion of cholesterol from the liver into the bile (i.e., activation of the reverse cholesterol transport system), etc. (Non-Patent Documents 2 to 7). Thus, development toward clinical applications is ongoing.

35 [0004] However, it is reported that glucosylceramide synthase knockout mice caused embryonic lethality (Non-Patent Document 8) and hence, there is also a concern of potential adverse effects of inhibitors of this enzyme. On the other hand, the life span of GM3 synthase knockout mice is equivalent to that of wild-type mice (Non-Patent Document 9). It is therefore expected that adverse effects caused by inhibition of GM3 synthase may be minimized.

40 [0005] GSLs found in insulin-responsible organs (such as muscle, liver and adipose tissue) are highly diverse and include glucosylceramides (GlcCer), lactosylceramides (LacCer) and gangliosides GM3 and GM2. Yamashita et al. established GM3 synthase (SAT-I) knockout mice and reported that gangliosides of the a- and b-series, which are normally expressed as shown in FIG 1, were not expressed in the mice, and when the onset of high-fat diet-induced insulin resistance was compared with the onset in the wild-type mice, insulin resistance was reduced in the SAT-I knockout mice (Non-Patent Document 9).

45 [Non-Patent Document 1] Inokuchi J. & Radin N. (1987) J. Lipid Res. 28, 565-571

[Non-Patent Document 2] Tagami S., Inokuchi J., Kabayama K., Yoshimura H., Kitamura F., Uemura S., Ogawa C., Ishii A., Saito M., Ohtsuka Y, et al. (2002) J. Biol. Chem. 277, 3085-3092

50 [Non-Patent Document 3] Zhao H., Przybylska M., Wu I.H., Zhang J., Siegel C., Komamitsky S., Yew N.S., & Cheng S.H. (2007) Diabetes 56, 1210-1218

[Non-Patent Document 4] Zhao H., Przybylska M., Wu I.H., Zhang J., Maniatis P., Pacheco J., Piepenhagen P., Copeland D., Arbeeny C., Shayman J.A., et al. (2009) Hepatology 50, 85-93

[Non-Patent Document 5] Aerts J. M., Ottenhoff R., Powlson A.S., Grefhorst A., van Eijk M., Dubbelhuis P.F., Aten J., Kuipers F., Serlie M. J., Wennekes T., et al. (2007) Diabetes 56, 1341-1349

55 [Non-Patent Document 6] van Eijk M., Aten J., Bijl N., Ottenhoff R., van Roomen C.P., Dubbelhuis P.F., Seeman I., Ghauharali-van der Vlugt K., Overkleeft H.S., Arbeeny C., et al. (2009) PLoS One 4, e4723. Epub. 2009 Mar. 4723

[Non-Patent Document 7] Bijl N., van Roomen C.P., Triantis V, Sokolovic M., Ottenhoff R., Scheij S., van Eijk M., Boot R.G, Aerts J.M., & Groen A.K. (2009) Hepatology 49, 637-645

[Non-Patent Document 8] YAMASHITA, T., et al, Proc. Natl. Acad. Sci. USA, Vol. 96, pp. 9142-9147, 1999

[Non-Patent Document 9] Yamashita T., Hashiramoto A., Haluzik M., Mizukami H., Beck S., Norton A., Kono M., Tsuji S., Daniotti J.L., Werth N., et al. (2003) Proc. Natl. Acad. Sci. U S A 100, 3445-3449

5 [0006] EP 1726313 relates to a prophylactic/therapeutic agent for atherosclerosis comprising GM3 synthase inhibitor or inhibitor for expression of GM3 synthase gene.

[0007] WO 02/084301 discloses a method for screening substances capable of modulating lipid metabolism by determining their ability to modulate the activity of proteins interacting with, and modulating the activity of, the protein expressed by the ABCA1.

10

DISCLOSURE OF THE INVENTION

Technical Problem

15 [0008] Therefore, the present inventors have examined lipid metabolism abnormality, as a target disease for treatment based on GM3 synthase regulation, in order to provide a means which provides a novel and highly safe drug for treating hyperlipidemia and a method for preventing/treating hyperlipidemia

Solution of Problem

20

[0009] The present inventors made extensive efforts to solve the problem described above, prepared the double knockout mice (SAT-I/apoE KO mice) where the gene involved in ganglioside biosynthesis (GM3 synthase gene) and the gene for apolipoprotein E (apoE) were knocked out, and analyzed, on an individual level, the functional role of gangliosides in various obesity-caused metabolic disorders. As a result, it has been found that a blood lipid level (including a blood cholesterol level and/or a blood triglyceride level) could be reduced through inhibiting GM3 synthase. The present invention has thus been accomplished.

25

[0010] Accordingly, the present invention provides a method for screening a substance that reduces a blood lipid level (including a blood cholesterol level and/or a blood triglyceride level), which is effective for treating hyperlipidemia as defined on the claims. A pharmaceutical composition effective for treating hyperlipidemia which selectively inhibits synthesis of gangliosides (particularly GM3), and others are also disclosed.

30

[0011] Specifically, the present invention describes the screening method, below.

1. A method for screening a GM3 synthase specific inhibitor having a blood lipid level lowering activity, which comprises:

35

- (i) contacting GM3 synthase with a substrate thereof in the presence and absence of a test substance,
- (ii) selecting a test substance that decreases the synthesized amount of GM3,
- (iii) administering the selected test substance to a non-human animal, and
- (iv) determining a cholesterol level and/or a triglyceride level in blood in the non-human animal, and selecting a test substance which reduces the cholesterol level and/or the triglyceride level by 10% or more as compared to a control.

40

2. A method for screening a substance that inhibits GM3 synthase expression and has a blood lipid level lowering activity, which comprises:

45

- (i) contacting a test substance with a cell expressing GM3 synthase,
- (ii) selecting a test substance that decreases the expression level of GM3 synthase as compared to a control,
- (iii) administering the selected test substance to a non-human animal, and
- (iv) determining a cholesterol level and/or a triglyceride level in blood in the non-human animal, and selecting a test substance which reduces the cholesterol level and/or the triglyceride level by 10% or more as compared to a control.

50

3. The method according to 2, wherein the cell expressing GM3 synthase is selected from the group consisting of adipocytes, hepatocytes, vascular endothelial cells, vascular smooth muscle cells, skeletal muscle cells, epithelial cells, neurocytes, fibroblasts, monocytes and macrophages.

55

4. The method according to 2, wherein the cell expressing GM3 synthase is a transformant produced by genetic engineering.

5. The screening method according to any one of 1 to 4, which further comprises selecting a test substance that reduces a blood cholesterol level and/or a blood triglyceride level.

5 6. The method according to any one of 1 to 5, wherein the non-human animal has a higher blood lipid level than normal level.

7. The method according to any one of 1 to 6, wherein the non-human animal is a mouse.

10 **[0012]** Also disclosed, but not part of the invention are :

[8] A pharmaceutical composition effective for treatment of hyperlipidemia, comprising a substance that specifically prevents a production of a ganglioside.

15 [9] A pharmaceutical composition effective for treatment of hyperlipidemia, comprising a substance that selectively suppresses a function of a ganglioside.

[10] The composition according to [9], wherein the ganglioside is GM3.

20 [11] A pharmaceutical composition effective for treatment of hyperlipidemia, comprising an antibody against GM3, a GM3 synthase inhibitor or a substance that inhibits GM3 synthase expression.

[12] A method for treating hyperlipidemia which comprises administering an effective dose of a pharmaceutical composition effective for treatment of hyperlipidemia to a non-human animal to reduce a blood lipid level.

25 [13] The method according to [12], wherein the pharmaceutical composition reduces a blood cholesterol level and/or a blood triglyceride level.

[14] The method according to [12] or [13], wherein the pharmaceutical composition comprises a substance that selectively inhibits production of a ganglioside.

30 [15] The method according to [14], wherein the ganglioside is GM3.

[16] The method according to [15], wherein the substance that selectively inhibits the ganglioside production is an antibody against GM3, a GM3 synthase inhibitor or a substance that inhibits GM3 synthase expression.

35 [17] A method for regulating a blood lipid level with an antibody against GM3 or a GM3 synthase inhibitor, or by inhibiting GM3 synthase expression.

40 **Advantageous Effect of Invention**

[0013] The pharmaceutical composition inhibits ganglioside GM3 biosynthesis to reduce a blood lipid level (including a blood cholesterol level and/or a blood triglyceride level), and can provide a new method for treating hyperlipidemia even from a clinical viewpoint.

45 **BRIEF DESCRIPTION OF DRAWINGS**

[0014]

FIG 1 shows the ganglioside biosynthesis pathway.

50 FIG 2 shows the method for preparing GM3 synthase gene knockout mice (SAT-I KO).

FIG 3 shows the preparation of SAT-I/apoE double knockout (DKO) mice and confirmation of the genes.

FIG 4 shows the assay results of the body weight and blood lipid level of SAT-I/apoE double knockout mice in FIG 3. FIG 5 shows the profiles of plasma lipoproteins from ApoE knockout mice and SAT-I/ApoE double knockout mice, which were fractionated with gel filtration high performance chromatography.

55 FIG 6 shows the change in the lipid composition in the livers of ApoE knockout mice and SAT-I/ApoE double knockout mice.

FIG 7 shows the analysis results of plasma glycosphingolipids from SAT-I knockout mice, ApoE knockout mice and SAT-I/ApoE double knockout mice.

FIG 8 shows the analysis results of glycosphingolipids in the livers of SAT-I knockout mice, ApoE knockout mice and SAT-I/ApoE double knockout mice.

FIG 9 shows the assay results of the ganglioside composition in the hearts, livers and muscles of SAT-I knockout mice and wild-type mice.

FIG 10 shows the assay results of plasma lipids in SAT-I knockout mice fed with high cholesterol diet.

EMBODIMENTS FOR CARRYING OUT THE INVENTION

1. Screening Method of the Invention

[0015] The present invention provides a method for screening a GM3 synthase specific inhibitor having a blood lipid level lowering activity, namely, an activity of reducing a blood cholesterol level and/or blood triglyceride level, which comprises inter alia (i) administering a test substance to a non-human animal, and (ii) determining the blood lipid level in the non-human animal. Other steps have been defined in the claims. Examples of the test substance include a peptide, a protein, a nonpeptidic compound, a synthetic compound, a fermentation product, a cell extraction liquid, a plant extraction liquid, an animal tissue extraction liquid, plasma, etc. The test substance administered may be used solely or in multiple combination manner. In determining the blood lipid level, there may be used known methods as in the aforesaid assay methods, etc.

[0016] As used herein, the term "substance or GM3 synthase specific inhibitor having a blood lipid level lowering activity" is intended to mean a substance that reduces a cholesterol level and/or a triglyceride level in blood. Specifically, the "substance or GM3 synthase specific inhibitor having a blood lipid level lowering activity", as used herein, refers to a substance that has the cholesterol level and/or the triglyceride level lowering activities in a sample (plasma) obtained from the subject, who received the substance, by 10% or more, preferably 20% or more, more preferably 40% or more, much more preferably 60% or more, and most preferably 80% or more, as compared to the control, when assayed according to the method described in KISO SEIKAGAKU JIKKENHO (Basic Biochemical Experiments), Vol. 5, Shishitsu, Toshitsu, Fukugotoshitsu (Lipids, Carbohydrates and Glycoconjugates) (edited by The Japanese Biochemical Society). In this case, the reduction is limited to nearly normal blood lipid level.

[0017] The blood lipid level in a sample from the subject can be determined through known methods utilizing the enzyme method, the immunodiffusion method or the like. Such methods include, for example, the methods described in KISO SEIKAGAKU JIKKENHO (Basic Biochemical Experiments), Vol. 5, Shishitsu, Toshitsu, Fukugotoshitsu (Lipids, Carbohydrates and Glycoconjugates) (edited by The Japanese Biochemical Society) described above. In addition, assay kits for triglycerides and cholesterol are also commercially available from various manufacturers. Examples are Cholesterol E-Test Wako (Wako Pure Chemical Industries, Ltd) for the cholesterol assay kit, TG-EN Kainos (Kainos Laboratories) for the triglyceride assay kit, and NEFA C-Test Wako (Wako Pure Chemical Industries, Ltd) for the free fatty acid assay kit.

[0018] The screening method of the present invention may involve any of the methods developed to date as the methods for assaying cholesterol and triglycerides. In more detail, the screening method of the present invention may further include the step with high performance liquid chromatography (HPLC) or gas chromatography. Regarding detailed principles or the like, see, e.g., KISO SEIKAGAKU JIKKENHO (Basic Biochemical Experiments), Vol. 5, Shishitsu, Toshitsu, Fukugotoshitsu (Lipids, Carbohydrates and Glycoconjugates) (edited by The Japanese Biochemical Society).

[0019] As used herein, the term "reduces a blood cholesterol level and/or a blood triglyceride level" is intended to mean to reduce the cholesterol level and/or the triglyceride level in blood by 10% or more, preferably 20% or more, more preferably 40% or more, much more preferably 60% or more, and most preferably 80% or more, in terms of the concentration of at least one of the cholesterol and triglyceride, as compared to the control, when assayed through the conventional method described above. However, the reduction is limited to nearly normal blood lipid level.

[0020] The substance that reduces a blood lipid level (including a blood cholesterol level and/or a blood triglyceride level) is effective for treating or preventing hyperlipidemia. As used herein, the term "hyperlipidemia" is also called hyperlipoproteinemia and refers to such a condition that a plasma lipid other than free fatty acids is elevated. Healthy adults have, in plasma, 150 to 220 mg/dl of cholesterols, 50 to 140 mg/dl of triacylglycerols and 150 to 220 mg/dl of phospholipids, and they are bound to apolipoproteins and dispersed as lipoproteins. Hyperlipidemia may be classified into hyperlipidemia where triacylglycerols are mainly increased (e.g., type I hyperlipidemia, type IV hyperlipidemia and type V hyperlipidemia), and hyperlipidemia where cholesterol is mainly increased (e.g., familial hypercholesteremia, familial complicated hyperlipidemia and type III hyperlipidemia) (Mariko Shiba, et al., SHIN SEIKAGAKU JIKKEN KOZA (New Series of Biochemical Experiments), Vol. 4, "Shishitsu I: Chusei Shishitsu-to-III Gata Koshiketsusho (Lipid I: Neutral Lipids and Lipoproteins)," published by Tokyo Kagaku Dojin Publishing Co., 1993, Chapter 11).

[0021] The substance that reduces the blood lipid level is not only effective for treating or preventing hyperlipidemia but also effective for treating or preventing at least one disease selected from arteriosclerosis, atherosclerosis, peripheral vascular disease, hyper LDLemia, hypo HDLemia, hypercholesteremia, hypertriglyceridemia, familial hypercholesteremia, cardiovascular disturbance, angina pectoris, ischemia, cardiac ischemia, thrombosis, myocardial infarction, reper-

fusion damage, angioplasty restenosis and hypertension.

[0022] Examples of the non-human animal as the subject used in the screening method described above include animals which are generally utilized in tests of pharmaceutical products, such as a monkey, chimpanzee, dog, cat, guinea pig, rat, mouse, rabbit, pig, sheep, horse, etc., preferably, a monkey, chimpanzee, rabbit, rat and mouse, and more preferably, a mouse or rat.

[0023] In the present invention, the substance having the blood lipid level lowering activity may be a substance that can specifically inhibit the function of GM3, or may be substance that can selectively inhibit a specific function in various functions of GM3. The glycosphingolipid as used herein refers to a lipid comprising a saccharide, a fatty acid and a sphingosine which is a long-chain base in the molecule. The ganglioside as used herein collectively refers to the family of sphingoglycolipids containing sialic acid, and is a molecule in which a sugar chain containing sialic acid is covalently bound to a lipid called ceramide. Today, gangliosides with various sugar chain structures are known, and GM3 is the first ganglioside molecule in the biosynthesis pathway (see FIG 1). That is, all endogenous ganglio-type gangliosides are biosynthesized through a series of enzyme reactions in which ceramide is used as a starting material and a GM3 synthase triggers the first step. During the series of reactions, GM3 is a molecule that is the origin of all gangliosides, and is synthesized from lactosylceramide by GM3 synthase (SAT-I). The sugar chain moiety is sequentially synthesized by glycosyltransferase in the lumen of the Golgi body in the cell using a sugar nucleotide as a donor.

[0024] In a specific biosynthesis pathway of ganglio-type gangliosides, GM3 is produced from Gal-Glc-Cer by GM3 synthase (SAT-I), and then, from this GM3, various gangliosides including the a-series (GM3, GM2, GM1, GD1 a and GT1a), the b-series (GD3, GD2, GD1b, GT1b and GQ1b) and the c-series (GT3, GT2, GT1c, GQ1c and GP1c) are produced, as shown in FIG 1. It is understood from the figure that production of these ganglioside molecules is dependent on GM3 synthase. In view of this, GM3 has the functions as the starting material for other gangliosides.

[0025] It is also reported that GM3 plays the functions of various growth factor receptors (Inokuchi J. and Kabayama K. (2007) Receptor Modifications in Glycobiology. Comprehensive Glycoscience 3, 733-744. (Elsevier Science & Technology)) or the functions to regulate molecular transporters (Glaros EN., et al. Glycosphingolipid Accumulation Inhibits Cholesterol Efflux via the ABCA1/Apolipoprotein A-I Pathway J. Biol. Chem. 280, 24515-24523, 2005). Furthermore, with regard to GM3 functions, a new molecular mechanism has recently been proposed on unresponsiveness to insulin signaling where GM3 serves to dissociate insulin receptors from caveolae (Kabayama K., et al., Glycobiology (2005) 15, 21-29, Kabayama K., et al., Proc. Natl. Acad. Sci. U S A (2007) 104, 13678-13683).

[0026] In the present invention, the substance having the blood lipid level lowering activity may be an antibody against GM3, which specifically or selectively inhibits a function of GM3. Such an antibody includes a known monoclonal antibody specific to GM3 (Kotani, M., et al., Biochem. Biophys. Acta, 1117, 97-103 (1992)). In one specific embodiment, the antibody may inhibit a function of GM3 on cell membranes of cells including primary culture cells obtained from adipocytes, hepatocytes, vascular endothelial cells, vascular smooth muscle cells, epithelial cells, fibroblasts, monocytes, macrophages, etc., and various cell lines (e.g., mouse 3T3-L1 in adipocytes; human HepG2 in hepatocytes; in endothelial cells, immortalized endothelial cell lines having an acetylated LDL uptake activity; RAW264.7 in macrophages; etc., but not limited to these cell lines). In the context of the present application, it is understood that the antibody includes a monoclonal antibody, a polyclonal antibody, an anti-idiotypic antibody, an antibody fragment (e.g., Fab, F(ab')₂, Fv variable region or complementarity-determining region). It is understood that, when the antibody is bound with K_a of 10^{-7} M or more, preferably 10^{-8} M or more, the antibody is specific to an antigen, namely, GM3 or GM3 synthase. The affinity of the monoclonal antibody can be easily determined by those skilled in the art (see Scatchard. Ann. N.Y. Acad. Sci. 51: 660-672, 1949).

[0027] In the present invention, the substance having the blood lipid level lowering activity may be "a GM3 synthase inhibitor," namely, a substance that specifically inhibits the GM3 biosynthesis. As used herein, the GM3 synthase (SAT-I) refers to a protein comprising the amino acid sequence of SEQ ID NO: 2 or 4, or the amino acid sequence of SEQ ID NO: 2 or 4 having mutations of deletion, substitution, insertion and/or addition of one to several (1 to 9, to 8, 1 to 7, 1 to 6, 1 to 5, 1 to 4, 1 to 3, 1 to 2, or 1) amino acid(s) therein. In the biosynthesis of GM3, glucosylceramide synthase acts on ceramide (Cer) to synthesize glucosylceramide (GlcCer), and then lactosylceramide (LacCer) is produced by lactosylceramide synthase, as shown in FIG 1. The GM3 synthase recognizes this LacCer as a substrate to synthesize GM3. That is, examples include those which can reduce the amount of GM3 synthesized via competitive inhibition of enzyme reactions with substrate analogs of glucosylceramide synthase, lactosylceramide synthase and GM3 synthase, or non-competitively through binding to these enzymes.

[0028] In the present invention, the substance having the blood lipid level lowering activity may be "a substance that specifically inhibits GM3 synthase expression." As used herein, cDNA of GM3 synthase (SAT-I) is represented by, for example, SEQ ID NO: 1 or 3. Examples of the substance that specifically inhibits GM3 synthase expression include an antisense nucleic acid, a ribozyme and a dsRNA having RNAi effect, etc.

[0029] As used herein, the term "nucleic acid" is interchangeably used with "polynucleotide", "gene" or "nucleic acid molecule", and is intended to mean a polymer of nucleotide. As used herein, the term "base sequence" is interchangeably used with "nucleic acid sequence" or "nucleotide sequence", and is represented as a sequence of deoxyribonucleotide

(abbreviated as A, G, C and T). Furthermore, the "polynucleotide comprising the base sequence of SEQ ID NO: 1 or a fragment thereof" is intended to mean a polynucleotide comprising a sequence represented by the respective deoxy-nucleotides A, G, C and/or T of SEQ ID NO: 1 or a fragment thereof.

[0030] The nucleic acid used in accordance with the present invention may be present in the form of RNA (e.g., mRNA) or DNA (e.g., cDNA or genomic DNA). DNA may be a double strand or a single strand. A single-stranded DNA or RNA may be a coding strand (also known as "sense strand") or a noncoding strand (also known as "antisense strand").

[0031] As used herein, the "substance that inhibits the expression of GM3 synthase" includes suppression of transcription of the GM3 synthase gene and suppression of translation into a protein. Moreover, it also includes complete stop of DNA expression as well as reduction of DNA expression.

[0032] One embodiment of the "substance that inhibits GM3 synthase expression" is a nucleic acid encoding the antisense chain complementary to the GM3 synthase gene. The antisense technology is publicly known as a method for specifically suppressing the expression of a particular endogenous gene, and is described in various documents (for example, see Hirashima and Inoue, "Shin Seikagaku Jikken Koza 2, Kakusan IV, Idenshi No Fukusei To Hatsugen (New Biochemistry Experimentation Lectures 2, Nucleic Acids IV, Replication and Expression of Gene)", edited by Japanese Biochemical Society, Tokyo Kagaku Dojin Co., Ltd., pp. 319-347, 1993; etc.). The antisense nucleic acid can be prepared, for example, based on information of the sequence of cDNA represented by SEQ ID NO: 1 or 3 according to the phosphorothionate method (Stein, Nucleic Acids Res., 16: 3209-3221, 1988) or others. The nucleic acid prepared can be utilized in the form directly applied to a cell according to a publicly-known method. Also, the nucleic acid can transform a desired cell in the form that the nucleic acid is incorporated into a vector having a known expression system and appropriately expressed. Preferably, the sequence of the antisense nucleic acid is complementary to the transcription product of an endogenous gene possessed by a cell to be transformed, while the sequence does not have to be completely complementary thereto as long as gene expression can be effectively inhibited. Preferably, the transcribed RNA has at least 90% (e.g., at least 95%, at least 96%, at least 97%, at least 98%, and at least 99%) complementarity to the transcription product of a target gene. For effectively inhibiting expression of the target gene using the antisense sequence, the length of the antisense nucleic acid is at least 15 bases, preferably at least 100 bases, and more preferably at least 500 bases. Usually, the length of the antisense nucleic acid to be used is less than 5kb, and preferably less than 2.5 kb.

[0033] In the present invention, the expression of an endogenous GM3 synthase gene can also be suppressed by utilizing DNA encoding a ribozyme. The ribozyme is intended to mean an RNA molecule having a catalytic activity, and cleaves the transcription product of a target DNA to inhibit the function of the gene. Regarding the design of ribozymes, reference can be made to various known documents (see, e.g., FEBS Lett. 228: 228, 1988; FEBS Lett. 239: 285, 1988; Nucl. Acids Res. 17: 7059, 1989; Nature 323: 349, 1986; Nucl. Acids Res. 19: 6751, 1991; Protein Eng 3: 733, 1990; Nucl. Acids Res. 19: 3875, 1991; Nucl. Acids Res. 19: 5125, 1991; Biochem. Biophys. Res. Commun. 186: 1271, 1992, etc.). Furthermore, the "polynucleotide encoding an RNA which suppresses DNA expression through a co-suppression effect" refers to a nucleotide which inhibits the function of a target DNA by 'co-suppression.'"

[0034] Moreover, the suppression of an endogenous gene expression in the present invention can also be achieved by transforming a desired cell with a gene having a dominant negative character of a target gene. The gene having a dominant negative character is intended to mean a gene having the function to extinguish or reduce the activity of an endogenous wild-type gene inherent to a desired cell, by expressing the gene.

[0035] Another embodiment of the nucleic acid to be used for suppressing the expression of GM3 synthase gene is RNA that suppresses DNA expression by RNAi effect. The term "RNAi" is intended to mean a phenomenon in which, when a double-stranded RNA having a sequence that is identical with or similar to a target gene sequence is introduced into a cell, expression of both the exogenous gene introduced and the target endogenous gene is suppressed. As used herein, RNA includes, for example, double-stranded RNA having 21 to 25 nucleotides causing RNA interference, such as dsRNA (double strand RNA), siRNA (small interfering RNA) and shRNA (short hairpin RNA). Such RNA may also be locally delivered to a desired site by the delivery system of liposomes or the like, and may be locally expressed using a vector in which the above-described double-strand RNA can be produced. Methods for preparing and using such a double-stranded RNA (dsRNA, siRNA or shRNA) are publicly known from many documents (see, e.g., Japanese Laid-Open Patent Application Publication (Translation of PCT Application) No. 2002-516062; USPA No. 2002/086356A; Nature Genetics, 24(2), 180-183, 2000 Feb.; Genesis, 26(4), 240-244, 2000 April; Nature, 407:6802, 319-20, 2002 Sep. 21; Genes & Dev., Vol. 16(8), 948-958, 2002 Apr. 15; Proc. Natl. Acad. Sci. USA., 99(8), 5515-5520, 2002 Apr. 16; Science, 296(5567), 550-553, 2002 Apr. 19; Proc Natl. Acad. Sci. USA, 99:9, 6047-6052, 2002 Apr. 30; Nature Biotechnology, Vol. 20 (5), 497-500, 2002 May; Nature Biotechnology, Vol. 20(5), 500-505, 2002 May; Nucleic Acids Res., 30:10, e46, 2002 May 15, etc.).

[0036] In a first aspect, the screening method of the present invention includes performing primary screening for a test substance *in vitro* before the test substance is administered to a non-human animal. In more detail, the method includes the steps of contacting glycosphingolipid synthase with a substrate thereof in the presence and absence of a test substance and selecting a test substance which reduces the amount of the product synthesized. The combination of glycosphingolipid synthase and a glycosphingolipid is preferably the combination of ganglioside synthase and a

ganglioside, and more preferably, the combination of GM3 synthase and GM3. Combinations of other ganglioside synthases and gangliosides are shown in FIG 1.

[0037] The screening method includes a method using an enzyme, as GM3 synthase used in the screening method described above, which is expressed in primary culture cells obtained from adipocytes, hepatocytes, vascular endothelial cells, vascular smooth muscle cells, skeletal muscle cells, epithelial cells, neurocytes, fibroblasts, monocytes, macrophages, etc. as well as in various cell lines (including, but not limited to, mouse 3T3-L1 in adipocytes; human HepG2 in hepatocytes; in endothelial cells, immortalized endothelial cell lines having an acetylated LDL uptake activity; RAW264.7 in macrophages; etc.), and a method using an enzyme produced by genetic engineering techniques, chemical synthesis, or the like are included. A preferred example is the enzyme derived from the nucleotide sequence described in SEQ ID NO: 1 or 3. The prepared GM3 synthase in this case may be in an isolated state or may not be in an isolated state. For instance, the above cells bearing the GM3 synthase may be used, or the GM3 synthase may be purified from the above cells. Alternatively, the GM3 synthase may also be prepared, for example, by using various culture cells, e.g., *Escherichia coli*, yeast, insect cells, mammal cells, etc. as a host and introducing an appropriate expression vector into the host, by genetic engineering. For these known methods, reference can be made to the methods described in, e.g., *Molecular Cloning 3rd Ed.* (J. Sambrook, et al, Cold Spring Harbour Laboratory Press, 2001), *Current Protocols in Molecular Biology*, John Wiley & Sons 1987-1997, etc.

[0038] In a second aspect, the screening method of the present invention includes, before administration of a test substance to a non-human animal, the step of contacting the test substance *in vitro* with cells expressing a preferred level of glycosphingolipid synthase, e.g., primary culture cells obtained from adipocytes, hepatocytes, vascular endothelial cells, vascular smooth muscle cells, skeletal muscle cells, epithelial cells, neurocytes, fibroblasts, monocytes, macrophages, etc. and various cell lines (including, but not limited to, mouse 3T3-L1 in adipocytes; human HepG2 in hepatocytes; in endothelial cells, immortalized endothelial cell lines having an acetylated LDL uptake activity; RAW264.7 in macrophages; etc.) etc., or with transformants prepared by genetic engineering, and determining the amount of glycosphingolipid selectively produced in these cells. The preferred combination of glycosphingolipid synthase to be determined and a glycosphingolipid includes the combination of ganglioside synthase and a ganglioside, and more preferably, the combination of GM3 synthase and GM3. The preferred expression level herein is intended to mean a normal or greater expression level in the organ from which each cell is derived. The other combinations of ganglioside synthases and gangliosides can be found in FIG 1. For the genetic engineering techniques for producing the transformants described above, known methods described in, e.g., *Molecular Cloning 3rd Ed.* (J. Sambrook, et al, Cold Spring Harbour Laboratory Press, 2001), *Current Protocols in Molecular Biology*, John Wiley & Sons 1987-1997, etc. may be used.

[0039] For purification and quantitative detection of gangliosides, known methods can be used. For example, the method of Macher, B. A. and Klock, J. C. (*J. Biol. Chem.* 255, 2092-2096, 1980) or the method of Ledeen et al. (*J. Neurochem.* 21, 829-839, 1973) can be referred to. It is also possible to employ quantitative detection using an antibody which specifically acts on various gangliosides, e.g., GM3. In such a screening method, the reduction level in the GM3 production amount by a test substance is at least 10% or more, preferably 20% or more, more preferably 40% or more, even more preferably 60% or more, still more preferably 80% or more, still more preferably 90% or more and 100%, when compared with the control.

[0040] In the screening method described above, the combination may be selective combination of at least one ganglioside described in FIG 1, specifically at least one ganglioside selected from the group consisting of GM3, GM2, GM1, GD1a and GT1a (a-series) as well as GD3, GD2, GD1b, GT1b and GQ1b (b-series), and each ganglioside synthase. In such a screening method, the reduction level in the ganglioside production amount by a test substance is at least 10% or more, preferably 20% or more, more preferably 40% or more, even more preferably 60% or more, still more preferably 80% or more, 90% or more and 100%, when compared with the control.

[0041] The screening method of the present invention may include contacting a test substance with cells expressing a suitable level of glycosphingolipid synthase, e.g., primary culture cells obtained from adipocytes, hepatocytes, vascular endothelial cells, vascular smooth muscle cells, skeletal muscle cell, epithelial cells, neurocytes, fibroblasts, monocytes, macrophages, etc. and various cell lines (including, but not limited to, mouse 3T3-L1 in adipocytes; human HepG2 in hepatocytes; in endothelial cells, immortalized endothelial cell lines having an acetylated LDL uptake activity; RAW264.7 in macrophages; etc.) etc., and determining the selective expression level of GM3 synthase in these cells. The method for determining the expression level of GM3 synthase includes a method through hybridization utilizing a sequence of GM3 synthase, for example, northern blotting, a method utilizing a probe immobilized on a carrier, a method using a gene chip, quantitative PCR, etc. In such a screening method, for example, the suppression level of the expression for GM3 synthase by a test substance is 10% or more, preferably 20% or more, more preferably 40% or more, even more preferably 60% or more, and still more preferably 80% or more, 90% or more and 100%.

[0042] The test substance in this embodiment of the screening method of the present invention includes a polynucleotide which specifically inhibits the expression of GM3 synthase. Examples of the form of such a polynucleotide which inhibits the expression of GM3 synthase are the aforesaid antisense nucleic acid, ribozyme, double-strand RNA, etc. These polynucleotides which inhibit the expression of GM3 synthase can be prepared by a publicly-known method. Moreover,

these nucleic acids may be subjected to various chemical modifications.

[0043] The screening method of the present invention also includes a method which comprises using a transgenic non-human animal-derived cell or recombinant cell, in which a reporter gene is connected to the downstream of a promoter of GM3 synthase; contacting a test substance with the cell; and measuring the expression level of the reporter gene in the cell. As the recombinant cell to be used in this method, various cells may be used, and preferably, primary culture cells obtained from adipocytes, hepatocytes, vascular endothelial cells, vascular smooth muscle cells, skeletal muscle cells, epithelial cells, neurocytes, fibroblasts, monocytes, macrophages, etc., and various established cell lines (including, but not limited to, mouse 3T3-L1 in adipocytes; human HepG2 in hepatocytes; in endothelial cells, immortalized endothelial cell lines having an acetylated LDL uptake activity; RAW264.7 in macrophages; etc.), and the like. The gene sequence at the 5' upstream of GM3 synthase gene is described in Kim J-W., et al., *Gene* 273, 163-171, 2001, and the promoter is described in Kim S-W., et al., *Biochim. Biophys. Acta* 1578, 84-89, 2002; Choi H-J., et al., *Biochem. Biophys. Res. Commun.* 313, 142-147, 2004. Specifically, putative binding sites of transcription factors such as AP4, MZF1, SP1, ATF/CREB, NFY, IK2 and LYF1 have been found, and CREB has been additionally found which functions as a PMA-inducible promoter. Preferred examples of proteins encoded by the reporter gene to be used include various known proteins such as firefly luciferase, Renilla luciferase, green fluorescent protein (GFP), β -galactosidase, alkaline phosphatase, etc. These reporter proteins can be detected using a known detection method or kit. In such a screening method, for example, the suppression level of reporter protein expression upon stimulation by a test substance is 10% or more, preferably 20% or more, more preferably 40% or more, 60% or more, and even more preferably 80% or more, 90% or more and 100%.

2. Pharmaceutical Composition

[0044] It is described a pharmaceutical composition useful for treating and/or preventing hyperlipidemia, comprising the substance having the blood lipid level lowering activity.

[0045] The pharmaceutical composition has the blood lipid level lowering activity, and preferably, can be used for the treatment or prevention of hyperlipidemia. Alternatively, the pharmaceutical composition can be used for the treatment or prevention of diseases that can be treated by the blood lipid level lowering activity, for example, at least one disease selected from arteriosclerosis, atherosclerosis, peripheral vascular disease, hyper LDLemia, hypo HDLemia, hypercholesteremia, hypertriglyceridemia, familial hypercholesteremia, cardiovascular disturbance, angina pectoris, ischemia, cardiac ischemia, thrombosis, myocardial infarction, reperfusion damage, angioplasty restenosis and hypertension, and preferably, arteriosclerosis, hyper LDLemia, hypo HDLemia, hypercholesteremia, hypertriglyceridemia and familial hypercholesteremia. However, the present pharmaceutical composition reduces an abnormal blood lipid level in the subject down to nearly normal level.

[0046] The pharmaceutical composition may contain a substance that specifically inhibits the functions of ganglioside. Preferably, the ganglioside may be GM3, and the substance that specifically inhibits the functions of ganglioside may be an anti-GM3 antibody. The pharmaceutical composition may further contain a substance that selectively inhibits the functions of ganglioside. The substance includes, for example, a substance, e.g., an antibody, which can selectively inhibit, with GM3, a particular function among various functions, including the function of regulating the functions of various growth factor receptors on cell membranes, transport proteins, adhesive molecule receptors or functional proteins, raw material supply functions to various other gangliosides.

[0047] The pharmaceutical composition may contain a substance that selectively inhibits ganglioside production. Such a substance may include a ganglioside synthase inhibitor, an expression inhibitor of this enzyme, etc., which is obtained by the screening method of the present invention described above.

[0048] Where the pharmaceutical composition is used, the composition may be administered, for example, orally, intravenously, through the oral mucosa, rectally, vaginally, transdermally, intranasally, by inhalation, etc., and preferably, administered orally. The active component of the pharmaceutical composition may be formulated solely or in combination. It is also possible to blend a pharmaceutically acceptable carrier or formulation additive therein to provide a formulation. In this case, the active component can be contained in the formulation, for example, in an amount of 0.1 to 99.9 wt %.

[0049] The pharmaceutically acceptable carrier or additive, including, for example, an excipient, a disintegrant, a disintegrating aid, a binder, a lubricant, a coating agent, a dye, a diluent, a dissolving agent, a dissolving aid, a tonicity agent, a pH adjuster, a stabilizer, etc. can be used.

[0050] Examples of formulations suitable for oral administration include a powdered drug, a tablet, a capsule, a fine grain agent, a granular agent, a liquid drug, syrup, and the like. In the case of oral administration, various excipients such as microcrystalline cellulose, sodium citrate, calcium carbonate, dipotassium phosphate and glycine can be used in combination with: starch, preferably starch of corn, potato or tapioca; various disintegrants such as alginic acid and some of silicates; and a granulating binder such as polyvinyl pyrrolidone, sucrose, gelatin and gum arabic. Further, in many cases, lubricants such as magnesium stearate, sodium lauryl sulfate and talc are very effective for tablet formation. Such solid composition may be used with a gelatin capsule filled therein. In this regard, examples of preferred substances

include lactose or milk sugar as well as a high-molecular-weight polyethylene glycol. When an aqueous suspension and/or elixir is desired for oral administration, active components are used in combination with a sweetener or a flavor and a colorant or a dye, and optionally in combination with an emulsifier and/or a suspending agent, and it can be used together with water, ethanol, propylene glycol, glycerin, etc., a diluent containing a combination thereof, etc.

[0051] Examples of the formulation suitable for parenteral administration include an injectable agent and a suppository. In the case of parenteral administration, a solution in which the active ingredient is dissolved in sesame oil or peanut oil, or a solution in which the active component is dissolved in an aqueous solution of propylene glycol can be used. An aqueous solution optionally may be buffered suitably (preferably pH 8 or higher), and a liquid diluent is firstly required to become isotonic. Such an aqueous solution is suitable for intravenous injection, and an oily solution is suitable for intraarticular injection, intramuscular injection and subcutaneous injection. Preparation of all of these solutions under aseptic conditions can be easily accomplished by the standard formulation technology well-known in the art. Moreover, the active ingredient can be administered topically on the skin or the like. In this case, it is desirably administered topically in the form of cream, jelly, paste or ointment according to the standard pharmaceutical practice.

[0052] The dose of the pharmaceutical composition is not particularly limited, and an appropriate dose can be chosen depending on various conditions such as the type of disease, age and symptoms of the patient, administration route, therapeutic goal, presence or absence of concurrent drugs, etc. The dose of the pharmaceutical composition is, for example, 1 to 5,000 mg, and preferably 10 to 1,000 mg per day for an adult (e.g., body weight: 60 kg). The composition of such daily dosage may be administered daily in 2 to 4 divided doses.

3. Method for Treatment

[0053] In the method for treatment described herein, the substance having the blood lipid level lowering activity is the substance capable of reducing a blood lipid level down to nearly normal level, as described above. Examples of the substance include a substance which suppresses the production level of glycosphingolipid expressed in primary culture cells obtained from adipocytes, hepatocytes, vascular endothelial cells, vascular smooth muscle cells, skeletal muscle cells, epithelial cells, neurocytes, fibroblasts, monocytes, macrophages, etc., and various cell lines (including, but not limited to, mouse 3T3-L1 in adipocytes; human HepG2 in hepatocytes; in endothelial cells, immortalized endothelial cell lines having an acetylated LDL uptake activity; RAW264.7 in macrophages; etc.), a substance which suppresses the production level of ganglioside expressed in these cells, a substance which suppresses the production level of GM3 expressed in these cells, GM3 synthase inhibitors, a substance which suppresses the expression of GM3 synthase, etc. The pharmaceutical composition comprising these substances having the blood lipid level lowering activity is prepared as described above, and can be administered to the subject in need thereof. In this case, dose, frequency of dosing, etc. may be appropriately controlled by a physician with considering age, medical history, drugs currently used, etc. of the subject. Thus, the prevention or treatment of hyperlipidemia, etc. or control of a lipid level in blood in the subject can be achieved.

EXAMPLES

[0054] Hereinafter, the present invention is described more specifically by referring to EXAMPLES. These descriptions are presented only for illustrative and should not be construed as limiting the present invention to these EXAMPLES.

1. Generating SAT-I KO Mice

[0055] The schematic diagram of a targeting vector used for generation of the SAT-I KO mice and confirmation of genetic defect in the KO mice are shown in FIG 2A and FIG 2B, respectively. In FIG 2A, the targeting vector is shown at the top, a wild-type gene at the second row, and a variant gene at the third row. Furthermore, FIG 2B shows determination of the SAT-I genotype by PCR. For identification of the wild-type SAT-I genomic allele (exon 2), 5'-GGAATC-CATCCCTTTTCTCACAGAG-3' (SEQ ID NO: 5) and 5'-TGAAGTCACTTGGCATTGCTGG-3' (SEQ ID NO: 6) were used as primers. To confirm the SAT-I knockout, the neomycin-resistant gene inserted in the genome and the primers of 5'-GGAATCCATCCCTTTTCTCACAGAG-3' (SEQ ID NO: 7) and 5'-TGAAGTCACTTGGCATTGCTGG-3' (SEQ ID NO: 8) were used. FIG 2C shows the results of TLC analysis of brain ganglioside. In the wild-type mice (+/+) and the heterozygous mice (+/-), GM1, GD1a, GD1b, GT1b, etc. were expressed, while in the knockout mice (-/-), these all disappeared, and GM1b and GD1a were expressed in a compensatory manner (see FIG 1).

2. Generating SAT-I/apoE Double Knockout Mice

[0056] ApoE is one of the major apolipoproteins and a constituent of lipoproteins including chylomicrons (CM), very low density lipoproteins (VLDL), intermediate density lipoproteins (IDL), low density lipoproteins (LDL), etc., and involved

in transport of cholesterol or triglycerides. ApoE is produced mainly in the liver and functions as a ligand for the lipoprotein when the lipoprotein binds to LDL receptors, VLDL receptors, LDL-receptor related protein (LRP) receptors, etc. to transport extracellular lipids into the cells.

[0057] SAT-I/apoE double knockout mice were generated by breeding SAT-I knockout mice generated independently (Yoshikawa M, Go S, Takasaki K, Kakazu Y, Ohashi M, Nagafuku M, Kabayama K, Sekimoto J, Suzuki S, Takaiwa K, et al. (2009) Proc. Natl. Acad. Sci. U S A 106, 9483-9488. Epub 2009 May 9422) (FIG 2) with spontaneously apoE deficient-hyperlipidemic model mice (hereafter, apoE deficient mice) established by Matsushima et al. (Matsushima Y, Hayashi S., & Tachibana M. (1999) Mamm. Genome 10, 352-357; Matsushima Y, Sakurai T., Ohoka A., Ohnuki T., Tada N., Asoh Y, & Tachibana M. (2001) J. Atheroscler. Thromb. 8, 71-79) (FIG 3). More specifically, the animal was prepared as follows.

[0058] The mice with either SAT-I or apoE deficient (F0 generation) were bred with each other to produce the F1 generation mice heterozygous for both genes. Subsequently, the F1 generation mice were bred with each other to produce SAT-I/apoE double knockout mice in the F2 generation (FIG 3A).

[0059] In the resulting SAT-I/apoE double knockout mice, it was confirmed by western blotting if apoE protein was expressed. In FIG 3B, the band at the position marked with the arrow shows apoE protein, lane C1 designates normal mouse and lane C2 designates a control sample for the apoE single knockout mouse.

[0060] The expression of SAT-I gene was also confirmed by PCR. Specific procedures are the same as in FIG 2B. In FIG 3C, lane C3 designates the control sample for the SAT-I single knockout mouse. Ex2 marked with the arrow designates normal SAT-I gene exon 2, and Neo marked with the arrow designates the neomycin resistant gene.

3. Assay for Body Weight and Blood Lipid Level in SAT-I/apoE Double Knockout Mice

[0061] Wild-type, SAT-I single knockout, apoE single knockout and SAT-I/apoE double knockout mice (male mice at 16-18 weeks old, all fed with normal diet) were examined for the body weight and blood lipid level. Specifically, after the wild-type, SAT-I single knockout, apoE single knockout and SAT-I/apoE double knockout mice (male mice at 16-18 weeks old) were fasted for 16 hours, the body weight was measured and blood was then collected under heparin. Lipids were assayed in plasma. As a result of the analysis, there was no significant difference in the body weight among all strains of mice (FIG 4A). The cholesterol level and TG level in plasma were 489 mg/dl and 89 mg/dl, respectively, for the apoE single knockout mice, as already reported, indicating markedly high levels when compared with those of wild-type mice (cholesterol level: 63 mg/dl, TG level: 72 mg/dl) (FIGs. 4A and C). On the other hand, the cholesterol and TG levels in plasma for the SAT-I/apoE double knockout mice were 284 mg/dl and 51 mg/dl, respectively, indicating remarkable reduction as compared with those of the apoE single knockout mice (FIGs. 4B and C). In particular, the TG level was normalized to the same level as in the wild-type. As such, it was revealed that abnormal cholesterol and TG levels in hyperlipidemic model mice were improved to normal levels by knocking out SAT-I. In addition, there was no significant change in free fatty acid level in blood among all strains of mice (FIG 4D).

[0062] The lipoproteins were isolated from the sera of these mice and the profiles were compared. The serum lipoproteins isolated were fractionated by particle size using gel filtration high performance chromatography (which was performed by the method described in Usui, S. et al., J. Lipid Res. 43,805-814, 2002), and the cholesterol and triglyceride in each fraction were quantified. The results are shown in the waveform graph (FIG 5) and TABLE 1 below. In the SAT-I/apoE double knockout (DKO) mice, a decrease in plasma total cholesterol and triglyceride was shown, as compared to the apoE knockout (apoE KO) mice, and the decrease was found to be mainly due to a decrease in lipoproteins of CM, VLDL and LDL.

[TABLE 1] Comparison of cholesterol and TG contents in serum lipoprotein fractions between apoE knockout mice and DKO mice

Genotype	Cholesterol (mg/dL)				
	total**	CM***	VLDL ***	LDL**	HDL*
apoE KO (n=3)	587±0.75	8.00±0.75	434±50	131.8±1.06	11.8±2.04
DKO (n=3)	259±52.9	2.59±0.76	168±64	78.3±8.05	10.0±0.51
%DKO/apoE KO	44.1%	32.4%	38.7%	59.4%	84.7%

(continued)

Genotype	TG (mg/dL)				
	total***	CM***	VLDL ***	LDL **	HDL *
apoE KO (n=3)	59.04	19.28	31.05	6.72	1.88
DKO (n=3)	34.65	12.13	15.98	5.27	1.28
%DKO/apoE KO	58.1%	62.9%	51.5%	78.2%	68.0%

Significance of the DKO value as compared to each value of apoE KO is indicated by *: p<0.05, **: p<0.01 and ***: p<0.00.

[0063] Next, the lipid level in the liver was examined. The liver was dissected from each mouse and PBS was added thereto. The liver was minced with scissors and then homogenized with a Polytron Homogenizer 3100. After the homogenized suspension was transferred to a centrifuging tube, chloroform/methanol (1:2, v/v) was added to the homogenate and incubated at 40°C to extract the lipids. Subsequently, the extract was centrifuged and the supernatant was recovered. After chloroform/methanol (1:2, v/v) was added to the centrifuge precipitate, incubation was performed in a similar manner followed by centrifugation. The supernatant was recovered and hexane was further added to the centrifuge precipitate followed by incubation at 40°C. After centrifugation, the supernatant was recovered. All of the supernatants were pooled to give the total lipid extract. The extract was then evaporated to dryness under nitrogen. To further purify the lipid components from the total lipid extract, the extract was evaporated to dryness under nitrogen and then CHCl₃ was added, followed by ultrasonication. Purified water was added to mix them and the mixture was centrifuged. After centrifugation, the mixture was separated into the upper layer (aqueous layer), intermediate layer (protein layer) and lower layer (chloroform layer). Only the lower layer was taken out of the three layers and evaporated to dryness under nitrogen. The total lipid extract purified was dissolved in an aqueous solution containing 2% Triton X-100 and 2% sodium cholate. The cholesterol level and the triglyceride level were then assayed using a Cholesterol E-Test Wako (Wako Pure Chemical Industries, Ltd) and a TG-EN Kainos (Kainos Laboratories), respectively (FIG 6). A significant increase both in cholesterol and triglyceride levels was observed in the apoE knockout mice, as compared to the wild-type mice (FIG 6; *: p<0.05, **: p<0.01, ***: p<0.001), whereas in the SAT-I/apoE double knockout mice, no significant difference was observed when compared with the apoE knockout mice (FIG 6; designated by n.s.).

[0064] Next, the expression of glycosphingolipids was analyzed in the plasma and liver of apoE knockout mice, SAT-I/apoE double knockout mice and wild-type mice. The method for analysis of glycosphingolipids is described below. To 1 ml of mouse plasma was added 8 mL of chloroform/methanol (1:1, v/v). The resulting mixture was ultrasonicated in a water bath with an ultrasonic generator, incubated at 40°C and centrifuged to recover the supernatant. To the centrifuge precipitate was again added 8 mL of chloroform/methanol (1:2, v/v). The mixture was ultrasonicated, incubated at 40°C and centrifuged to recover the supernatant. The supernatant was pooled with the supernatant from the previous centrifugation. The pooled supernatant was then evaporated to dryness under nitrogen to give the total lipid extract. To separate the total lipid extract into neutral lipids and acidic lipids, a DEAE column obtained by packing a glass filter-equipped column (φ10mm) with DEAE-Sephadex A-25 (GE Healthcare) was used. The total lipid extract was dissolved in chloroform/methanol/purified water (30:60:8, v/v). The solution was charged onto the DEAE column. Then, chloroform/methanol/purified water (30:60:8, v/v) was passed through the column to elute the neutral lipids. Subsequently, chloroform/methanol/1N sodium acetate aqueous solution (30:60:8, v/v) was passed through the column to elute the acidic lipids. The respective eluates were evaporated to dryness under nitrogen. Next, alkaline methanolysis was performed to degrade the glycerol skeleton fats contained in each lipid fraction. After addition of 0.1 M sodium hydroxide/methanol solution, the mixture was ultrasonicated and incubated at 40°C. Thereafter, 1N hydrogen chloride/methanol solution was added to neutralize the solution. Subsequently, 50 mM sodium chloride aqueous solution was added thereto. The resulting mixture was passed through a Sep-Pak C18 Cartridge (Waters Associates, Inc.), and then purified water was passed for desalting. The first round of lipids was eluted with methanol. Furthermore, the second round of elution was performed using chloroform/methanol (2:1, v/v) for the neutral lipid fraction and using chloroform/methanol (1:2, v/v) for the acidic lipid fraction. After elution, the eluate was evaporated to dryness and the residue was dissolved in a small amount of chloroform/methanol. The solution was spotted onto a silica gel plate for TLC (Merck & Co., Inc.). Separation of the lipids was performed as follows. A first development was carried out with chloroform alone. After drying the plate, a second development for the acidic lipid fraction was carried out with chloroform/methanol/0.2% calcium chloride aqueous solution (55:45:10, v/v). For the neutral lipid fraction, a second development was performed with chloroform/methanol/purified water (60:40:10, v/v). Following the development, orcinol-sulfuric reagent (coloring reagent for compounds with a sugar chain structure) was sprayed and heated to approximately 120°C to produce a color. The bands detected were quantified with a network scanner and an image processing software (Image J).

[0065] As shown in FIG 7, the major glycosphingolipid in plasma was GM2 in the wild-type mice, and GM2, GlcCer and LacCer were increased in the neutral fraction of the apoE knockout mice. On the other hand, GM2 was detected also in the acidic fraction of SAT-I knockout mice, even though small, as compared to the wild-type mice. Also in the neutral fraction of SAT-I knockout mice, compensatory increase of LacCer and GA2 was observed. The same expression patterns as in the SAT-I knockout mice were found also in the acidic fraction and neutral fraction of SAT-I/apoE double knockout mice.

[0066] Next, the analysis results of glycosphingolipids in the liver are shown in FIG 8. It was confirmed that the major glycosphingolipid in the liver was GM2 in the wild-type mice. On the other hand, GM2 was detected also in the acidic fraction of SAT-I knockout mice, even though small, as compared to the wild-type mice. Also in the neutral fraction of SAT-I knockout mice, compensatory increase of LacCer and GA2 was observed. Further in the apoE knockout mice and the SAT-I/apoE double knockout mice, the same expression patterns as in the wild-type and the SAT-I knockout mice were found, respectively; there was no difference in expression level between the two groups.

[0067] The analysis results of gangliosides in the heart, liver and muscle of the SAT-I knockout mice and wild-type mice are given in FIG 9. The expression of GM2 was observed only in the liver. In the SAT-I knockout mice employed in this study, exon 2 of the SAT-I gene and its flanking regions are replaced with the neomycin-resistant gene (FIG 2). As a result of the gene analysis of SAT-I gene in the liver, the possibility was suggested that SAT-I variants, which do not require exon 2, might be specifically expressed in the liver. It was further confirmed, in the other SAT-I knockout mouse lacking exon 6 which encodes the enzymatically catalytic domain in SAT-I, that GM2 was not expressed in the liver, which also indicates that SAT-I is the only one enzyme for the synthesis of GM3. From the foregoing, GM2 observed in the plasma and liver of SAT-I knockout mice is considered to be derived from the liver.

[0068] The foregoing results are summarized as follows.

1) The plasma total cholesterol and triglyceride levels were reduced in the SAT-I/apoE double knockout mice down to about a half the level of the apoE knockout mice. The decrease was due to the CM, VLDL and LDL fractions.

2) With regard to the glycosphingolipid composition in the plasma and liver, the ganglioside level was decreased by the SAT-I gene defect, and the neutral glycosphingolipids including GlcCer, LacCer, GA2, GA1, etc. were increased in a compensatory manner.

3) In the liver where endogenous cholesterol and triglycerides are synthesized and VLDL that transports them is synthesized, it was examined if there is a difference in the cholesterol level and the triglyceride level. As a result, no significant difference was seen between the SAT-I/apoE double knockout mice and the apoE knockout mice.

[0069] Based on 1) and 2), the possibility was suggested that the synthesis, metabolism and excretion of cholesterol and triglycerides as well as the synthesis, release and tissue-uptaking of VLDL, etc. might be affected in the plasma or liver by the decreased gangliosides or the increased glycosphingolipids in a compensatory manner. Based on 3), even if there was no difference between cholesterol and triglyceride levels in the liver, differences were observed in their synthesis, metabolism, excretion, etc. Thus, these results in total suggested that the increase of plasma total cholesterol and plasma triglycerides is prevented.

[0070] To further confirm the results, the SAT-I knockout mice were fed with high cholesterol diet, and the cholesterol and triglyceride levels in the sera were compared with those of the wild-type mice.

[0071] After the wild-type mice and the SAT-I KO mice were fed with high cholesterol diet for 10 weeks, the sera were collected therefrom. The total cholesterol and free fatty acids were quantitatively determined using a kit available from Wako Pure Chemicals. Statistical analysis of the total cholesterol was performed using Scheffe's F test, and statistical analysis of free fatty acids was according the Tukey-Kramer method. The results indicate that the increase in serum cholesterol level observed when the wild-type mice were fed with high cholesterol diet was not observed at all when the SAT-I KO mice were fed with high cholesterol diet (FIG 10).

[0072] The foregoing results reveal, for the first time, that hyperlipidemia could be treated by suppressing the biosynthesis of gangliosides through regulation of the SAT-I gene expression (not the regulation of all GSLs expressions).

INDUSTRIAL APPLICABILITY

[0073] The present invention is a novel and revolutionary invention, in which blood lipid level can be reduced to nearly normal level through the selective regulation of ganglioside GM3. Based on the present invention, it is expected to develop a novel method for preventing/treating hyperlipidemia and furthermore, a novel method for preventing/treating atherosclerosis, hyper LDLemia, hypo HDLemia, hypercholesteremia, hypertriglyceridemia and familial hypercholesteremia.

[Sequence Listing]

SEQUENCE LISTING

5 **[0074]**

<110> Japan Science and Technology Agency

10 <120> Screening method for substance reducing lipid level in blood

<130> PCT10-0059

<150> JP 2009-269593

15 <151> 2009-11-27

<160> 8

<170> PatentIn version 3.5

20 <210> 1

<211> 1188

<212> DNA

<213> Homo sapiens

25 <220>

<221> misc_feature

<222> (1)..(1188)

<223> Human SAT-I

30 <220>

<221> CDS

<222> (1)..(1188)

35 <400> 1

40

45

50

55

EP 2 505 660 B1

	atg gct tct gtt cca atg cca agt gag tac acc tat gtg aaa ctg aga	48
	Met Ala Ser Val Pro Met Pro Ser Glu Tyr Thr Tyr Val Lys Leu Arg	
	1 5 10 15	
5	agt gat tgc tcg agg cct tcc ctg caa tgg tac acc cga gct caa agc	96
	Ser Asp Cys Ser Arg Pro Ser Leu Gln Trp Tyr Thr Arg Ala Gln Ser	
	20 25 30	
10	aag atg aga agg ccc agc ttg tta tta aaa gac atc ctc aaa tgt aca	144
	Lys Met Arg Arg Pro Ser Leu Leu Leu Lys Asp Ile Leu Lys Cys Thr	
	35 40 45	
15	ttg ctt gtg ttt gga gtg tgg atc ctt tat atc ctc aag tta aat tat	192
	Leu Leu Val Phe Gly Val Trp Ile Leu Tyr Ile Leu Lys Leu Asn Tyr	
	50 55 60	
20	act act gaa gaä tgt gac atg aaa aaa atg cat tat gtg gac cct gac	240
	Thr Thr Glu Glu Cys Asp Met Lys Lys Met His Tyr Val Asp Pro Asp	
	65 70 75 80	
25	cat gta aag aga gct cag aaa tat gct cag caa gtc ttg cag aag gaa	288
	His Val Lys Arg Ala Gln Lys Tyr Ala Gln Gln Val Leu Gln Lys Glu	
	85 90 95	
30	tgt cgt ccc aag ttt gcc aag aca tca atg gcg ctg tta ttt gag cac	336
	Cys Arg Pro Lys Phe Ala Lys Thr Ser Met Ala Leu Leu Phe Glu His	
	100 105 110	
35	agg tat agc gtg gac tta ctc cct ttt gtg cag aag gcc ccc aaa gac	384
	Arg Tyr Ser Val Asp Leu Leu Pro Phe Val Gln Lys Ala Pro Lys Asp	
	115 120 125	
40	agt gaa gct gag tcc aag tac gat cct cct ttt ggg ttc cgg aag ttc	432
	Ser Glu Ala Glu Ser Lys Tyr Asp Pro Pro Phe Gly Phe Arg Lys Phe	
	130 135 140 145 150	

EP 2 505 660 B1

	130		135		140														
	tcc	agt	aaa	gtc	cag	acc	ctc	ttg	gaa	ctc	ttg	cca	gag	cac	gac	ctc			480
5	Ser	Ser	Lys	Val	Gln	Thr	Leu	Leu	Glu	Leu	Leu	Pro	Glu	His	Asp	Leu			
	145					150					155					160			
	cct	gaa	cac	ttg	aaa	gcc	aag	acc	tgt	cgg	cgc	tgt	gtg	gtt	att	gga			528
	Pro	Glu	His	Leu	Lys	Ala	Lys	Thr	Cys	Arg	Arg	Cys	Val	Val	Ile	Gly			
					165					170					175				
10	agc	gga	gga	ata	ctg	cac	gga	tta	gaa	ctg	ggc	cac	acc	ctg	aac	cag			576
	Ser	Gly	Gly	Ile	Leu	His	Gly	Leu	Glu	Leu	Gly	His	Thr	Leu	Asn	Gln			
				180					185					190					
15	ttc	gat	ggt	gtg	ata	agg	tta	aac	agt	gca	cca	ggt	gag	gga	tat	tca			624
	Phe	Asp	Val	Val	Ile	Arg	Leu	Asn	Ser	Ala	Pro	Val	Glu	Gly	Tyr	Ser			
			195					200					205						
20	gaa	cat	ggt	gga	aat	aaa	act	act	ata	agg	atg	act	tat	cca	gag	ggc			672
	Glu	His	Val	Gly	Asn	Lys	Thr	Thr	Ile	Arg	Met	Thr	Tyr	Pro	Glu	Gly			
		210					215					220							
25	gca	cca	ctg	tct	gac	ctt	gaa	tat	tat	tcc	aat	gac	tta	ttt	ggt	gct			720
	Ala	Pro	Leu	Ser	Asp	Leu	Glu	Tyr	Tyr	Ser	Asn	Asp	Leu	Phe	Val	Ala			
	225					230					235				240				
30	ggt	tta	ttt	aag	agt	ggt	gat	ttc	aac	tgg	ctt	caa	gca	atg	gta	aaa			768
	Val	Leu	Phe	Lys	Ser	Val	Asp	Phe	Asn	Trp	Leu	Gln	Ala	Met	Val	Lys			
				245					250						255				
35	aag	gaa	acc	ctg	cca	ttc	tgg	gta	cga	ctc	ttc	ttt	tgg	aag	cag	gtg			816
	Lys	Glu	Thr	Leu	Pro	Phe	Trp	Val	Arg	Leu	Phe	Phe	Trp	Lys	Gln	Val			
			260						265					270					
40	gca	gaa	aaa	atc	cca	ctg	cag	cca	aaa	cat	ttc	agg	att	ttg	aat	cca			864
	Ala	Glu	Lys	Ile	Pro	Leu	Gln	Pro	Lys	His	Phe	Arg	Ile	Leu	Asn	Pro			
			275				280						285						
45	ggt	atc	atc	aaa	gag	act	gcc	ttt	gac	atc	ctt	cag	tac	tca	gag	cct			912
	Val	Ile	Ile	Lys	Glu	Thr	Ala	Phe	Asp	Ile	Leu	Gln	Tyr	Ser	Glu	Pro			
		290					295					300							
50	cag	tca	agg	ttc	tgg	ggc	cga	gat	aag	aac	gtc	ccc	aca	atc	ggt	gtc			960
	Gln	Ser	Arg	Phe	Trp	Gly	Arg	Asp	Lys	Asn	Val	Pro	Thr	Ile	Gly	Val			
	305					310					315				320				
55	att	gcc	ggt	gtc	tta	gcc	aca	cat	ctg	tgc	gat	gaa	gtc	agt	ttg	gcg			1008
	Ile	Ala	Val	Val	Leu	Ala	Thr	His	Leu	Cys	Asp	Glu	Val	Ser	Leu	Ala			
				325						330					335				
60	ggt	ttt	gga	tat	gac	ctc	aat	caa	ccc	aga	aca	cct	ttg	cac	tac	ttc			1056
	Gly	Phe	Gly	Tyr	Asp	Leu	Asn	Gln	Pro	Arg	Thr	Pro	Leu	His	Tyr	Phe			
				340					345					350					
65	gac	agt	caa	tgc	atg	gct	gct	atg	aac	ttt	cag	acc	atg	cat	aat	gtg			1104
	Asp	Ser	Gln	Cys	Met	Ala	Ala	Met	Asn	Phe	Gln	Thr	Met	His	Asn	Val			
			355					360					365						
70	aca	acg	gaa	acc	aag	ttc	ctc	tta	aag	ctg	gtc	aaa	gag	gga	gtg	gtg			1152
	Thr	Thr	Glu	Thr	Lys	Phe	Leu	Leu	Lys	Leu	Val	Lys	Glu	Gly	Val	Val			
			370				375					380							
75	aaa	gat	ctc	agt	gga	ggc	att	gat	cgt	gaa	ttt	tga							1188
	Lys	Asp	Leu	Ser	Gly	Gly	Ile	Asp	Arg	Glu	Phe								
	385					390					395								

EP 2 505 660 B1

<210> 2
 <211> 395
 <212> PRT
 <213> Homo sapiens

5

<400> 2

10	Met	Ala	Ser	Val	Pro	Met	Pro	Ser	Glu	Tyr	Thr	Tyr	Val	Lys	Leu	Arg
	1				5					10					15	
15	Ser	Asp	Cys	Ser	Arg	Pro	Ser	Leu	Gln	Trp	Tyr	Thr	Arg	Ala	Gln	Ser
				20					25					30		
20	Lys	Met	Arg	Arg	Pro	Ser	Leu	Leu	Leu	Lys	Asp	Ile	Leu	Lys	Cys	Thr
			35					40						45		
25	Leu	Leu	Val	Phe	Gly	Val	Trp	Ile	Leu	Tyr	Ile	Leu	Lys	Leu	Asn	Tyr
		50					55						60			
30	Thr	Thr	Glu	Glu	Cys	Asp	Met	Lys	Lys	Met	His	Tyr	Val	Asp	Pro	Asp
	65					70					75					80
35	His	Val	Lys	Arg	Ala	Gln	Lys	Tyr	Ala	Gln	Gln	Val	Leu	Gln	Lys	Glu
					85					90					95	
40	Cys	Arg	Pro	Lys	Phe	Ala	Lys	Thr	Ser	Met	Ala	Leu	Leu	Phe	Glu	His
				100					105					110		
45	Arg	Tyr	Ser	Val	Asp	Leu	Leu	Pro	Phe	Val	Gln	Lys	Ala	Pro	Lys	Asp
			115					120					125			
50	Ser	Glu	Ala	Glu	Ser	Lys	Tyr	Asp	Pro	Pro	Phe	Gly	Phe	Arg	Lys	Phe
		130					135					140				
55	Ser	Ser	Lys	Val	Gln	Thr	Leu	Leu	Glu	Leu	Leu	Pro	Glu	His	Asp	Leu
	145					150					155					160
60	Pro	Glu	His	Leu	Lys	Ala	Lys	Thr	Cys	Arg	Arg	Cys	Val	Val	Ile	Gly
					165					170					175	
65	Ser	Gly	Gly	Ile	Leu	His	Gly	Leu	Glu	Leu	Gly	His	Thr	Leu	Asn	Gln
				180					185						190	
70	Phe	Asp	Val	Val	Ile	Arg	Leu	Asn	Ser	Ala	Pro	Val	Glu	Gly	Tyr	Ser
			195					200					205			
75	Glu	His	Val	Gly	Asn	Lys	Thr	Thr	Ile	Arg	Met	Thr	Tyr	Pro	Glu	Gly
	210						215					220				

EP 2 505 660 B1

Ala Pro Leu Ser Asp Leu Glu Tyr Tyr Ser Asn Asp Leu Phe Val Ala
 225 230 235 240

5 Val Leu Phe Lys Ser Val Asp Phe Asn Trp Leu Gln Ala Met Val Lys
 245 250 255

10 Lys Glu Thr Leu Pro Phe Trp Val Arg Leu Phe Phe Trp Lys Gln Val
 260 265 270

15 Ala Glu Lys Ile Pro Leu Gln Pro Lys His Phe Arg Ile Leu Asn Pro
 275 280 285

20 Val Ile Ile Lys Glu Thr Ala Phe Asp Ile Leu Gln Tyr Ser Glu Pro
 290 295 300

25 Gln Ser Arg Phe Trp Gly Arg Asp Lys Asn Val Pro Thr Ile Gly Val
 305 310 315 320

Ile Ala Val Val Leu Ala Thr His Leu Cys Asp Glu Val Ser Leu Ala
 325 330 335

30 Gly Phe Gly Tyr Asp Leu Asn Gln Pro Arg Thr Pro Leu His Tyr Phe
 340 345 350

35 Asp Ser Gln Cys Met Ala Ala Met Asn Phe Gln Thr Met His Asn Val
 355 360 365

Thr Thr Glu Thr Lys Phe Leu Leu Lys Leu Val Lys Glu Gly Val Val
 370 375 380

40 Lys Asp Leu Ser Gly Gly Ile Asp Arg Glu Phe
 385 390 395

<210> 3

<211> 1245

<212> DNA

45 <213> Mus musculus

<220>

<221> misc_feature

<222> (1)..(1245)

50 <223> Mouse SAT-I

<220>

<221> CDS

<222> (1)..(1245)

55 <400> 3

EP 2 505 660 B1

atg cac aca gag gcg gtg ggc ggc gcg gcg cgg agg ccc cag aag ctg 48
Met His Thr Glu Ala Val Gly Gly Ala Ala Arg Arg Pro Gln Lys Leu
1 5 10 15

5 cga agc caa gca gcg gca cct gcc tgc cga gca atg cca agt gag ttc 96

10

15

20

25

30

35

40

45

50

55

EP 2 505 660 B1

	Arg	Ser	Gln	Ala	Ala	Ala	Pro	Ala	Cys	Arg	Ala	Met	Pro	Ser	Glu	Phe	
				20					25					30			
	acc	tct	gca	aag	ctg	aga	agt	gat	tgc	tca	agg	acc	tcc	ctg	caa	tgg	144
5	Thr	Ser	Ala	Lys	Leu	Arg	Ser	Asp	Cys	Ser	Arg	Thr	Ser	Leu	Gln	Trp	
			35					40					45				
	tac	acc	cga	acc	cag	cac	aag	atg	aga	aga	ccc	agc	ttg	tta	ata	aaa	192
	Tyr	Thr	Arg	Thr	Gln	His	Lys	Met	Arg	Arg	Pro	Ser	Leu	Leu	Ile	Lys	
10		50					55					60					
	gac	atc	tgc	aag	tgc	acg	ttg	gtt	gca	ttt	gga	gtc	tgg	ctc	ctg	tac	240
	Asp	Ile	Cys	Lys	Cys	Thr	Leu	Val	Ala	Phe	Gly	Val	Trp	Leu	Leu	Tyr	
	65					70					75					80	
	atc	ctc	att	ttg	aat	tac	acc	gct	gaa	gaa	tgt	gac	atg	aaa	aga	atg	288
15	Ile	Leu	Ile	Leu	Asn	Tyr	Thr	Ala	Glu	Glu	Cys	Asp	Met	Lys	Arg	Met	
					85					90					95		
	cac	tat	gtg	gac	cct	gac	cgg	ata	aag	aga	gct	cag	agc	tat	gct	cag	336
	His	Tyr	Val	Asp	Pro	Asp	Arg	Ile	Lys	Arg	Ala	Gln	Ser	Tyr	Ala	Gln	
20				100					105					110			
	gaa	gtc	ttg	cag	aag	gaa	tgt	cgg	ccc	agg	tac	gcg	aag	acg	gct	atg	384
	Glu	Val	Leu	Gln	Lys	Glu	Cys	Arg	Pro	Arg	Tyr	Ala	Lys	Thr	Ala	Met	
			115					120					125				
	gct	ctg	tta	ttt	gag	gac	agg	tac	agc	atc	aac	ttg	gag	cct	ttt	gtg	432
25	Ala	Leu	Leu	Phe	Glu	Asp	Arg	Tyr	Ser	Ile	Asn	Leu	Glu	Pro	Phe	Val	
		130					135					140					
	cag	aag	gtc	ccc	acg	gcc	agt	gaa	gct	gag	ctc	aag	tat	gac	ccg	cct	480
	Gln	Lys	Val	Pro	Thr	Ala	Ser	Glu	Ala	Glu	Leu	Lys	Tyr	Asp	Pro	Pro	
30						150					155					160	
	ttt	gga	ttc	cgg	aag	ttc	tcc	agt	aaa	gtc	cag	agc	ctc	ttg	gat	atg	528
	Phe	Gly	Phe	Arg	Lys	Phe	Ser	Ser	Lys	Val	Gln	Ser	Leu	Leu	Asp	Met	
					165					170					175		
	ctg	ccc	gaa	cat	gac	ttt	cct	gaa	cac	ttg	aga	gcc	aag	gcc	tgc	aag	576
35	Leu	Pro	Glu	His	Asp	Phe	Pro	Glu	His	Leu	Arg	Ala	Lys	Ala	Cys	Lys	
				180					185					190			
	cgc	tgt	gtg	gtt	gtt	ggg	aac	ggg	ggc	atc	ctg	cac	gga	cta	gag	ctg	624
	Arg	Cys	Val	Val	Val	Gly	Asn	Gly	Gly	Ile	Leu	His	Gly	Leu	Glu	Leu	
40			195					200					205				
	ggg	cac	gac	ctc	aac	cag	ttc	gat	gtg	gta	ata	agg	ttg	aac	agt	gcg	672
	Gly	His	Ala	Leu	Asn	Gln	Phe	Asp	Val	Val	Ile	Arg	Leu	Asn	Ser	Ala	
		210				215						220					
	cca	ggt	gag	ggt	tac	tct	gaa	cac	ggt	ggg	aat	aaa	act	act	ata	agg	720
45	Pro	Val	Glu	Gly	Tyr	Ser	Glu	His	Val	Gly	Asn	Lys	Thr	Thr	Ile	Arg	
						230					235					240	
	atg	act	tac	cca	gag	ggt	gcg	cca	ctg	tcg	gac	ggt	gaa	tac	tac	gcc	768
	Met	Thr	Tyr	Pro	Glu	Gly	Ala	Pro	Leu	Ser	Asp	Val	Glu	Tyr	Tyr	Ala	
50					245					250					255		
	aat	gat	ttg	ttc	gtt	act	gtt	tta	ttt	aag	agt	ggt	gat	ttc	aag	tgg	816
	Asn	Asp	Leu	Phe	Val	Thr	Val	Leu	Phe	Lys	Ser	Val	Asp	Phe	Lys	Trp	
				260				265						270			
55	ctt	caa	gca	atg	gta	aaa	aat	gaa	agc	ctg	ccc	ttt	tgg	ggt	cgc	ctc	864
	Leu	Gln	Ala	Met	Val	Lys	Asn	Glu	Ser	Leu	Pro	Phe	Trp	Val	Arg	Leu	

EP 2 505 660 B1

	275	280	285	
5	ttc ttt tgg aag caa gtg gca gaa aaa gtc cca ctc cag cca aag cac Phe Phe Trp Lys Gln Val Ala Glu Lys Val Pro Leu Gln Pro Lys His 290 295 300			912
10	ttc agg att ttg aac cca gtt atc atc aaa gaa act gcc ttc gac atc Phe Arg Ile Leu Asn Pro Val Ile Ile Lys Glu Thr Ala Phe Asp Ile 305 310 315 320			960
15	ctt cag tac tca gag cct cag tca aga ttc tgg ggc cat gat aag aac Leu Gln Tyr Ser Glu Pro Gln Ser Arg Phe Trp Gly His Asp Lys Asn 325 330 335			1008
20	atc ccc acg atc ggc gtc att gcc gtt gtc ttg gct aca cat ctg tgt Ile Pro Thr Ile Gly Val Ile Ala Val Val Leu Ala Thr His Leu Cys 340 345 350			1056
25	gat gaa gtc agc ctg gca ggc ttt ggc tac gac ctc agt caa ccc agg Asp Glu Val Ser Leu Ala Gly Phe Gly Tyr Asp Leu Ser Gln Pro Arg 355 360 365			1104
30	acc cct ctg cac tac ttt gac agt cag tgc atg ggc gcc atg cac tgg Thr Pro Leu His Tyr Phe Asp Ser Gln Cys Met Gly Ala Met His Trp 370 375 380			1152
35	cag gtc atg cac aat gtg acc aca gag acc aag ttc ctc ctg aag ctc Gln Val Met His Asn Val Thr Thr Glu Thr Lys Phe Leu Leu Lys Leu 385 390 395 400			1200
40	ctc aag gag ggc gtg gtg gag gac ctc agc ggc ggc atc cac tga Leu Lys Glu Gly Val Val Glu Asp Leu Ser Gly Gly Ile His 405 410			1245

<210> 4
 <211> 414
 <212> PRT
 <213> Mus musculus

<400> 4

EP 2 505 660 B1

Met His Thr Glu Ala Val Gly Gly Ala Ala Arg Arg Pro Gln Lys Leu
 1 5 10 15

5 Arg Ser Gln Ala Ala Ala Pro Ala Cys Arg Ala Met Pro Ser Glu Phe
 20 25 30

10 Thr Ser Ala Lys Leu Arg Ser Asp Cys Ser Arg Thr Ser Leu Gln Trp
 35 40 45

15 Tyr Thr Arg Thr Gln His Lys Met Arg Arg Pro Ser Leu Leu Ile Lys
 50 55 60

Asp Ile Cys Lys Cys Thr Leu Val Ala Phe Gly Val Trp Leu Leu Tyr
 65 70 75 80

20 Ile Leu Ile Leu Asn Tyr Thr Ala Glu Glu Cys Asp Met Lys Arg Met
 85 90 95

25

30

35

40

45

50

55

EP 2 505 660 B1

His Tyr Val Asp Pro Asp Arg Ile Lys Arg Ala Gln Ser Tyr Ala Gln
 100 105 110

5
 Glu Val Leu Gln Lys Glu Cys Arg Pro Arg Tyr Ala Lys Thr Ala Met
 115 120 125

10
 Ala Leu Leu Phe Glu Asp Arg Tyr Ser Ile Asn Leu Glu Pro Phe Val
 130 135 140

15
 Gln Lys Val Pro Thr Ala Ser Glu Ala Glu Leu Lys Tyr Asp Pro Pro
 145 150 155 160

20
 Phe Gly Phe Arg Lys Phe Ser Ser Lys Val Gln Ser Leu Leu Asp Met
 165 170 175

25
 Leu Pro Glu His Asp Phe Pro Glu His Leu Arg Ala Lys Ala Cys Lys
 180 185 190

30
 Arg Cys Val Val Val Gly Asn Gly Gly Ile Leu His Gly Leu Glu Leu
 195 200 205

35
 Gly His Ala Leu Asn Gln Phe Asp Val Val Ile Arg Leu Asn Ser Ala
 210 215 220

40
 Pro Val Glu Gly Tyr Ser Glu His Val Gly Asn Lys Thr Thr Ile Arg
 225 230 235 240

45
 Met Thr Tyr Pro Glu Gly Ala Pro Leu Ser Asp Val Glu Tyr Tyr Ala
 245 250 255

50
 Asn Asp Leu Phe Val Thr Val Leu Phe Lys Ser Val Asp Phe Lys Trp
 260 265 270

55
 Leu Gln Ala Met Val Lys Asn Glu Ser Leu Pro Phe Trp Val Arg Leu
 275 280 285

60
 Phe Phe Trp Lys Gln Val Ala Glu Lys Val Pro Leu Gln Pro Lys His
 290 295 300

65
 Phe Arg Ile Leu Asn Pro Val Ile Ile Lys Glu Thr Ala Phe Asp Ile
 305 310 315 320

70
 Leu Gln Tyr Ser Glu Pro Gln Ser Arg Phe Trp Gly His Asp Lys Asn
 325 330 335

75
 Ile Pro Thr Ile Gly Val Ile Ala Val Val Leu Ala Thr His Leu Cys
 340 345 350

EP 2 505 660 B1

Asp Glu Val Ser Leu Ala Gly Phe Gly Tyr Asp Leu Ser Gln Pro Arg
 355 360 365

5 Thr Pro Leu His Tyr Phe Asp Ser Gln Cys Met Gly Ala Met His Trp
 370 375 380

10 Gln Val Met His Asn Val Thr Thr Glu Thr Lys Phe Leu Leu Lys Leu
 385 390 395 400

Leu Lys Glu Gly Val Val Glu Asp Leu Ser Gly Gly Ile His
 405 410

15 <210> 5
 <211> 25
 <212> DNA
 <213> Artificial Sequence

20 <220>
 <223> Primer

25 <400> 5
 ggaatccatc ccttttctca cagag 25

30 <210> 6
 <211> 22
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Primer

35 <400> 6
 tgaactcact tggcattgct gg 22

40 <210> 7
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Primer

45 <400> 7
 ggaatccatc ccttttctca cagag 25

50 <210> 8
 <211> 22
 <212> DNA
 <213> Artificial Sequence

55 <220>
 <223> Primer

<400> 8
 tgaactcact tggcattgct gg 22

Claims

1. A method for screening a GM3 synthase specific inhibitor having a blood lipid level lowering activity, which comprises:

- 5 (i) contacting GM3 synthase with a substrate thereof in the presence and absence of a test substance,
(ii) selecting a test substance that decreases the synthesized amount of GM3,
(iii) administering the selected test substance to a non-human animal, and
(iv) determining a cholesterol level and/or a triglyceride level in blood in the non-human animal, and selecting
10 a test substance which reduces the cholesterol level and/or the triglyceride level by 10% or more as compared
to a control.

2. A method for screening a substance that inhibits GM3 synthase expression and has a blood lipid level lowering activity, which comprises:

- 15 (i) contacting a test substance with a cell expressing GM3 synthase,
(ii) selecting a test substance that decreases the expression level of GM3 synthase as compared to a control,
(iii) administering the selected test substance to a non-human animal, and
(iv) determining a cholesterol level and/or a triglyceride level in blood in the non-human animal, and selecting
20 a test substance which reduces the cholesterol level and/or the triglyceride level by 10% or more as compared
to a control.

3. The method according to claim 2, wherein the cell expressing GM3 synthase is selected from the group consisting
of adipocytes, hepatocytes, vascular endothelial cells, vascular smooth muscle cells, skeletal muscle cells, epithelial
25 cells, neurocytes, fibroblasts, monocytes and macrophages.

4. The method according to claim 2, wherein the cell expressing GM3 synthase is a transformant produced by genetic
engineering.

5. The screening method according to any one of claims 1 to 4, which further comprises selecting a test substance
30 that reduces a blood cholesterol level and/or a blood triglyceride level.

6. The method according to any one of claims 1 to 5, wherein the non-human animal has a higher blood lipid level
than normal level.

7. The method according to any one of claims 1 to 6, wherein the non-human animal is a mouse.

Patentansprüche

40 1. Verfahren zum Screenen eines GM3-Synthase-spezifischen Inhibitors mit einer Blut-Lipidspiegel-reduzierenden
Aktivität, welches umfasst:

- (i) Inkontaktbringen von GM3-Synthase mit einem Substrat davon in der Gegenwart und Abwesenheit einer
Testsubstanz,
45 (ii) Auswählen einer Testsubstanz, die die synthetisierte Menge von GM3 reduziert,
(iii) Verabreichen der ausgewählten Testsubstanz an ein nicht-humanes Tier, und
(iv) Bestimmen eines Cholesterolspiegels und/oder Triglyceridspiegels in Blut in dem nicht-humanen Tier, und
Auswählen einer Testsubstanz, die den Cholesterolspiegel und/oder den Triglyceridspiegel um 10% oder mehr
50 im Vergleich zu einer Kontrolle reduziert.

2. Verfahren zum Screenen einer Substanz, die GM3-Synthase-Expression inhibiert und eine Blut-Lipidspiegel-ver-
ringende Aktivität aufweist, welches umfasst:

- (i) Inkontaktbringen einer Testsubstanz mit einer GM3-Synthase exprimierenden Zelle,
55 (ii) Auswählen einer Testsubstanz, die den Expressionsspiegel von GM3-Synthase im Vergleich zu einer Kon-
trolle reduziert,
(iii) Verabreichen der ausgewählten Testsubstanz an ein nicht-humanes Tier, und
(iv) Bestimmen eines Cholesterolspiegels und/oder eines Triglyceridspiegels in Blut in dem nicht-humanen Tier,

EP 2 505 660 B1

und Auswählen einer Testsubstanz, die den Cholesterolspiegel und/oder den Triglyceridspiegel um 10% oder mehr im Vergleich zu einer Kontrolle reduziert.

- 5 3. Verfahren nach Anspruch 2, wobei die GM3-Synthase exprimierende Zelle ausgewählt ist aus der Gruppe, bestehend aus Adipocyten, Hepatocyten, vaskularen Endothelzellen, vaskularen glatten Muskelzellen, Skelettmuskelzellen, Epithelzellen, Neurozyten, Fibroblasten, Monozyten und Makrophagen.
- 10 4. Verfahren nach Anspruch 2, wobei die GM3-Synthase exprimierende Zelle eine durch genetische Manipulation hergestellte Transformante ist.
- 15 5. Screeningverfahren nach einem der Ansprüche 1 bis 4, die weiterhin Auswählen einer Testsubstanz umfasst, die einen Blut-Cholesterolspiegel und/oder einen Blut-Triglyceridspiegel reduziert.
6. Verfahren nach einem der Ansprüche 1 bis 5, wobei das nicht-humane Tier einen höheren Blut-Lipidspiegel als normalen Spiegel aufweist.
7. Verfahren nach einem der Ansprüche 1 bis 6, wobei das nicht-humane Tier eine Maus ist.

20 Revendications

- 25 1. Méthode de criblage d'un inhibiteur spécifique de GM3 synthase ayant une activité d'abaissement du taux de lipides dans le sang, qui comprend :
 - (i) la mise en contact de GM3 synthase avec un substrat de celle-ci en présence et absence d'une substance d'essai,
 - (ii) la sélection d'une substance d'essai qui diminue la quantité synthétisée de GM3,
 - (iii) l'administration de la substance d'essai sélectionnée à un animal non humain, et
 - 30 (iv) la détermination d'un taux de cholestérol et/ou d'un taux de triglycéride dans le sang chez l'animal non humain, et la sélection d'une substance d'essai qui réduit le taux de cholestérol et/ou le taux de triglycéride de 10 % ou plus comparé à un témoin.
- 35 2. Méthode de criblage d'une substance qui inhibe l'expression de GM3 synthase et a une activité d'abaissement du taux de lipides dans le sang, qui comprend :
 - (i) la mise en contact d'une substance d'essai avec une cellule exprimant la GM3 synthase,
 - (ii) la sélection d'une substance d'essai qui diminue le taux d'expression de la GM3 synthase comparé à un témoin,
 - (iii) l'administration de la substance d'essai sélectionnée à un animal non humain, et
 - 40 (iv) la détermination d'un taux de cholestérol et/ou d'un taux de triglycéride dans le sang chez l'animal non humain, et la sélection d'une substance d'essai qui réduit le taux de cholestérol et/ou le taux de triglycéride de 10 % ou plus comparé à un témoin.
- 45 3. Méthode selon la revendication 2, dans laquelle la cellule exprimant la GM3 synthase est choisie dans le groupe consistant en les adipocytes, les hépatocytes, les cellules endothéliales vasculaires, les cellules du muscle lisse vasculaire, les cellules du muscle squelettique, les cellules épithéliales, les neurocytes, les fibroblastes, les mono-cytes et les macrophages.
- 50 4. Méthode selon la revendication 2, dans laquelle la cellule exprimant la GM3 synthase est une cellule mutante produite par génie génétique.
5. Méthode de criblage selon l'une quelconque des revendications 1 à 4, comprenant en outre la sélection d'une substance d'essai qui réduit un taux de cholestérol dans le sang et/ou un taux de triglycéride dans le sang.
- 55 6. Méthode de criblage selon l'une quelconque des revendications 1 à 5, dans laquelle l'animal non humain a un taux de lipides dans le sang plus élevé que le taux normal.
7. Méthode de criblage selon l'une quelconque des revendications 1 à 6, dans laquelle l'animal non humain est une

souris.

5

10

15

20

25

30

35

40

45

50

55




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7





FIG. 8






FIG. 10

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- EP 1726313 A [0006]
- WO 02084301 A [0007]
- JP 2002516062 PCT [0035]
- US 2002086356A A [0035]
- JP 2009269593 A [0074]

Non-patent literature cited in the description

- **INOKUCHI J. ; RADIN N.** *J. Lipid Res.*, 1987, vol. 28, 565-571 [0005]
- **TAGAMI S. ; INOKUCHI J. ; KABAYAMA K. ; YOSHIMURA H. ; KITAMURA F. ; UEMURA S. ; OGAWA C. ; ISHII A. ; SAITO M. ; OHTSUKA Y et al.** *J. Biol. Chem.*, 2002, vol. 277, 3085-3092 [0005]
- **ZHAO H. ; PRZYBYLSKA M. ; WU I.H. ; ZHANG J. ; SIEGEL C. ; KOMAMITSKY S. ; YEW N.S. ; CHENG S.H.** *Diabetes*, 2007, vol. 56, 1210-1218 [0005]
- **ZHAO H. ; PRZYBYLSKA M. ; WU I.H. ; ZHANG J. ; MANIATIS P. ; PACHECO J. ; PIEPENHAGEN P. ; COPELAND D. ; ARBEENY C. ; SHAYMAN J.A. et al.** *Hepatology*, 2009, vol. 50, 85-93 [0005]
- **AERTS J. M. ; OTTENHOFF R. ; POWLSON A.S. ; GREFFHORST A. ; VAN EIJK M. ; DUBBELHUIS P.F. ; ATEN J. ; KUIPERS F. ; SERLIE M. J. ; WENNEKES T. et al.** *Diabetes*, 2007, vol. 56, 1341-1349 [0005]
- **VAN EIJK M. ; ATEN J. ; BIJL N. ; OTTENHOFF R. ; VAN ROOMEN C.P. ; DUBBELHUIS P.F. ; SEEMAN I. ; GHAUHARALI-VAN DER VLUGT K. ; OVERKLEEFTH.S. ; ARBEENY C. et al.** *PLoS One*, March 2009, vol. 4, 4723 [0005]
- **BIJL N. ; VAN ROOMEN C.P. ; TRIANTIS V. ; SOKOLOVIC M. ; OTTENHOFF R. ; SCHEIJ S. ; VAN EIJK M. ; BOOT R.G. ; AERTS J.M. ; GROEN A.K.** *Hepatology*, 2009, vol. 49, 637-645 [0005]
- **YAMASHITA, T. et al.** *Proc. Natl. Acad. Sci. USA*, 1999, vol. 96, 9142-9147 [0005]
- **YAMASHITA T. ; HASHIRAMOTO A. ; HALUZIK M. ; MIZUKAMI H. ; BECK S. ; NORTON A. ; KONO M. ; TSUJI S. ; DANIOTTI J.L. ; WERTH N. et al.** *Proc. Natl. Acad. Sci. U S A*, 2003, vol. 100, 3445-3449 [0005]
- **KISO SEIKAGAKU JIKKENHO.** Shishitsu, Toshitsu, Fukugotoshitsu. vol. 5 [0016] [0017] [0018]
- **SHIN SEIKAGAKU JIKKEN KOZA. MARIKO SHIBA et al.** Shishitsu I: Chusei Shishitsu-to-III Gata Koshiketsusho. Tokyo Kagaku Dojin Publishing Co, 1993, vol. 4 [0020]
- Receptor Modifications in Glycobiology. **INOKUCHI J. ; KABAYAMA K.** Comprehensive Glycoscience. Elsevier Science & Technology, 2007, vol. 3, 733-744 [0025]
- **GLAROS EN. et al.** Glycosphingolipid Accumulation Inhibits Cholesterol Efflux via the ABCA1/Apolipoprotein A-I Pathway. *J. Biol. Chem.*, 2005, vol. 280, 24515-24523 [0025]
- **KABAYAMA K. et al.** *Glycobiology*, 2005, vol. 15, 21-29 [0025]
- **KABAYAMA K. et al.** *Proc. Natl. Acad. Sci. U S A*, 2007, vol. 104, 13678-13683 [0025]
- **KOTANI, M. et al.** *Biochem. Biophys. Acta*, 1992, vol. 1117, 97-103 [0026]
- **SCATCHARD.** *Ann. N.Y. Acad. Sci.*, 1949, vol. 51, 660-672 [0026]
- **HIRASHIMA ; INOUE.** Shin Seikagaku Jikken Koza 2, Kakusan IV, Idenshi No Fukusei To Hatsugen. Tokyo Kagaku Dojin Co., Ltd, 1993, 319-347 [0032]
- **STEIN.** *Nucleic Acids Res.*, 1988, vol. 16, 3209-3221 [0032]
- *FEBS Lett.*, 1988, vol. 228, 228 [0033]
- *FEBS Lett.*, 1988, vol. 239, 285 [0033]
- *Nucl. Acids Res.*, 1989, vol. 17, 7059 [0033]
- *Nature*, 1986, vol. 323, 349 [0033]
- *Nucl. Acids Res.*, 1991, vol. 19, 6751 [0033]
- *Protein Eng.*, 1990, vol. 3, 733 [0033]
- *Nucl. Acids Res.*, 1991, vol. 19, 3875 [0033]
- *Nucl. Acids Res.*, 1991, vol. 19, 5125 [0033]
- *Biochem. Biophys. Res. Commun.*, 1992, vol. 186, 1271 [0033]
- *Nature Genetics*, February 2000, vol. 24 (2), 180-183 [0035]
- *Genesis*, April 2000, vol. 26 (4), 240-244 [0035]
- *Nature*, 21 September 2002, vol. 407 (6802), 319-20 [0035]
- *Genes & Dev.*, 15 April 2002, vol. 16 (8), 948-958 [0035]
- *Proc. Natl. Acad. Sci. USA.*, 16 April 2002, vol. 99 (8), 5515-5520 [0035]
- *Science*, 19 April 2002, vol. 296 (5567), 550-553 [0035]

EP 2 505 660 B1

- *Proc Natl. Acad. Sci. USA*, 30 April 2002, vol. 99 (9), 6047-6052 [0035]
- *Nature Biotechnology*, May 2002, vol. 20 (5), 497-500 [0035]
- *Nature Biotechnology*, May 2002, vol. 20 (5), 500-505 [0035]
- *Nucleic Acids Res.*, 15 May 2002, vol. 30 (10), 46 [0035]
- **J. SAMBROOK et al.** *Molecular Cloning*. Cold Spring Harbour Laboratory Press, 2001 [0037] [0038]
- *Current Protocols in Molecular Biology*. John Wiley & Sons, 1987 [0037] [0038]
- **MACHER, B. A. ; KLOCK, J. C.** *J. Biol. Chem.*, 1980, vol. 255, 2092-2096 [0039]
- **LEDEEN et al.** *J. Neurochem.*, 1973, vol. 21, 829-839 [0039]
- **KIM J-W. et al.** *Gene*, 2001, vol. 273, 163-171 [0043]
- **KIM S-W. et al.** *Biochim. Biophys. Acta*, 2002, vol. 1578, 84-89 [0043]
- **CHOI H-J. et al.** *Biochem. Biophys. Res. Commun.*, 2004, vol. 313, 142-147 [0043]
- **YOSHIKAWA M ; GO S ; TAKASAKI K ; KAKAZU Y ; OHASHI M ; NAGAFUKU M ; KABAYAMA K ; SEKIMOTO J ; SUZUKI S ; TAKAIWA K et al.** *Proc. Natl. Acad. Sci. USA*, May 2009, vol. 106, 9483-9488 [0057]
- **MATSUSHIMA Y ; HAYASHI S. ; TACHIBANA M.** *Mamm. Genome*, 1999, vol. 10, 352-357 [0057]
- **MATSUSHIMA Y ; SAKURAI T. ; OHOKA A. ; OHNUKI T. ; TADA N. ; ASOH Y ; TACHIBANA M.** *J. Atheroscler. Thromb.*, 2001, vol. 8, 71-79 [0057]
- **USUI, S. et al.** *J. Lipid Res.*, 2002, vol. 43, 805-814 [0062]